



OPEN ACCESS

Original research

Repetitive and forceful movements of the hand as predictors of treatment for pain in the distal upper extremities

Jonathan Aavang Petersen ¹, Charlotte Brauer,¹ Lau Caspar Thygesen,² Esben Meulengracht Flachs,¹ Christina Bach Lund,¹ Jane Frølund Thomsen¹

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/oemed-2021-107543>).

¹Department of Occupational and Environmental Medicine, Bispebjerg Hospital, Copenhagen, Denmark
²National Institute of Public Health, University of Southern Denmark, Odense, Denmark

Correspondence to

Dr Jonathan Aavang Petersen, Department of Occupational and Environmental Medicine, Bispebjerg Hospital, 2400 København, Denmark; jonathanaavang@hotmail.com

Received 11 March 2021
Accepted 2 August 2021
Published Online First
20 August 2021

ABSTRACT

Objective To investigate repetitive movements and the use of hand force as causes of treatment for distal upper extremities musculoskeletal disorders

Methods A cohort of 202 747 workers in a pension health scheme from 2005 to 2017 in one of 17 jobs (eg, office work, carpentry, cleaning) was formed. Representative electro-goniometric measurements of wrist angular velocity as a measure for repetition and expert-rated use of hand force were used in a job exposure matrix (JEM). Job titles were retrieved from the Danish registers. Outcome was first treatment in the distal upper extremities. In a Poisson regression model, incidence rate ratios (IRRs) of treatment were adjusted for age, calendar-year, diagnosis of rheumatoid arthritis and arm fractures. In further analyses, wrist velocity or hand force was added.

Results In men, wrist velocity had an IRR of 1.48 (95% CI 1.15 to 1.91) when the highest exposure level was compared with the lowest but with no clear exposure-response pattern. The effect became insignificant when adjusted for hand force. Hand force had an IRR of 2.65 (95% CI 2.13 to 3.29) for the highest versus the lowest exposure with an exposure-response pattern, which remained after adjustment for wrist velocity. Among women, no increased risk was found for hand force, while wrist velocity showed a significantly protective association with treatment.

Conclusions In men, occupational exposure to hand force more than doubled the risk of seeking treatment. The results for exposure to repetition were less clear. In women, we could not find any indications of an increased risk neither for force nor for repetition.

INTRODUCTION

Distal upper extremity musculoskeletal disorders (D-UEMSD) are highly prevalent among the working population with epicondylitis of the elbow and tenosynovitis of the hand or wrist as the most commonly recognised occupational diseases in the European Union.¹ Musculoskeletal disorders are a leading cause of work disability and loss of productivity across Europe.² Incidence of epicondylitis and hand or wrist tendinitis ranges from 0.45 to 7.0 per 100 workers depending on case definition and population.³ The most consistent non-occupational risk factors include age, female sex, obesity, psychosocial factors, previous injuries and inflammatory arthritis.^{4 5}

Key messages

What is already known about this subject?

► Some evidence exists of an association between measures of hand force and repetition and tendinitis of the wrist and epicondylitis of the elbow; however, these studies have often been limited by bias because of self-reported exposure and an outcome sensitive to time fluctuations.

What are the new findings?

► We found a clear exposure-response association between expert-rated use of hand force in 17 different jobs and treatment for pain in the arms and hands in a private health scheme among men.
► An effect of high repetition using technical measurements of wrist angular velocity was less evident and disappeared when controlling for use of hand force.
► Among women we did not find any increased risk.

How might this impact on policy or clinical practice in the foreseeable future?

► Preventive strategies should focus on reducing the use of hand force in the working population.

Prospective studies have examined the work-relatedness of epicondylitis and hand or wrist tendinitis.^{5–12} Follow-up time ranges from 2 to 6 years and case definition varies from self-report to findings based on physical examination. Different definitions of exposure have been used to describe ranges of repetition, use of force, posture or hand-arm vibrations. Reviews have found some evidence of an exposure-response effect between force and risk of hand or wrist tendinitis and inconsistent associations with measures of repetition and posture alone.^{13 14} Combined exposures may be a multiplicative risk factor but a low incidence of disease in the studies combined with an uneven distribution of severe exposures in the study populations has prevented the investigation into such interactions.¹⁴ The risk of epicondylitis has been related to repetitive movements of the elbow, wrist or forearm and high force in combination with repetitive movements or awkward positions, but



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Aavang Petersen J, Brauer C, Thygesen LC, et al. *Occup Environ Med* 2022;**79**:55–62.

the evidence of force, posture and repetition as single risk factors is less clear.^{14 15}

It is paramount to separate measures of exposure from outcome in a prospective design to draw causal conclusions. However, obtaining enough statistical power can be difficult when objectively measured exposures and outcomes are used. Often, registers will ensure the objective outcomes, but not when examining temporally fluctuating and under-reported diseases and conditions such as epicondylitis and hand or wrist tendinitis.^{16 17} In that case, close monitoring and follow-up is needed, and this is costly and time-consuming.

Quantitative measurements for the exposure can be obtained by direct observation, self-report or technical measurements. Register information on job title and industry and measurements of job-related exposures can be combined in a job exposure matrix (JEM).¹⁸ This separates exposure and outcome and enables formation of large cohorts that can be followed for a long period of time.^{19–22} Using this method in the Danish registers an exposure-response effect of increasing wrist angular velocity on carpal tunnel syndrome has been found, but associations with force and combinations of force and repetition could not be examined due to lack of available force measurements.¹⁹ Expert-rated force has been used recently in JEMs, and may allow for investigation into combinations of force and repetition.^{23 24}

In parallel with the public treatments of diseases, private sector health programmes have emerged that offer treatments for musculoskeletal pain. Using such a health scheme as an outcome measure, where access to treatment is easily available and not dependent on a doctor's referral, may reveal more cases and strengthen the investigation into causality.

The purpose of this study was to investigate the association between repetitive movements of the wrist and the use of hand force and seeking treatment for pain in the distal upper extremities, using a JEM in a population covered in a health scheme.

METHODS

Study population

We established a cohort of persons aged 18–65 years covered by a private Danish pension fund, the PensionDanmark Health Scheme (PDHS), between October 2005 and December 2017. The Danish Civil registration system allowed for linkage to national registers at Statistics Denmark. Annual information on the main occupational titles were added by using the Occupation and Industry Register in the Danish Occupational Cohort with eXposure data based on the Danish version of the International Standard Classification of Occupation and the Danish Industrial Classification of All Economic Activities.^{25–27} Educational level was derived from the Danish version of the International Standard Classification of Education.²⁸

Exposure measurements

As a measure of repetition, we used the results of goniometric measurements from studies pooled in the EMINGO database, hosted by the Department of Occupational and Environmental Medicine, Lund, Sweden. Seventeen jobs were covered by the PDHS. Of these, 14 were measured in Denmark and 3 in Sweden, with the use of identical measurement protocols and equipment. The job functions in the Swedish measurements were comparable to Danish conditions.

The measurement programme has previously been described in detail.^{19 21 29} Briefly, companies and employees with exposures believed to be typical for the job titles selected for the study were asked to participate (see online supplemental table 1).

Measurements were performed on healthy, right-handed individuals with no musculoskeletal complaints. Jobs were selected in order to secure large contrasts in exposure. Measurements in at least 10 men or 10 women or both (if job functions differed with sex) were considered to provide accurate job exposure estimates.³⁰ Mean measurement time was 5.7 hours (3.1–6.7).

Whole working day (excluding pauses) wrist measurements were performed using biaxial goniometers (SG75, Biometrics, Newport, UK) and person-worn data-loggers (Logger Teknologi, Åkarp, Sweden) with a sampling frequency of 20 Hz. Angular velocity (°/s) was selected as measure for repetition. The goniometers were placed on the dorsal side of the wrist with the proximal part in the midline between radius and ulna and the distal part over the third metacarpal bone. Analyses were made for both sides but in this study, we only report results from the right (and dominant) side.

As a measure of force, the following Expert-JEM was used: five physicians, specialised in occupational medicine rated the level of the daily duration of significant use of hand force in 33 job titles with representative goniometric measurements whereof 17 were covered in the PDHS. This was rated in five categories and noted on rating cards for each job title: 'no or an insignificant part of the time', 'a small part of the time', 'some of the time', 'much of the time' and 'most of the time', scored 1–5. It was believed that ordering by quantitative category boundaries was unlikely to add any important information as it is easier to assess duration on a qualitative than on a quantitative scale. Each card included a short description of the work tasks performed during the measurements. The expert placed the cards in the relevant rating categories with a further split into two subcategories, low and high. The latter group was then scored with an addition of 0.5 points to the main category score, resulting in 10 scores (1–5.5 by steps of 0.5 score points). The experts received a written instruction and the instruction was further discussed at a meeting to ensure a common understanding of the rating task. Approximately 1 month after the first exposure rating the experts repeated their ratings, blinded to their first rating. The partial Spearman's correlations and overall agreement (weighted kappa) were both above 0.80 (details will be published separately). The combined mean value of the ratings was used as the exposure value.

Outcome and health-related confounders

The PDHS covers mainly employees in blue collar work, skilled and unskilled. The health scheme provides members with access to medical services including physiotherapy and chiropractors at numerous private health clinics (for further details see Pedersen and Arendt³¹). The PDHS is owned by a not-for-profit labour market pension fund and operated by a private health-care provider. The PDHS has existed since 2005 with step-wise inclusion of jobs in additional collective agreements. The purpose of the PDHS was to prevent and treat work-related illness but injuries originating from leisure time activities have since been included. Membership is provided by collective agreements between companies and workers. The mandatory, tax-deductible annual premium is €48 per member and is automatically deducted in the pension contributions. The use of the service is not reported to neither employer nor labour union. Number of treatments received are typically not restricted and no physician referral is required. All individual treatments in the PDHS are registered at every visit. Type of treatment and body region of concern are registered in the invoice sent by the healthcare provider to the PDHS. We used chiropractic and

physiotherapeutic treatments of hands, wrists, elbows and arms as outcome. To ensure that the registered treatment was more than just a trial visit in a newly available service, we defined outcome as having a second treatment within 30 days after the first treatment.

First diagnosis of rheumatoid arthritis (RA, International Classification of Diseases, 10th Edition (ICD-10): M05) and any upper extremity fractures (ICD-10: S42, S47, S52, S54, S57, S58, S62) were identified through the Danish National Patient Register.³² RA was included as a confounding risk factor from the date of diagnosis and onwards and fractures were included for a period of 2 years after diagnosis.

Statistical analysis

The cohort was followed from the first day of coverage in one of the 17 jobs in the PDHS from 1 October 2005 until the date of the second treatment within 30 days after the first treatment, the last day of registered coverage or 31 December 2017, whichever came first. A person was allowed as many periods of coverage as possible but a minimum of 30 days in total was required.

Poisson regression models were used to determine incidence risk ratios (IRRs) of treatment in relation to wrist angular velocity, use of hand force and job. The logarithm of risk time was used as offset value.

The main analyses included the exposures (wrist velocity and use of hand force) as continuous and categorical variables. The data revealed an apparent skewness of sex-distribution across quintiles of exposure, requiring us to split analyses into men and women. We also performed Spearman's rank correlation analyses between wrist velocity and use of hand force in quintiles.

We also analysed sex-specific associations between job groups and seeking treatment. To ensure enough power, only jobs with >300 persons were included. For male job groups, forklift operators were the reference group and for female job groups office workers were the reference group (ie, the lowest exposed job group).

As sensitivity analyses, we adjusted results for educational level (elementary, high school, vocational, higher or unknown). The importance of outcome definition was evaluated by calculating IRRs based on first ever treatment. We had outcome information available for the year 2018, but without register information on the diagnoses of RA and fractures. The effect of an additional year of follow-up was investigated, but only with adjustment for age and calendar-year. Effect of time covered in the PDHS was investigated by restricting analysis to only those covered for >90% of the time. We examined sex-specific interactions between high/low exposure using median split.

We report results as crude and adjusted IRRs with 95% CIs. Adjustments were made based on a directed acyclic graph for age, age \times age, calendar year, diagnosis of RA and upper extremity fractures within the last 2 years (model 1). Finally, we multiple adjusted for the two exposures (repetition or hand force) (model 2).

Data were analysed using SAS V.9.4 (SAS Institute, Cary, North California, USA).

RESULTS

Cohort characteristics

A total of 202 747 members were followed for 616 759.8 person-years (figure 1). Table 1 shows the characteristics of the 17 jobs whereof 3 jobs had measurements on both men and women. Almost 70% of the cohort was men. The age at entry was quite similar across jobs with hairdressers and pig farm workers as




Figure 1 Flow chart of the creation of the cohort.

the youngest groups. The cohort members were followed for an average of 3.0 years with time covered as a truck drivers as the longest (4.4 years). Cohort members were covered in the PDHS for an average of 88% of the follow-up time.

There was low correlation between quintiles of wrist angular velocity and hand/wrist force (Spearman's correlation coefficient 0.04). Sex distribution varied considerably across quintiles of exposures (table 2). Men had more risk time with fractures than women and persons with RA were present in the cohort for only a very short time after the diagnosis. More than half of the cohort was skilled workers, 35.7% unskilled and 3.5% higher educated.

A total of 1085 first-time treatments was found, with elbows and arms (forearm or upper-arm) accounting for 86.4% of the treatments. Hands and wrists accounted for 4.6% and 9.0% of the treatments, respectively. The body regions did not differ between men and women. Men had a lower incidence rate than women of 15.3 vs 24.0 treatments per 10 000 person-years (table 3).

Wrist angular velocity

Analysing wrist angular velocity as a continuous variable, we found a small but significant association among men in model 1 (table 3). Adjusting for force did not alter this result (model 2). Wrist angular velocity as categorical exposure had a significantly increased risk among men in the highest tertile compared with the lowest tertile, with an IRR of 1.48 (95% CI 1.15 to 1.92) in model 1. Estimates became insignificant after adjusting for hand force in model 2.

Among women, a significantly protective association with increasing wrist angular velocity was found in both models. Only the highest tertile was significantly different from the lowest in wrist angular velocity. However, the same jobs were categorised in high velocity and medium hand force and therefore the effect of adjusting for hand force was small.

Wrist force

The association between use of hand force and treatment was significant and unaffected by adjusting for wrist angular velocity among men, with an IRR of 1.50 (95% CI 1.38 to 1.63). Hand force had a clear exposure-response effect on the outcome in model 1, with an IRR of 2.23 (95% CI 1.83 to 2.72) and 2.65 (95% CI 2.13 to 3.29) in the second and third tertiles, respectively (table 3). This effect was still evident after adjusting for angular velocity.

Table 1 Characteristics of the cohort: persons covered in the PensionDanmark Health Scheme in the period 2005–2017 aged 18–65 years

Job title	Measured sex (N)	N, at entry Men/Women	Median age at entry (IQR)	Total risk time, person- years	Wrist angular velocity (°/s), 50th percentile, (SD)	Wrist force, expert mean*
Office worker	Men (18)	3667	34.5 (25.3–47.4)	4961.2	3.5 (1.9)	1.00
Fork-lift operator	Men (10)	6105/460	38.2 (28.5–47.3)	15 737.1	5.6 (2.6)	1.30
Office worker	Women (25)	7499	38.3 (27.1–48.6)	14 748.1	6.1 (3.5)	1.00
Truck driver	Men (11)	32 476/518	39.7 (30.1–48.4)	145 085.8	6.4 (3.8)	2.15
Plumber	Men (11)	4199/5	38.1 (30.0–46.1)	22 118.5	9.3 (2.2)	4.00
Smith	Men (12)	6262/43	37.6 (27.5–47.6)	9732.8	10.2 (2.2)	4.25
Wood processing worker	Men (10)	2515/411	40.8 (29.9–49.8)	6327.8	10.4 (3.0)	3.10
Gardener	Men (11)	2681	34.4 (25.3–47.1)	4763.9	11.6 (2.4)	4.15
Carpenter	Men (10)	23 772/53	30.9 (23.8–43.4)	82 594.7	11.6 (2.3)	4.20
Scaffolder	Men (10)	5284/52	38.2 (28.7–48.5)	10 316.1	12.9 (1.6)	4.40
Construction worker	Men (10)	15 789/155	39.0 (27.8–48.8)	42 444.5	14.0 (3.8)	4.15
Bricklayer	Men (10)	10 096/36	34.4 (25.4–47.7)	35 651.7	14.2 (4.7)	4.15
Garbage collector	Men (11)	3916/109	39.1 (30.2–47.6)	12 287.1	14.3 (2.4)	3.10
Pig farm worker	Men (10)	2770/915	22.7 (19.0–29.7)	5128.5	14.6 (4.8)	3.15
Hairdresser	Women (10)	631/10 325	25.1 (21.7–32.8)	44 347.8	16.3 (2.8)	1.25
Gardener	Women (9)	532	34.4 (26.0–43.7)	911.1	18.4 (7.2)	4.15
Kitchen assistant	Women (10)	11 994/12 282	28.0 (22.0–38.9)	45 584.1	21.5 (4.0)	2.45
Cleaning assistant	Women (24)	15 753/27 954	37.9 (28.7–48.0)	101 735.8	27.9 (3.2)	2.55
Laundry worker	Men (10)	1066	33.2 (25.2–44.1)	3383.4	30.1 (8.5)	2.35
Laundry worker	Women (13)	2387	39.8 (30.2–49.0)	8890.2	31.8 (7.0)	2.35
Total		148 976/63 736	35.1 (25.7–46.4)	616 759.8	14.5	2.96
Minimum–maximum	9–25				3.5–31.8	1.0–4.4

Exposure intensities of wrist angular velocity are based on the electro-goniometric measurements. Wrist force is based on expert rating. Jobs are arranged according to increasing wrist velocity.

*Full range was 1–5.5.

N, number.

Neither as a continuous nor a categorical variable was the use of wrist force positively associated with treatment among women in both models.

Job groups

Figure 2 shows the sex-specific associations between job group and treatment in jobs with >300 persons represented. Incidence rates ranged from 6.2 to 65.9 per 10 000 person-years. Compared with fork-lift operators, several jobs showed a significantly higher adjusted IRR ranging from 1.71 to 3.30 among men (figure 2A). Truck drivers had a significantly lower risk of treatment for D-UEMSD. Among women, an increased risk, although non-significant, compared with office workers was found only among gardeners, IRR 2.03 (95% CI 0.87 to 4.72). Cleaning assistants and laundry workers had a significantly decreased risk with an IRR of 0.64 (95% CI 0.47 to 0.87) and 0.43 (95% CI 0.25 to 0.76), respectively (figure 2B).

Sensitivity analyses

Adjusting for educational level attenuated results, leaving only results among men statistically significant. Using the first ever treatment as outcome did not change the overall results, although the inverse association with categorical wrist angular velocity among women became statistically significant in model 2. An additional year of follow-up or restricting analysis to those covered for >90% of the time did not change the results. Fully adjusted interaction analyses with dichotomised high (H)/low (L) force (F)/velocity (V) for men showed an IRR of 3.88 (95% CI 3.09 to 4.87) for HF/LV, 2.36 (95% CI 1.77 to 3.15) for LF/HV and 2.66 (95% CI 2.08 to 4.87) for HF/HV. For women

IRRs of HF/LV, LF/HV and HF/HV were 0.44 (95% CI 0.14 to 1.42), 0.69 (95% CI 0.50 to 0.99) and 0.77 (95% CI 0.56 to 0.99), respectively.

DISCUSSION

Key results

The aim of this study was to analyse whether repetitive and forceful movements were associated with pain requiring treatment in the arms or hands. With the use of a JEM based on objective measurements of repetitive wrist movements and expert-rated hand force, we found that the use of hand force, controlled for repetitive movements, increased the risk with an exposure-response pattern in men, but not in women. Increasing velocity of the movements of the hand, controlled for force, did not increase the risk of pain requiring treatment in neither men nor women. In fact, high velocity had a significant protective effect in women.

Limitations

Some methodological limitations must be addressed. Outcome definition was based on the reported anatomical region of concern by the treating physiotherapist or chiropractor at the time of treatment. This gives the advantage of being a real-time observation but may limit the specificity of the outcome. We anticipated that seeking treatment for pain in the hands, wrists, elbows and arms primarily represented disorders such as epicondylitis of the elbow and tenosynovitis of the hand or wrist. Treatment for functional limitations is also a possibility, although expected to be a minor part. However, we were not able to

Table 2 Distribution of potential confounders across quintiles of exposures

	0–<20th percentile 3.5–≤7.9	20th–≤40th percentile 7.9–≤11.6	40th–≤60th percentile 11.6–≤14.2	60th–≤80th percentile 14.2–≤19.9	80th–≤100th percentile 19.9–≤31.8	Total
Wrist angular velocity, °/s						
RT—total	180 532.0	120 773.8	93 176.1	62 684.4	159 593.6	616 759.8
Men	162 870.2	119 461.9	92 688.8	17 889.7	61 889.6	454 800.2
Women	17 661.7	1311.9	487.3	44 794.8	97 704.0	161 959.6
Mean age (IQR)	43.1 (34.3–52.0)	38.1 (27.8–47.8)	40.1 (29.3–50.3)	33.7 (24.5–41.2)	38.5 (28.2–48.0)	39.6 (29.1–49.4)
Educational level						
Unskilled*, %	51.3	7.2	31.9	30.1	44.8	35.7
Skilled, %	41.4	91.4	62.5	65.6	25.2	53.4
Higher, %	3.0	0.8	1.3	1.5	8.8	3.5
Unknown, %	4.3	0.7	4.3	2.8	21.8	7.4
RT with rheumatoid arthritis	651.7	301.7	204.1	221.6	549.3	1928.4
Men	553.0	293.9	201.1	72.5	107.3	1227.7
Women	98.7	7.8	3.1	149.1	442.1	700.7
RT with fractures†	660.9	686.7	451.6	156.8	355.4	2311.5
Men	618.5	682.8	451.6	99.8	173.2	2026.9
Women	42.4	2.9	0	57.0	182.2	284.5
	0–<20th percentile 1.00–≤1.73	20th–≤40th percentile 1.73–≤2.50	40th–≤60th percentile 2.50–≤3.58	60th–≤80th percentile 3.58–≤4.15	80th–≤100th percentile 4.15–≤4.40	Total
Wrist force						
RT—total	79 804.2	202 943.3	125 479.1	105 889.6	102 643.6	616 759.8
Men	21 414.7	168 765.5	57 701.6	104 564.5	102 354.0	454 800.2
Women	58 389.5	34 177.9	67 777.6	1325.1	289.6	161 959.6
Mean age (IQR)	36.5 (26.1–45.6)	41.4 (31.6–50.9)	39.8 (30.0–49.8)	40.3 (30.0–50.0)	37.5 (27.1–47.1)	39.6 (29.1–49.4)
Educational level						
Unskilled*, %	30.9	51.7	44.9	25.4	9.2	35.7
Skilled, %	61.5	39.5	23.9	70.9	88.3	53.4
Higher, %	5.4	2.8	8.3	1.2	0.8	3.5
Unknown, %	2.3	6.0	22.9	2.6	2.7	7.4
RT with rheumatoid arthritis	304.0	708.8	427.9	243.4	244.3	1928.4
Men	74.0	527.4	148.0	240.3	238.0	1227.7
Women	230.0	181.4	279.9	3.1	6.3	700.7
RT with fractures	161.5	715.5	318.3	461.4	654.7	2311.5
Men	83.7	636.9	191.2	0.9	654.7	2026.9
Women	77.8	78.6	127.2	460.5	0	284.5

*Unskilled defined as elementary or high school.

†Fractures defined as with effect 2 years from date of diagnosis.

RT, risk time in years.

perform analyses on elbow and hands separately. This lack of specificity in outcome might have diluted the associations. The PDHS as a source of outcome is unvalidated and future studies are needed to confirm our results.

A possible bias could have been different care-seeking behaviour in job groups or between sexes, for example, because of accessibility or information.³³ However, the population was homogenous regarding educational level why this bias may have been limited. The use of the PDHS was comparable to the use of physical therapy in the general population. According to Statistics Denmark, approximately 6%–8% of the Danish population had at least one treatment with doctor-referred physiotherapy annually in 2008–2018 compared with 5%–7% of the members in the PDHS in the same period.³⁴ Also, leisure time physical activity was not known and may be an unmeasured confounder, but previously the confounding effect on pain outcomes have been minor.³⁵

A JEM entails the risk of non-differential misclassification of individual exposure due to variation within groups compared with the variation between groups.^{36–37} The job groups were

selected to represent large exposure variation to counteract the effect of such misclassification. Angular velocity was chosen over mean power frequency as the measurement of repetition, as it has been shown to be consistently associated with carpal tunnel syndrome, tendinitis and elbow disorders.^{19–38} The fewer job groups and a lower absolute number of events might explain the results among the women.

The expert-rated use of force did not take sex into account, which may have caused misclassification. As an example, male and female gardeners had quite different angular velocity but the same level of force in our JEM. This might represent a misclassification, as male gardeners typically perform more strenuous tasks than females. Comparison with a gold standard using objective measures of force, such as electromyography, is difficult due to lack of comparable studied jobs. However, as an example, hairdressers have been studied with electromyography and relatively high levels of muscular activity and low levels of rest were found compared with jobs traditionally considered to be forceful such as meat cutters.³⁸ This was not reflected in the expert-rating of force of the hairdressers, exposing the difficulties in choosing the

Table 3 The risk of treatment for pain in the distal upper extremities with increasing levels of wrist angular velocity and exertion of force

Physical exposures	Events	Person-years	IR	IRR, crude (95% CI)	IRR, model 1* (95% CI)	IRR, model 2 (95% CI)
Wrist angular velocity						
<i>Continuous variable</i>						
Men	698	454 800.2	15.3	1.02 (1.01 to 1.03)	1.02 (1.00 to 1.03)	1.02 (1.00 to 1.03)
Women	387	161 959.6	24.0	0.99 (0.97 to 1.00)	0.98 (0.97 to 0.99)	0.97 (0.96 to 0.99)
<i>Categorical variable†</i>						
Low (ref)						
Men (3.5–<10.4 °/s)	212	199 898.5	10.6	1	1	1
Women (3.5–<10.4 °/s)	67	18 757.3	35.7	1	1	1
Medium						
Men (10.4–14.3 °/s)	390	187 154.1	20.8	1.96 (1.66 to 2.32)	2.08 (1.73 to 2.50)	1.20 (0.94 to 1.54)
Women (10.4–18.3 °/s)	89	44 539.4	20.0	0.56 (0.41 to 0.77)	0.82 (0.59 to 1.14)	0.82 (0.59 to 1.14)
High						
Men (>14.3 °/s)	96	67 747.8	14.2	1.34 (1.05 to 1.70)	1.48 (1.15 to 1.92)	1.13 (0.86 to 1.49)
Women (>18.3 °/s)	231	97 704.0	23.6	0.66 (0.50 to 0.87)	0.70 (0.53 to 0.93)	0.71 (0.53 to 0.93)
Wrist force						
<i>Continuous variable</i>						
Men	698	454 800.2	15.3	1.48 (1.37 to 1.61)	1.50 (1.38 to 1.63)	1.50 (1.38 to 1.63)
Women	387	161 959.6	24.0	0.97 (0.84 to 1.13)	0.87 (0.74 to 1.02)	1.13 (0.89 to 1.43)
<i>Categorical variable†</i>						
Low (ref)						
Men (1.0–2.45)	765	190 180.1	8.7	1	1	1
Women (1.0–2.15)	147	59 919.1	24.5	1	1	1
Medium						
Men (>2.45–4.15)	313	162 266.1	19.3	2.22 (1.84 to 2.68)	2.23 (1.83 to 2.72)	1.99 (1.55 to 2.56)
Women (>2.15–2.55)	231	97 704.0	23.6	0.96 (0.78 to 1.19)	0.79 (0.63 to 0.99)	0.71 (0.53 to 0.94)
High						
Men (>4.15)	220	102 354.0	21.5	2.48 (2.02 to 3.03)	2.65 (2.13 to 3.29)	2.31 (1.72 to 3.09)
Women (>2.55)	9	3377.6	26.6	1.09 (0.55 to 2.13)	1.07 (0.54 to 2.10)	1.06 (0.54 to 2.08)

Estimates in bold significant at $p < 0.05$. Model 1: adjusted for age and calendar-year, diagnoses of rheumatoid arthritis and fractures of the wrist, elbow or arm. Model 2: adjusted for model 1 and the other exposure, that is, force or repetition.

*No cases of incident treatment among women with rheumatoid arthritis. This variable was left out in the analyses for women.

†Tertiles of exposure were defined for sexes separately.

IR, incidence rate per 10 000 person-years; IRR, incidence rate ratio.

relevant exposure measure for force. The possible misclassification using the expert-rated force will in such case result in too low estimates and may to some extent explain the negative findings among women. The strength of this study is the prospective design and the use of objective exposure and outcome, avoiding recall bias. However, the methods used are crude and introduces non-differential biases and entails risk of attenuation of results and negative findings, such as among the women. Therefore, the positive associations found among the men are more reliable.

Interpretation

Our results among the men was in line with the results from other more recent prospective studies using both individual and group-based exposures, where a pattern of an exposure-response effect between force and a risk of D-UEMSD has been found. Thomsen *et al* used a task exposure matrix, prospectively, on the job titles of 3123 employees in 19 industrial settings and found insignificant effect of high repetition. Force, on an observer assessed 5-point scale, remained a significant risk factor when repetition was included in the model.³⁹ Force was significantly associated with both self-reported pain and possible tendinitis based on physical examination. Harris *et al* followed 413 hand-intensive workers for 28 months in 3 months intervals for wrist tendinosis and found similar results of no association with

any individualised measure of repetition adjusted for force.⁵ Per cent time spent in heavy pinch grip showed an exposure-response relationship after adjustment for repetition, although statistical power was low due to sample size. Nordander *et al* used wrist angular velocity and wrist extensor muscular activity from pooled studies in a cross-sectional design.³⁸ Associations were found between angular velocity and prevalence rates of complaints in the right elbow and hand and symptoms of carpal tunnel syndrome in both sexes and medial epicondylitis among men only. However, the cross-sectional design did not allow for causal conclusions and no adjustments for force were made.

A systematic review where only 2/12 studies were prospective and a wide range of outcomes were included have found that force and repetition interact significantly.⁴⁰ As combinations of exposures were limited in our study, interpretations of our interaction results should be done with caution, and the interaction effects between force and repetition needs to be confirmed in well-designed studies reducing the risk of recall bias. Our study cohort mainly consisted of organised, skilled and unskilled workers, and might only be generalisable to similar populations. However, as we used generic exposure variables and the range of exposure was wide, especially among men, covering both very low and very high exposed workers, the results may have broad application.




Figure 2 Association between treatment for pain in the upper extremities and job title, adjusted incidence rate ratio (IRR) sorted by increasing wrist velocity. REF indicates reference group. (A) Men: job titles with >300 men. (B) Women: job titles with >300 women. Adjusted for age, age \times age, calendar-year and diagnoses of rheumatoid arthritis and fractures of the wrist, elbow or arm.

Future research on occupational exposure and D-UEMSD should focus on further disentanglement of repetitive movements, use of hand force to explore exposure-response associations and possible interactions. This should be possible by including additional jobs with goniometric measurements or applying expert-rated JEMs on large populations in prospective designs in order to close the exposure information gaps that now exist.

CONCLUSION

The purpose of this study was to investigate the association between repetitive movements of the wrist and the use of hand force and pain in the arm and hand. We found a weak association with repetition among men but without a clear exposure-response pattern and no association among women. We found that exertion of force was convincingly associated with seeking treatment among men. This was not found in women, probably due to some degree of exposure misclassification and a lower number of investigated job groups.

Acknowledgements The authors thank Henrik Enquist and colleagues at the Department of Occupational and Environmental Medicine, Lund, Sweden for hosting and providing access to the EMINGO database. They also thank PensionDanmark for providing access to the PDHS database for research purposes. They also thank Tine Steen Erichsen, Ellen Bøtcher Pedersen and Jens Peter Bonde, Department of Occupational and Environmental Medicine, Bispebjerg and Frederiksberg Hospital, University of Copenhagen for expert-rating for the JEM together with the authors Charlotte Brauer and Jane Frølund Thomsen. Finally, they thank Thomas Heilskov-Hansen, Department of Occupational and Environmental Medicine, Bispebjerg and Frederiksberg Hospital, University of and Annett Dalbøge, Department of

Occupational Medicine, Danish Ramazzini Centre, Aarhus University Hospital, for the collection of technical exposure measurements in Denmark.

Contributors JFT, CBL, LCT, CB, EMF and JAP designed the study and JAP, JFT and EMF acquired the data; JAP and EMF analysed the data and JAP drafted the manuscript; JFT, LCT, CBL, CB, EMF and JAP interpreted the data. All authors critically reviewed the draft for important intellectual content. All authors agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding This study was supported by grants from the Danish Working Environment Research Fund (grant # 20175100917) and The Research Centre for Hairdressers and Beauticians (grant # 150217).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval In Denmark, register and questionnaire studies do not require approval by committees on biomedical research ethics. The project was approved by the Knowledge Centre on Data Protection Compliance under the records of processing regarding health science research projects within the Capital Region of Denmark (VD-2018-521).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data from the JEM are available upon reasonable request. Data from the Danish national registers are available through online access at Statistics Denmark under standard conditions.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Jonathan Aavang Petersen <http://orcid.org/0000-0003-0635-4144>

REFERENCES

- European Agency for Safety and Health at Work. *Osh in figures: work-related musculoskeletal disorders in the EU-Facts and figures*. Luxembourg, 2010.
- Bevan S. Economic impact of musculoskeletal disorders (MSDs) on work in Europe. *Best Pract Res Clin Rheumatol* 2015;29:356–73.
- da Costa JT, Baptista JS, Vaz M. Incidence and prevalence of upper-limb work related musculoskeletal disorders: a systematic review. *Work* 2015;51:635–44.
- Roquelaure Y, Ha C, Rouillon C, et al. Risk factors for upper-extremity musculoskeletal disorders in the working population. *Arthritis Rheum* 2009;61:1425–34.
- Harris C, Eisen EA, Goldberg R, et al. 1st place, PREMUS best paper competition: workplace and individual factors in wrist tendinitis among blue-collar workers—the San Francisco study. *Scand J Work Environ Health* 2011;37:85–98.
- Descatha A, Albo F, Leclerc A, et al. Lateral epicondylitis and physical exposure at work? A review of prospective studies and meta-analysis. *Arthritis Care Res* 2016;68:1681–7.
- Herquelot E, Guéguen A, Roquelaure Y, et al. Work-related risk factors for incidence of lateral epicondylitis in a large working population. *Scand J Work Environ Health* 2013;39:578–88.
- Descatha A, Dale AM, Jaegers L, et al. Self-reported physical exposure association with medial and lateral epicondylitis incidence in a large longitudinal study. *Occup Environ Med* 2013;70:670–3.
- Garg A, Kapellusch JM, Hegmann KT, et al. The strain index and TLV for HAL: risk of lateral epicondylitis in a prospective cohort. *Am J Ind Med* 2014;57:286–302.
- Fan ZJ, Silverstein BA, Bao S, et al. The association between combination of hand force and forearm posture and incidence of lateral epicondylitis in a working population. *Hum Factors* 2014;56:151–65.
- Harris-Adamson C, You D, Eisen EA, et al. The impact of posture on wrist tendinitis among blue-collar workers: the San Francisco study. *Hum Factors* 2014;56:143–50.
- Nambiema A, Bertrais S, Bodin J, et al. Proportion of upper extremity musculoskeletal disorders attributable to personal and occupational factors: results from the French pays de la Loire study. *BMC Public Health* 2020;20:1–13.

- 13 NIOSH. Musculoskeletal disorders and workplace factors. A critical review of epidemiologic evidence for work-related musculoskeletal disorders of the neck, upper extremity, and low back, 1997. Available: <https://www.cdc.gov/niosh/docs/97-141/>
- 14 Keir PJ, Farias Zuniga A, Mulla DM, *et al.* Relationships and mechanisms between occupational risk factors and distal upper extremity disorders. *Hum Factors* 2021;63:5–31.
- 15 Seidel DH, Ditchen DM, Hoehne-Hückstädt UM, *et al.* Quantitative measures of physical risk factors associated with work-related musculoskeletal disorders of the elbow: a systematic review. *Int J Environ Res Public Health* 2019;16:130–23.
- 16 Gummesson C, Isacson S-O, Isacson AH, *et al.* The transition of reported pain in different body regions – a one-year follow-up study. *BMC Musculoskelet Disord* 2006;7:1–7.
- 17 Fan ZJ, Bonauto DK, Foley MP, *et al.* Underreporting of work-related injury or illness to workers' compensation: individual and industry factors. *J Occup Environ Med* 2006;48:914–22.
- 18 Plato N, Steineck G. Methodology and utility of a job-exposure matrix. *Am J Ind Med* 1993;23:491–502.
- 19 Lund CB, Mikkelsen S, Thygesen LC, *et al.* Movements of the wrist and the risk of carpal tunnel syndrome: a nationwide cohort study using objective exposure measurements. *Occup Environ Med* 2019;76:519–26.
- 20 Heilskov-Hansen T, Mikkelsen S, Svendsen SW, *et al.* Exposure-response relationships between movements and postures of the wrist and carpal tunnel syndrome among male and female house painters: a retrospective cohort study. *Occup Environ Med* 2016;73:401–8.
- 21 Dalbøge A, Hansson Gert-Åke, Frost P, *et al.* Upper arm elevation and repetitive shoulder movements: a general population job exposure matrix based on expert ratings and technical measurements. *Occup Environ Med* 2016;73:553–60.
- 22 Tabatabaeifar S, Svendsen SW, Frost P. Carpal tunnel syndrome as sentinel for harmful hand activities at work: a nationwide Danish cohort study. *J Occup Environ Med* 2020;62:375–82.
- 23 Kirkeby L, Svendsen SW, Hansen TB, *et al.* Surgery for trapeziometacarpal osteoarthritis in relation to cumulative occupational hand force requirements: a Danish nationwide cohort study. *Occup Environ Med* 2021;78:92–7.
- 24 Svendsen SW, Johnsen B, Fuglsang-Frederiksen A, *et al.* Ulnar neuropathy and ulnar neuropathy-like symptoms in relation to biomechanical exposures assessed by a job exposure matrix: a triple case-referent study. *Occup Environ Med* 2012;69:773–80.
- 25 Pedersen CB. The Danish civil registration system. *Scand J Public Health* 2011;39:22–5.
- 26 Flachs EM, Petersen SEB, Kolstad HA, *et al.* Cohort Profile: DOC*X: a nationwide Danish occupational cohort with eXposure data - an open research resource. *Int J Epidemiol* 2019;48:1413–1413k.
- 27 Petersson F, Baadsgaard M, Thygesen LC. Danish registers on personal labour market affiliation. *Scand J Public Health* 2011;39:95–8.
- 28 Jensen VM, Rasmussen AW. Danish education registers. *Scand J Public Health* 2011;39:91–4.
- 29 Heilskov-Hansen T, Svendsen SW, Frølund Thomsen J, *et al.* Sex differences in task distribution and task exposures among Danish house painters: an observational study combining questionnaire data with biomechanical measurements. *PLoS One* 2014;9:e110899.
- 30 Hoozemans MJ, Burdorf A, van der Beek AJ, *et al.* Group-Based measurement strategies in exposure assessment explored by bootstrapping. *Scand J Work Environ Health* 2001;27:125–32.
- 31 Pedersen MS, Arendt JN. Bargaining for health: a case study of a collective agreement-based health program for manual workers. *J Health Econ* 2014;37:123–36.
- 32 Lynge E, Sandegaard JL, Rebolj M. The Danish national patient register. *Scand J Public Health* 2011;39:30–3.
- 33 Thompson AE, Anisimowicz Y, Miedema B, *et al.* The influence of gender and other patient characteristics on health care-seeking behaviour: a QUALICOPC study. *BMC Fam Pract* 2016;17:1–7.
- 34 Engmann TS. Flere danskere går TIL fysioterapeut (opdateret). Copenhagen, 2019. Available: <https://www.dst.dk/da/Statistik/bagtal/2019/2019-04-11-flere-danskere-gaar-til-fysioterapeut>
- 35 van den Heuvel SG, van der Beek AJ, Blatter BM, *et al.* Do work-related physical factors predict neck and upper limb symptoms in office workers? *Int Arch Occup Environ Health* 2006;79:585–92.
- 36 Armstrong BG. Effect of measurement error on epidemiological studies of environmental and occupational exposures. *Occup Environ Med* 1998;55:651–6.
- 37 Peters S. Although a valuable method in occupational epidemiology, job-exposure -matrices are no magic fix. *Scand J Work Environ Health* 2020;46:231–4.
- 38 Nordander C, Ohlsson K, Akesson I, *et al.* Exposure-response relationships in work-related musculoskeletal disorders in elbows and hands - A synthesis of group-level data on exposure and response obtained using uniform methods of data collection. *Appl Ergon* 2013;44:241–53.
- 39 Thomsen JF, Mikkelsen S, Andersen JH, *et al.* Risk factors for hand-wrist disorders in repetitive work. *Occup Environ Med* 2007;64:527–33.
- 40 Gallagher S, Heberger JR. Examining the interaction of force and repetition on musculoskeletal disorder risk: a systematic literature review. *Hum Factors* 2013;55:108–24.