

Treatment After Anterior Cruciate Ligament Injury

Panther Symposium ACL Treatment Consensus Group

Theresa Diermeier, MD, Benjamin B. Rothrauff, MD, PhD, Lars Engebretsen, MD, PhD, Andrew D. Lynch, PhD, PT, Olufemi R. Ayeni, MD, PhD, Mark V. Paterno, PT, PhD, John W. Xerogeanes, MD, Freddie H. Fu, MD, DSc, DPs (Hon), Jon Karlsson, MD, PhD, Volker Musahl,* MD, and the Panther Symposium ACL Treatment Consensus Group

Investigation performed at University of Pittsburgh, Pittsburgh, Pennsylvania, USA

Treatment strategies for anterior cruciate ligament (ACL) injuries continue to evolve. Evidence supporting best-practice guidelines for the management of ACL injury is to a large extent based on studies with low-level evidence. An international consensus group of experts was convened to collaboratively advance toward consensus opinions regarding the best available evidence on operative versus nonoperative treatment for ACL injury. The purpose of this study was to report the consensus statements on operative versus nonoperative treatment of ACL injuries developed at the ACL Consensus Meeting Panther Symposium 2019. There were 66 international experts on the management of ACL injuries, representing 18 countries, who were convened and participated in a process based on the Delphi method of achieving consensus. Proposed consensus statements were drafted by the scientific organizing committee and session chairs for the 3 working groups. Panel participants reviewed preliminary statements before the meeting and provided initial agreement and comments on the statement via online survey. During the meeting, discussion and debate occurred for each statement, after which a final vote was then held. Ultimately, 80% agreement was defined a priori as consensus. A total of 11 of 13 statements on operative versus nonoperative treatment of ACL injury reached consensus during the symposium. Overall, 9 statements achieved unanimous support, 2 reached strong consensus, 1 did not achieve consensus, and 1 was removed because of redundancy in the information provided. In highly active patients engaged in jumping, cutting, and pivoting sports, early anatomic ACL reconstruction is recommended because of the high risk of secondary meniscal and cartilage injuries with delayed surgery, although a period of progressive rehabilitation to resolve impairments and improve neuromuscular function is recommended. For patients who seek to return to straight-plane activities, nonoperative treatment with structured, progressive rehabilitation is an acceptable treatment option. However, with persistent functional instability, or when episodes of giving way occur, anatomic ACL reconstruction is indicated. The consensus statements derived from international leaders in the field will assist clinicians in deciding between operative and nonoperative treatment with patients after an ACL injury.

Keywords: ACL injury; ACL reconstruction; nonoperative treatment

Anterior cruciate ligament (ACL) injuries are one of the most common injuries of the knee, with an incidence of approximately 85 per 100,000 in patients aged between 16 and 39 years.^{26,34,46} The ACL is the primary stabilizer of the knee limiting anterior tibial translation and internal rotation, with deficiency resulting in anterior and rotatory instability.^{51,90} The most common mode of injury is a noncontact mechanism during pivoting, cutting, and jumping with the knee slightly flexed and in a valgus position.^{1,5}

Both operative and nonoperative treatments of an ACL injury continue to evolve.^{21,22,32,75} Improved understanding of the structure and function of the native ACL has supported the development and adoption of anatomic ACL reconstruction techniques.³ In parallel, increased recognition of the resilience of the neuromuscular system in achieving dynamic, functional knee stability despite ACL deficiency has concurrently supported nonoperative treatment as a viable strategy in some patients.^{13,23}

Successful outcomes after both operative and nonoperative treatments necessitate progressive rehabilitation, which entails staged and phase-adjusted physical therapy with the aim to address impairments, achieve functional stability, and safely return to sport.⁶¹ The acute phase after

The Orthopaedic Journal of Sports Medicine, 8(6), 2325967120931097
DOI: 10.1177/2325967120931097
© The Author(s) 2020

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For article reuse guidelines, please visit SAGE's website at <http://www.sagepub.com/journals-permissions>.

TABLE 1
Anatomic ACL Reconstruction Checklist
Based on van Eck et al^{80a}

1	Individualization of surgery for each patient
2	Use of 30° scope
3	Use of an accessory medial portal
4	Direct visualization of the femoral insertion site
5	Measuring the femoral insertion site dimensions
6	Visualizing the lateral intercondylar ridge
7	Visualizing the lateral bifurcate ridge
8	Placing the femoral tunnel(s) in the femoral ACL insertion site
9	Transportal drilling
10	Direct visualization of the tibial insertion site
11	Measuring the tibial insertion site dimensions
12	Placing the tibial tunnel(s) in the tibial ACL insertion site
13	Femoral fixation
14	Tibial fixation
15	Knee flexion angle during femoral tunnel drilling
16	Graft type
17	Graft tensioning

^aACL, anterior cruciate ligament.

the injury or surgery focuses on the elimination of residual symptoms (effusion, pain) and impairments (range of motion, quadriceps activation and strength). Subsequently, neuromuscular and perturbation training are implemented to improve knee stabilization.^{9,19} The last phase aims to further optimize muscular strength, return to preinjury sport level through sport-specific exercises, and assess psychological readiness for the return to sport.³ Any discussion of nonoperative treatment within this consensus document implies the completion of a progressive, staged rehabilitation protocol.

Similarly, any discussion of operative treatment implies anatomic ACL reconstruction (Table 1), which intends to restore the ACL to its native dimensions, collagen orientation, and insertion sites.⁸¹

Anatomic ACL reconstruction includes both single- and double-bundle techniques, followed by a progressive rehabilitation program that considers the natural healing cascade and ligamentization of the graft.⁶² After fixation during ACL reconstruction, a biological graft transitions from a tendon to a structure with ultrastructural, biochemical, and mechanical properties more similar to the native ACL.⁷¹ These properties of the graft depend on the phase of ligamentization, with the minimum graft strength occurring between 4 and

12 weeks postoperatively.^{62,71} Comprehensive rehabilitation after operative ACL reconstruction is also paramount for clinical outcome and return to sport.

Whereas operative treatment aims to reduce laxity, non-operative treatment aims to reduce functional instability and both thereby prevent further damage to the menisci and cartilage, which may contribute to posttraumatic osteoarthritis (PTOA).^{55,82} Functional bracing, intended to reduce the risk of ACL injury by decreasing peak ligament strain, has not yet been conclusively shown to achieve this goal, as the evidence is still limited.^{29,72}

There is still uncertainty as to which patients should undergo immediate surgery and which patients may be successfully treated nonoperatively. There have been 3 different patient responses after ACL injury described: (1) a copers can return to the preinjury level without surgery and subjective instability, (2) an adapter reduces his/her level of activity to avoid subjective instability, and (3) a noncopers cannot return to preinjury activity level because of subjective instability and episodes of giving way.⁵⁸ A screening tool to differentiate potential copers from noncopers was developed and included a combination of hop tests, questionnaires on general knee function, and the frequency of giving-way episodes.^{18,57} Patients categorized as potential copers thereafter participated in structured progressive rehabilitation with additional perturbation training.^{9,19} Regardless of this 3-response concept, there is a strong historical view that the treatment approach should be determined through a shared decision-making process between the patient and the provider.⁸ In particular, the physician should share information on the evidence-based treatment options while also considering the patient's expectations and goals. While the patient and provider are the primary stakeholders in the shared decision-making process, the potential influence of secondary stakeholders, such as family and coaches, should be anticipated so as to minimize interests potentially conflicting with the health of the patient.

Taken as a whole, the current body of evidence regarding the treatment of ACL injury is to a large extent based on low-level evidence. Therefore, an international, multidisciplinary group of experts was assembled to develop expert- and evidence-based consensus statements to assist clinicians in managing this difficult abnormality. The purpose of this article is to report the results of the consensus group addressing the best available evidence on operative versus nonoperative treatment of ACL injury that were developed at the ACL Consensus Meeting Panther Symposium 2019.

*Address correspondence to Volker Musahl, MD, University of Pittsburgh Medical Center, 3200 South Water Street, Pittsburgh, PA 15203, USA (email: musahlv@upmc.edu).

All authors are listed in the Authors section at the end of the article.

This article has been copublished in *Knee Surgery, Sports Traumatology, Arthroscopy*; *Journal of ISAKOS*; and *British Journal of Sports Medicine*. Minor differences exist between this version and others to be consistent with *OJSM* editorial style.

Final revision submitted March 9, 2020; accepted April 3, 2020.

One or more of the authors has declared the following potential conflict of interest or source of funding: O.R.A. has received honoraria from DJO. J.W.X. has received consulting fees from Arthrex and Trice Medical, nonconsulting fees from Arthrex, and royalties from Arthrex. F.H.F. has received educational support and hospitality payments from Smith & Nephew. V.M. has received consulting fees and speaking fees from Smith & Nephew.

Contributing authors have the following disclosures: C.H.B. has received hospitality payments from Smith & Nephew. D.L.J. has received consulting fees, nonconsulting fees, and royalties from Smith & Nephew. J.A.M. has received nonconsulting fees from Arthrex and Smith & Nephew and honoraria from the Musculoskeletal Transplant Foundation and Vericel. K.O. has received educational support from Arthrex. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.




Figure 1. ACL Consensus Meeting Panther Symposium 2019. ACL, anterior cruciate ligament.

METHODS

An international and multidisciplinary group of experts on ACL injury, including orthopaedic surgeons, sports medicine physicians, physical therapists, and scientists, were convened in a 1-year consensus-building effort, which culminated in the consensus meeting, at the University of Pittsburgh, Pittsburgh, Pennsylvania, USA (Figure 1).

The symposium included experts from 18 countries, spanning 6 continents. Experts were assigned to 1 or more of the 3 consensus groups defined by a specific subtopic within ACL injury. The operative versus nonoperative treatment consensus group consisted of 34 participants. A modified Delphi method was used to develop the consensus statements.

The scientific organizing committee and session chairs proposed a series of statements on the basis of a literature review. These were drafted with the aim of addressing areas of current controversy within the treatment of ACL injury, intended to assist clinicians in the management of

this injury. Before the meeting, the proposed statements were presented to the panelists via a web-based survey. Each panelist indicated the extent of agreement or disagreement with each statement and was asked to provide comments on each statement. On the third day of the ACL Consensus Meeting Panther Symposium 2019, after 2 days of presentations by symposium delegates on current knowledge, a consensus discussion was held.

A total of 13 statements on the operative versus nonoperative treatment of ACL injury were discussed. The session was moderated by 2 experts (L.E. and A.D.L.). Initial results and comments from the web-based survey were presented for each statement followed by discussion, debate, and revision by the working group. Consensus was determined by show of hands. Satisfactory consensus was defined as 80% agreement. Opposing views were documented and discussed. Statements with less than 80% agreement were included in the consensus document, noting the percentage of agreement. Statements felt to be irrelevant or redundant were excluded from this final article.

TABLE 2
Consensus Statements on Nonoperative and Operative Treatment of ACL Injury^a

Agreed Statements		Agreement, %
1	Operative and nonoperative treatments are both acceptable treatment options for ACL injury.	100.0
2	Operative versus nonoperative treatment should be reached via a shared decision-making process that considers the patient's presentation, goals, and expectations as well as a balanced presentation of the available evidence-based literature.	82.6
3	The (injury) status of other stabilizing and supporting structures (eg, menisci, other ligaments, cartilage) affects the decision to pursue operative or nonoperative treatment.	100.0
4	Individual anatomic differences (eg, tibial slope, femoral morphology, alignment, etc) may affect the stability of the knee after ACL injury and should be considered in the decision-making process for operative versus nonoperative treatment.	95.7
5	After an ACL injury, patients may be offered a period of progressive rehabilitation to improve impairments and improve overall function.	100.0
6	Patients presenting with instability in their desired activity despite optimal rehabilitation should be referred for operative treatment.	100.0
7	Development of osteoarthritis after an ACL injury is multifactorial, and evidence is inconclusive after operative or nonoperative treatment.	100.0
8	In active patients wishing to return to jumping, cutting, and pivoting sports (eg, soccer, football, handball, basketball), operative treatment is the preferred option to maintain athletic participation in the medium to long term (1- \geq 5 years after injury).	100.0
9	In active patients wishing to return to jumping, cutting, and pivoting sports (eg, soccer, football, handball, basketball), return to cutting and pivoting sports without surgery places the knee at risk of secondary injury (meniscus, cartilage, etc).	100.0
11	In active patients wishing to return to straight-plane activities (eg, running, cycling, swimming, weight lifting, etc), nonoperative treatment is an option.	100.0
12	In active patients wishing to return to straight-plane activities (eg, running, cycling, swimming, weight lifting, etc), in the case of persistent instability in daily life, operative treatment is appropriate for a return to nonrotational activities.	100.0
Not Agreed Statement		Agreement, %
10	In active patients wishing to return to cutting and pivoting sports (eg, soccer, football, handball, basketball), delayed operative treatment may be an option for temporary return to athletic participation after nonoperative treatment, accepting the risk of additional injury.	43.4

^aACL, anterior cruciate ligament.

This consensus group was assigned 2 liaisons (T.D. and B.B.R.), who were responsible for amending each statement as requested over the course of the discussion. Liaisons transcribed the discussion and subsequently completed a literature review of MEDLINE for each finalized statement. To reduce the potential for bias in the data analysis and/or literature review, liaisons did not submit answers to the online questionnaire, nor did they partake in the voting process.

RESULTS

Of the 13 statements discussed by this working group, 9 achieved unanimous consensus, 2 achieved nonunanimous consensus, 1 did not achieve consensus, and 1 was excluded because of redundancy in the information provided (Table 2). The 12 finalized statements, with supporting literature, are as follows.

Operative and nonoperative treatments are both acceptable treatment options for ACL injury (agree 23/23; 100%)

After ACL injury, some patients are able to regain good functional knee stability after nonoperative treatment entailing progressive rehabilitation and are able to return to preinjury sport activity level without an ACL reconstruction

(copers),^{27,28} but the identification of these patients has been challenging.⁷⁷ In a prospective study, the combination of hop tests, muscle strength, subjective instability (episodes of giving way), and knee function was found to be a moderate predictive tool for identification of potential copers.^{18,28,32,57} A randomized controlled trial comparing operative and nonoperative treatment in 121 young, active, nonelite patients with isolated ACL tears demonstrated no superiority of either treatment with regard to patient-reported outcomes at 2- and 5-year follow-up.^{21,22} However, almost 40% of the patients who were initially assigned to the nonoperative treatment group required delayed ACL reconstruction, and 32% of the patients (29 menisci in 19 patients) had subsequent surgery for meniscal injuries during the 2-year follow-up period. In contrast, 34 patients (56%) who underwent early ACL reconstruction also had meniscal treatment (24 partial resection, 10 fixation) simultaneously with the ACL reconstruction, but only 10% (6 meniscal injuries in 5 patients) in the operatively treated group had meniscal injuries that required surgical treatment during follow-up.²¹ With regard to knee laxity, as measured by KT-1000 arthrometer and pivot-shift test, nonoperative treatment resulted in a larger anterior tibial translation (9.0 vs 6.6 mm, respectively) and higher rate of rotatory laxity (positive pivot shift: 78% vs 25%, respectively). A matched-pair study based on the Swedish National ACL

Registry comparing operative and nonoperative treatment after ACL injury reported superior results for quality of life, knee function, and symptoms at 1, 2, and 5 years' follow-up for ACL reconstruction compared with nonoperative treatment.³⁸ Another prospective trial with highly active patients included 832 patients at baseline with subacute ACL tear, whereas 345 patients were initially screened for possibility of nonoperative treatment. Based on the results of various hop tests, subjective instability, and general knee function, 146 patients were classified as potential copers; at final follow-up after 10 years, only 25 patients had not undergone ACL reconstruction.³²

CONCLUSION. Operative and nonoperative are both acceptable treatment options after ACL injury, and a decision based on concomitant injuries, risk factors, level of activity, and patient's expectations and goals is recommended, as demonstrated in the following statements.

Operative versus nonoperative treatment should be reached via a shared decision-making process that considers the patient's presentation, goals, and expectations as well as a balanced presentation of the available evidence-based literature (agree 19/23; 82.6%)

Before a particular treatment approach is pursued, the provider (physician and/or physical therapist) should present the evidence for operative and nonoperative treatment options for an ACL injury to the patient. Based on the patient's activity level, goals, and expectations, a decision should be made with the patient (and parents/guardians for minors) and provider as the primary stakeholders.⁸ Physicians and physical therapists must be aware that personal and situational factors such as level of competition, time in season, playing status, and role in the team could affect the injured athletes' treatment decision. Parents and coaches are often the first people from whom athletes seek support or advice.⁵⁶ However, the coach may be conflicted by the interests of the team and the athlete's immediate and future health.^{20,83} For some athletes, reactions and comments of parents related to the athlete's injury were reported to negatively affect the athlete's treatment decision, with pressure to return to sport.⁵⁶ Because of the possible conflict of interest, secondary stakeholders such as family, coaches, and agents, among others, should not be directly involved in the decision-making process, although their indirect involvement may be considered.

CONCLUSION. Shared decision-making of the treatment option should be based on the evidence for operative and nonoperative treatments as well as patient's expectations and goals with the provider and patient as the primary stakeholders.

The (injury) status of other stabilizing and supporting structures (eg, menisci, other ligaments, cartilage) affects the decision to pursue operative or nonoperative treatment (agree 23/23; 100%)

ACL injuries often occur together with concomitant injury to other knee structures, with meniscal injuries reported in

23% to 42%, cartilage lesions in 27%, and combined meniscal and chondral lesions in 15% of cases (Figure 2).^{6,11,38}

However, most studies investigating nonoperative ACL treatment or studies comparing nonoperative and operative treatment are limited to isolated ACL tears.^{21,22,32} Based on clinical and biomechanical studies, an ACL reconstruction with concomitant meniscal repair may restore knee kinematics and results in improved patient-reported outcomes at short- and long-term follow-up.^{44,66,70,89} In contrast, simultaneously performed meniscectomy with ACL reconstruction is associated with poorer clinical outcome, inferior knee kinematics, and a high rate (48%-100%) of osteoarthritis in the long-term follow-up.^{12,30,47,50,88} In case of delayed ACL reconstruction, a meniscectomy is more often performed than a meniscal repair.³⁷ The presence of concomitant knee injuries should therefore always be considered in the decision-making process, given the worse outcomes for meniscal injuries with delayed ACL reconstruction and higher rate of osteoarthritis in the long-term follow-up. In case of concomitant meniscal injury repair, anatomic ACL reconstruction with additional treatment of the meniscal injury is recommended.

In case of multiple ligament injuries involving the ACL and at least 1 other ligament, the literature has consistently demonstrated that operative management is superior to nonoperative management.^{42,63,68} Based on a recent systematic review, early (within 3 weeks after injury) reconstruction in a multiple ligament-injured knee was superior to delayed reconstruction with regard to clinical outcome measurements (Lysholm score: 90 vs 82, respectively, of 100 points) and resulted in higher rate of excellent/good International Knee Documentation Committee (IKDC) scores (47% vs 31%, respectively).⁴² Although failure after ligament reconstruction is not consistently defined in the literature (ie, the need for revision vs objective laxity vs rerupture on imaging vs Knee injury and Osteoarthritis Outcome Score [KOOS] <44), the failure rate in a multiple ligament-injured knee is lower for reconstruction (6%-9%) compared with repair techniques (37%-40%).^{41,75}

CONCLUSION. The presence of a repairable meniscal lesion or a multiple ligament injury is an indication for an early anatomic ACL reconstruction with concomitant treatment of the other injured structures (meniscal repair, ligament repair/augmentation).

Individual anatomic differences (eg, tibial slope, femoral morphology, alignment, etc) may affect the stability of the knee after ACL injury and should be considered in the decision-making process for operative versus nonoperative treatment (agree 22/23; 95.7%)

Bony morphology and soft tissue injury patterns have been demonstrated to influence knee joint laxity. An increased posterior tibial slope is associated with increased anterior tibial translation as well as with increased rotatory instability (Figure 3).^{67,85}

In addition, an increased lateral femoral condyle ratio resulted in increased rotatory instability.^{64,65} Severe varus limb alignment (>5°) was demonstrated to increase the risk




Figure 2. As seen in T2-weighted magnetic resonance imaging sequences, the patient sustained a (A) complete anterior cruciate ligament rupture and (B) associated lateral meniscus root tear.




Figure 3. (A, B) Posterior tibial slope varies among patients, with greater slope increasing the risk of failure after anterior cruciate ligament (ACL) reconstruction. (C, D) Notch dimension varies among patients, with small notch width dimensions constituting a relative contraindication for double-bundle ACL reconstruction.

for more rapid degeneration of the medial compartment in the ACL-deficient knee and is also a risk factor for secondary failure after an ACL reconstruction.^{59,79} Whereas lateral meniscal tears and a complete lateral meniscectomy result in increased rotatory instability,^{31,52} a complete medial meniscectomy more strongly affects anterior tibial translation. However, general joint laxity (Beighton hypermobility score >4) is not associated with increased rotatory laxity in the ACL-deficient knee.⁷⁶

CONCLUSION. Bony morphology features (increased posterior tibial slope, severe varus limb alignment, etc) and concomitant injuries associated with increased or persistent knee instability should be considered in the decision-making process and are a relative indication for operative treatment.

After an ACL injury, patients may be offered a period of progressive rehabilitation to improve impairments and improve overall function (agree 23/23; 100%)

Knee joint effusion, limited range of motion, and decreased quadriceps strength in the injured leg are common impairments initially after an ACL injury.^{10,45} Effusion can limit quadriceps function and in turn affect knee joint mechanics.⁶⁰ Progressive rehabilitation is useful in treating these initial impairments.³² In patients with the possibility for nonoperative treatment (absence of concomitant meniscal injuries or multiligament injuries requiring surgical treatment), before the evaluation of knee instability, a phase of rehabilitation is recommended to treat the initial impairments. Afterward, evaluation by hop

tests and assessment of strength, overall knee function, and subjective instability is recommended to quantify the patients' potential for nonoperative treatment. If progressive rehabilitation does not provide a satisfactory outcome, then operative intervention needs to be pursued, and the progressive rehabilitation will have enhanced the postoperative outcome.¹⁴ In a cohort study with 2187 patients after resolution of impairments, 1 group was treated with neuromuscular training (ie, strengthening and neuromuscular training) before ACL reconstruction and was compared with immediate ACL reconstruction. At 2 years' follow-up, preoperative progressive rehabilitation before ACL reconstruction resulted in better patient-reported outcome (KOOS and IKDC) compared with ACL reconstruction without preoperative rehabilitation.¹⁴ Whereas 63% of the patients without preoperative rehabilitation returned to sport at 2 years' follow-up, which is similar to the reported rate (65%) in a meta-analysis from 2016, the rate increased to 72% in the group that completed preoperative rehabilitation.¹⁴

CONCLUSION. Preoperative resolution of impairments and a period of rehabilitation are recommended for operative and nonoperative treatments.

Patients presenting with instability in their desired activity despite optimal rehabilitation should be referred for operative treatment (agree 23/23; 100%)

Persistent instability is a risk factor for further damage to the meniscus and cartilage.³³ Although the definitions of recurrent instability and episodes of instability vary in the current literature, a correlation between persistent and recurrent instability after ACL injury and meniscal and cartilage lesions has been demonstrated in several studies.^{2,36,74} In a cohort study of 62 patients with acute ACL reconstructions, 37 with subacute ACL reconstructions, and 36 with chronic ACL reconstructions, 1 episode of giving way was associated with 3-fold higher odds for lateral meniscal tears. Timing of surgery and episodes of instability influenced the incidence of lateral meniscal tears, with 1.45 higher odds in subacute (6-12 weeks) ACL reconstruction and 2.82 higher odds in chronic (>12 weeks) ACL reconstruction.² Moreover, frequent episodes of instability are correlated with medial meniscal tears and chondral injuries.³⁶ Chondral defects and meniscectomy have been demonstrated as predictive factors for the development of osteoarthritis after ACL reconstruction.^{15,35}

A partial ACL injury progressed to a complete ACL tear in 39% of young active patients treated nonoperatively, with half of the complete tears presenting with a concomitant meniscal lesion at the time of reconstruction. Age ≤ 20 years and participation in pivoting contact sports were identified as significant risk factors for progression to a complete tear.¹⁶

CONCLUSION. If patient-reported instability or severe episodes of giving way occur during the progressive rehabilitation, patients should be referred for anatomic ACL reconstruction.

Development of osteoarthritis after an ACL injury is multifactorial, and evidence is inconclusive after operative or nonoperative treatment (agree 23/23; 100%)

Osteoarthritis is the most common joint disease, affecting not only the cartilage but all other tissues of the joint as well.²⁴ The pathomechanism of PTOA has not been fully elucidated, but based on current research, the process of development of osteoarthritis is multifactorial.²⁴ Injuries, like ACL ruptures, can affect the joint biomechanics and cause chondral and meniscal lesions and thereby reduce the sustainability of the joint. Matrix metalloproteinases are responsible for cartilage destruction and synovial inflammation and have been shown to be elevated after ACL injury and reconstruction.^{78,84} A meta-analysis of 24 observational studies found a 4-fold increased risk for PTOA after knee injuries, although the definition of an injury was largely heterogeneous among the analyzed studies.⁵³ After ACL injury, the prevalence of PTOA is increased after both operative and nonoperative treatments as compared with those without injury.^{17,48,55,69} Based on a recent systematic review with 41 included studies, the rate of osteoarthritis after ACL reconstruction varied between 1% and 80%, with meniscectomy as the consistent risk factor for the development of osteoarthritis.⁴³ Although long-term outcome studies after ACL reconstruction are available, the technique has evolved in recent years, with a shift from non-anatomic ACL reconstruction to anatomic ACL reconstruction, limiting conclusions on the possible protective effect of anatomic ACL reconstruction.

CONCLUSION. Osteoarthritis after ACL injury is seen after both operative and nonoperative treatments. Therefore, there is still a need for prospective randomized controlled trials to evaluate the hypothesized preventive effect of anatomic ACL reconstruction on the development of PTOA.

In active patients wishing to return to jumping, cutting, and pivoting sports (eg, soccer, football, handball, basketball), operative treatment is the preferred option to maintain athletic participation in the medium to long term (1- ≥ 5 years after injury) (agree 23/23; 100%)

In active patients wishing to return to pivoting and cutting sports, ACL reconstruction is the preferred treatment option to maintain participation in the medium to long term. However, overall, only 65% of patients return to their preinjury sport level after ACL reconstruction, and only 55% return to competitive-level sports.⁴ Although the exact reasons are still unknown, younger age, male sex, professional sport level, and positive psychological response were demonstrated to be associated with a successful return to preinjury sport level after ACL reconstruction. In general, elite athletes return to their preinjury level of sport after ACL reconstruction more often than recreational athletes.^{39,87} For instance, over 90% of elite soccer players were reported to return to the preinjury level after ACL reconstruction.⁸⁷ Similarly, in a recent systematic review, the return-to-sport rate in elite football and basketball players was 78% and 82%, respectively.³⁹ In contrast, only 12.8% of

high-level athletes returned to the preinjury sport level with nonoperative treatment, with a high rate of secondary meniscal and cartilage damage; after 20 years, 95% of the patients underwent meniscectomy, during which 68% of patients were found to have chondral lesions.^{18,54} Overall, athletes returned to their preinjury sport level between 6 and 13 months after ACL reconstruction.³⁹

CONCLUSION. In active patients, anatomic ACL reconstruction is the preferred treatment because of the higher rate of return to preinjury sport level.

In active patients wishing to return to jumping, cutting, and pivoting sports (eg, soccer, football, handball, basketball), return to cutting and pivoting sports without surgery places the knee at risk of secondary injury (meniscus, cartilage, etc) (agree 23/23; 100%)

In a prospective randomized controlled trial, patients with high activity level (median Tegner activity score of 9) with isolated ACL tears received early operative treatment or nonoperative treatment with the option of delayed ACL reconstruction. Although no differences were evident for patient-reported outcomes, at 2-year follow-up, patients in the “optional” operative treatment group had more self-reported and clinical laxity of the involved knee and more meniscal surgery over a 5-year follow-up period.²¹ In a separate cohort, the risk for sustaining at least 1 additional intra-articular injury increased by 0.6% with each month of delay in operative treatment.⁷ The odds of secondary cartilage lesions increased by nearly 1% for each month of delay.²⁵ A delay in ACL reconstruction of at least 12 months almost doubled the risk for meniscal tears.^{7,40} Increased risk of secondary injury is especially noted in young (<12 years) and skeletally immature patients.²

CONCLUSION. Nonoperative treatment increases the risk for secondary injuries if the patient wants to return to jumping, cutting, and pivoting sports because of the increased risk of further episodes of instability.

In active patients wishing to return to cutting and pivoting sports (eg, soccer, football, handball, basketball), delayed operative treatment may be an option for temporary return to athletic participation after nonoperative treatment, accepting the risk of additional injury (agree 10/23; 43.4%)

No consensus was reached for this statement. Some professional athletes and active patients want to delay ACL reconstruction to temporarily return to athletic participation (competition). Based on current evidence, the risk of secondary damage to the knee (eg, meniscus, cartilage) is high, especially in high-demand sports with jumping, cutting, and pivoting. In a recent cross-sectional study, 860 patients were included, with 47.2% being professional athletes. With regard to prevalence of meniscal tears, medial, lateral, and combined lesions were found more often with increasing time from injury (TFI) to surgery (with TFI of 0-36 weeks, medial meniscal tear prevalence was 48.2%, and with TFI of >61 weeks, it was 59.3%). Not only did the prevalence of injury increase with time, the rate of

meniscectomy also increased (with TFI of 0-36 weeks, medial meniscectomy rate was 7.5%, and with TFI of >61 weeks, it was 12.8%).⁷³

CONCLUSION. Delayed ACL reconstruction in active patients may be a treatment option, but the provider, as well as the patient, must be aware of the risk of secondary injuries with worse long-term outcomes.

In active patients wishing to return to straight-plane activities (eg, running, cycling, swimming, weight lifting, etc), nonoperative treatment is an option (agree 23/23; 100%)

Straight-plane activities are less demanding on the ligamentous stabilizers of the knee and therefore are amenable to nonoperative treatment. The anteroposterior stability during straight-plane activities might be maintained by muscular control, but coronal and rotational stability could not be compensated.⁸⁶ With specific neuromuscular training (perturbation training) additional to standard rehabilitation, unphysiological muscular co-contractions during walking can be minimized and normalize the knee kinematics in the ACL-deficient knee.⁹ In a matched-pair study, nonoperative treatment resulted in an earlier return (3-4 months vs 6-12 months, respectively) and a higher return to level II sports (88.9% vs 77.8%, respectively) as compared with operative treatment.²⁸ Another study demonstrated a significantly higher number of nonoperatively treated patients returned to level II and level III sports compared with operative treatment.²⁷

CONCLUSION. For return to straight-plane activities, nonoperative treatment is an option.

In active patients wishing to return to straight-plane activities (eg, running, cycling, swimming, weight lifting, etc), in the case of persistent instability in daily life, operative treatment is appropriate for a return to nonrotational activities (agree 23/23; 100%)

Straight-plane activities are less demanding to the ligamentous stabilizers of the knee and are therefore amenable to nonoperative treatment. If during the nonoperative treatment subjective instability persists or episodes of giving way occur, referral for consideration of anatomic ACL reconstruction is recommended.^{21,49} Moreover, current evidence for the efficacy of nonoperative treatment is limited to isolated ACL tears.

CONCLUSION. Based on current evidence, persistent instability in activities of daily living is an indication for anatomic ACL reconstruction to restore knee laxity and prevent secondary injuries.

CONCLUSION

The expert panel at the ACL Consensus Meeting Panther Symposium 2019 reached consensus, defined as >80% agreement, on 11 of 12 statements in terms of operative versus nonoperative treatment for ACL injuries. Consensus was reached that both treatment options may be acceptable, depending on patient characteristics, including the

type of sporting demands and the presence of concomitant injuries. In highly active patients engaged in jumping, cutting, and pivoting sports, early anatomic ACL reconstruction is recommended because of the high risk of secondary meniscal and cartilage injuries with delayed surgery, although a period of progressive rehabilitation to resolve impairments and improve neuromuscular function may be recommended. For patients who want to return to straight-plane activities, nonoperative treatment with structured, progressive rehabilitation is an acceptable treatment option. However, with persistent functional instability, or when episodes of giving way occur, anatomic ACL reconstruction is indicated.

Despite strong consensus by experts, there is a need for larger randomized trials with longer term follow-up in which early surgery (followed by rehabilitation) is compared with a strategy of early rehabilitation and delayed surgery. There are insufficient data to guide treatment in instances when there are concomitant meniscal and collateral ligament injuries. Data on long-term clinical outcomes are needed to better understand the effect of ACL treatment on injuries, subsequent injuries to menisci and cartilage, and the development of osteoarthritis.

AUTHORS

Theresa Diermeier, MD (Department of Orthopaedic Surgery, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA); Department of Orthopaedic Sports Medicine, Technical University of Munich, Munich, Germany); Benjamin B. Rothrauff, MD, PhD (Department of Orthopaedic Surgery, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA); Lars Engebretsen, MD, PhD (Department of Orthopaedic Surgery, Oslo University Hospital, Oslo, Norway); Andrew D. Lynch, PhD, PT (Department of Orthopaedic Surgery, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA); Olufemi R. Ayeni, MD, PhD (Division of Orthopaedic Surgery, McMaster University, Hamilton, Ontario, Canada); Mark V. Paterno, PT, PhD (Division of Sports Medicine, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA); John W. Xerogeanes, MD (Emory University School of Medicine, Atlanta, Georgia, USA); Freddie H. Fu, MD, DSc, DPs (Hon) (Department of Orthopaedic Surgery, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA); Jon Karlsson, MD, PhD (Department of Orthopaedics, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden); Volker Musahl, MD (Department of Orthopaedic Surgery, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA); and the Panther Symposium ACL Treatment Consensus Group: Charles H. Brown Jr, MD (International Knee and Joint Centre, Abu Dhabi, UAE); Terese L. Chmielewski, PT, PhD, SCS (TRIA Orthopaedic Center, Bloomington, Minnesota, USA); Mark Clatworthy, MBChB, FRACS (Department of Orthopaedic Surgery, Middlemore Hospital, Auckland, New Zealand); Stefano Della Villa, MD (Education Research Department, Isokinetic Medical Group,

Bologna, Italy); Lucio Ernlund, MD, MSc (Instituto de Joelho e Ombro, Curitiba, Paraná, Brasil); Christian Fink, MD, Prof. (Gelenkpunkt-Sports and Joint Surgery Innsbruck, Innsbruck, Austria; Research Unit for Orthopaedic Sports Medicine and Injury Prevention [OSMI], UMIT, Hall in Tirol, Austria); Alan Getgood, MD, FRCS(Tr&Orth), DipSEM (Fowler Kennedy Sport Medicine Clinic London, London, Ontario, Canada); Timothy E. Hewett, PhD (Mayo Clinic, Rochester, Minnesota, USA; The Ohio State University, Columbus, Ohio, USA); Yasuyuki Ishibashi, MD (Department of Orthopaedic Surgery, Hirosaki University Graduate School of Medicine, Hirosaki, Aomori, Japan); Darren L. Johnson, MD (Department of Orthopaedic Surgery and Sports Medicine, University of Kentucky, Lexington, Kentucky, USA); Jeffrey A. Macalena, MD (Department of Orthopedic Surgery, University of Minnesota, Minneapolis, Minnesota, USA); Robert G. Marx, MD, MSc, FRCSC (Medicine Institute, Hospital for Special Surgery, New York, USA); Jacques Menetrey, MD, Prof. (University Hospitals of Geneva, Geneva, Switzerland); Sean J. Meredith, MD (Department of Physical Therapy, University of Delaware, Newark, Delaware, USA); Kentaro Onishi, DO (Department of Physical Medicine and Rehabilitation, University of Pittsburgh, Pittsburgh, Pennsylvania, USA); Thomas Rauer, MD (Department of Trauma Surgery, University Hospital Zurich, Zurich, Switzerland); Benjamin B. Rothrauff, MD, PhD (UPMC Freddie Fu Sports Medicine Center, Department of Orthopaedic Surgery, University of Pittsburgh, Pittsburgh, Pennsylvania, USA); Laura C. Schmitt, PT, PhD (Division of Physical Therapy, School of Health and Rehabilitation Sciences, The Ohio State University, Columbus, Ohio, USA); Romain Seil, MD, Prof. (Department of Orthopaedic Surgery, Clinique d'Eich-Centre Hospitalier de Luxembourg, Luxembourg); Eric H. Senorski, PT, PhD, MSc (Department of Health and Rehabilitation, Institute of Neuroscience and Physiology, University of Gothenburg, Gothenburg, Sweden); Rainer Siebold, MD (ATOS Clinic, Heidelberg, Germany); Lynn Snyder-Mackler, PT, ScD, FAPTA (Department of Physical Therapy, University of Delaware, Newark, Delaware, USA); Tim Spalding, MD, FRCS(Tr&Orth) (University Hospitals Coventry & Warwickshire, Coventry, UK); Eleonore Svantesson, MD (Department of Orthopaedics, Institute of Clinical Sciences, University of Gothenburg, Gothenburg, Sweden); Kevin E. Wilk, PT, DPT, FAPTA (Champion Sports Medicine, American Sports Medicine Institute, Birmingham, Alabama, USA).

REFERENCES

1. Alentorn-Geli E, Myer GD, Silvers HJ, et al. Prevention of non-contact anterior cruciate ligament injuries in soccer players, part 1: mechanisms of injury and underlying risk factors. *Knee Surg Sports Traumatol Arthrosc.* 2009;17:705-729.
2. Anderson AF, Anderson CN. Correlation of meniscal and articular cartilage injuries in children and adolescents with timing of anterior cruciate ligament reconstruction. *Am J Sports Med.* 2015;43:275-281.
3. Ardern CL, Österberg A, Tagesson S, Gauffin H, Webster KE, Kvist J. The impact of psychological readiness to return to sport and

- recreational activities after anterior cruciate ligament reconstruction. *Br J Sports Med.* 2014;48:1613-1619.
4. Ardern CL, Taylor NF, Feller JA, Webster KE. Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *Br J Sports Med.* 2014;48:1543.
 5. Boden BP, Dean GS, Feagin JA Jr, Garrett WE Jr. Mechanisms of anterior cruciate ligament injury. *Orthopedics.* 2000;23:573-578.
 6. Borchers JR, Kaeding CC, Pedroza AD, et al. Intra-articular findings in primary and revision anterior cruciate ligament reconstruction surgery: a comparison of the MOON and MARS study groups. *Am J Sports Med.* 2011;39:1889-1893.
 7. Brambilla L, Pulici L, Carimati G, et al. Prevalence of associated lesions in anterior cruciate ligament reconstruction: correlation with surgical timing and with patient age, sex, and body mass index. *Am J Sports Med.* 2015;43:2966-2973.
 8. Charles C, Gafni A, Whelan T. Shared decision-making in the medical encounter: what does it mean? (or it takes at least two to tango). *Soc Sci Med.* 1997;44:681-692.
 9. Chmielewski TL, Hurd WJ, Rudolph KS, Axe MJ, Snyder-Mackler L. Perturbation training improves knee kinematics and reduces muscle co-contraction after complete unilateral anterior cruciate ligament rupture. *Phys Ther.* 2005;85:740-749.
 10. Chmielewski TL, Stackhouse S, Axe MJ, Snyder-Mackler L. A prospective analysis of incidence and severity of quadriceps inhibition in a consecutive sample of 100 patients with complete acute anterior cruciate ligament rupture. *J Orthop Res.* 2004;22:925-930.
 11. Cimino PM. The incidence of meniscal tears associated with acute anterior cruciate ligament disruption secondary to snow skiing accidents. *Arthroscopy.* 1994;10(2):198-200.
 12. Cohen M, Amaro JT, Eijnsman B, et al. Anterior cruciate ligament reconstruction after 10 to 15 years: association between meniscectomy and osteoarthritis. *Arthroscopy.* 2007;23:629-634.
 13. Eitzen I, Moksnes H, Snyder-Mackler L, Risberg MA. A progressive 5-week exercise therapy program leads to significant improvement in knee function early after anterior cruciate ligament injury. *J Orthop Sports Phys Ther.* 2010;40:705-721.
 14. Failla MJ, Logerstedt DS, Grindem H, et al. Does extended preoperative rehabilitation influence outcomes 2 years after ACL reconstruction? A comparative effectiveness study between the MOON and Delaware-Oslo ACL cohorts. *Am J Sports Med.* 2016;44:2608-2614.
 15. Fairbank TJ. Knee joint changes after meniscectomy. *J Bone Joint Surg Br.* 1948;30-B:664-670.
 16. Fayard JM, Sonnery-Cottet B, Vrgoc G, et al. Incidence and risk factors for a partial anterior cruciate ligament tear progressing to a complete tear after nonoperative treatment in patients younger than 30 years. *Orthop J Sports Med.* 2019;7:2325967119856624.
 17. Fithian DC, Paxton EW, Stone ML, et al. Prospective trial of a treatment algorithm for the management of the anterior cruciate ligament-injured knee. *Am J Sports Med.* 2005;33:335-346.
 18. Fitzgerald GK, Axe MJ, Snyder-Mackler L. A decision-making scheme for returning patients to high-level activity with nonoperative treatment after anterior cruciate ligament rupture. *Knee Surg Sports Traumatol Arthrosc.* 2000;8:76-82.
 19. Fitzgerald GK, Axe MJ, Snyder-Mackler L. The efficacy of perturbation training in nonoperative anterior cruciate ligament rehabilitation programs for physically active individuals. *Phys Ther.* 2000;80:128.
 20. Flint FA, Weiss MR. Returning injured athletes to competition: a role and ethical dilemma. *Can J Sport Sci.* 1992;17(1):34-40.
 21. Frobell R, Roos E, Roos H, Ranstam J, Lohmander L. A randomized trial of treatment for acute anterior cruciate ligament tears. *N Engl J Med.* 2010;363(4):331-342.
 22. Frobell RB, Roos HP, Roos EM, Roemer FW, Ranstam J, Lohmander LS. Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. *BMJ.* 2013;346:F232.
 23. Fu FH, van Eck CF, Tashman S, Irrgang JJ, Moreland MS. Anatomic anterior cruciate ligament reconstruction: a changing paradigm. *Knee Surg Sports Traumatol Arthrosc.* 2015;23:640-648.
 24. Glyn-Jones S, Palmer AJ, Agricola R, et al. Osteoarthritis. *Lancet.* 2015;386(9991):376-387.
 25. Granan L-P, Bahr R, Lie SA, Engebretsen L. Timing of anterior cruciate ligament reconstructive surgery and risk of cartilage lesions and meniscal tears: a cohort study based on the Norwegian National Knee Ligament Registry. *Am J Sports Med.* 2009;37:955-961.
 26. Granan L-P, Bahr R, Steindal K, Furnes O, Engebretsen L. Development of a national cruciate ligament surgery registry: the Norwegian National Knee Ligament Registry. *Am J Sports Med.* 2007;36:308-315.
 27. Grindem H, Eitzen I, Engebretsen L, Snyder-Mackler L, Risberg MA. Nonsurgical or surgical treatment of ACL injuries: knee function, sports participation, and knee reinjury. The Delaware-Oslo ACL cohort study. *J Bone Joint Surg Am.* 2014;96:1233-1241.
 28. Grindem H, Eitzen I, Moksnes H, Snyder-Mackler L, Risberg MA. A pair-matched comparison of return to pivoting sports at 1 year in anterior cruciate ligament-injured patients after a nonoperative versus an operative treatment course. *Am J Sports Med.* 2012;40:2509-2516.
 29. Hangalur G, Brenneman E, Nicholls M, Bakker R, Laing A, Chandrasekar N. Can a knee brace reduce the strain in the anterior cruciate ligament? A study using combined in vivo/in vitro method. *Prosthet Orthot Int.* 2016;40:394-399.
 30. Hertel P, Behrend H, Cierpinski T, Musahl V, Widjaja G. ACL reconstruction using bone-patellar tendon-bone press-fit fixation: 10-year clinical results. *Knee Surg Sports Traumatol Arthrosc.* 2005;13:248-255.
 31. Hoshino Y, Miyaji N, Nishida K, et al. The concomitant lateral meniscus injury increased the pivot shift in the anterior cruciate ligament-injured knee. *Knee Surg Sports Traumatol Arthrosc.* 2019;27(2):646-651.
 32. Hurd WJ, Axe MJ, Snyder-Mackler L. A 10-year prospective trial of a patient management algorithm and screening examination for highly active individuals with anterior cruciate ligament injury, part 2: determinants of dynamic knee stability. *Am J Sports Med.* 2008;36:48-56.
 33. Jonsson H, Riklund-Ahlstrom K, Lind J. Positive pivot shift after ACL reconstruction predicts later osteoarthritis: 63 patients followed 5-9 years after surgery. *Acta Orthop Scand.* 2004;75:594-599.
 34. Kaeding CC, Leger-St-Jean B, Magnussen RA. Epidemiology and diagnosis of anterior cruciate ligament injuries. *Clin Sports Med.* 2017;36:1-8.
 35. Keays SL, Newcombe PA, Bullock-Saxton JE, Bullock MI, Keays AC. Factors involved in the development of osteoarthritis after anterior cruciate ligament surgery. *Am J Sports Med.* 2010;38:455-463.
 36. Kluczynski MA, Marzo JM, Bisson LJ. Factors associated with meniscal tears and chondral lesions in patients undergoing anterior cruciate ligament reconstruction: a prospective study. *Am J Sports Med.* 2013;41(12):2759-2765.
 37. Krutsch W, Zellner J, Baumann F, Pfeifer C, Nerlich M, Angele P. Timing of anterior cruciate ligament reconstruction within the first year after trauma and its influence on treatment of cartilage and meniscus pathology. *Knee Surg Sports Traumatol Arthrosc.* 2017;25:418-425.
 38. Kvist J, Kartus J, Karlsson J, Forsblad M. Results from the Swedish National Anterior Cruciate Ligament Register. *Arthroscopy.* 2014;30:803-810.
 39. Lai CCH, Ardern CL, Feller JA, Webster KE. Eighty-three per cent of elite athletes return to preinjury sport after anterior cruciate ligament reconstruction: a systematic review with meta-analysis of return to sport rates, graft rupture rates and performance outcomes. *Br J Sports Med.* 2018;52:128.
 40. Lawrence JT, Argawal N, Ganley TJ. Degeneration of the knee joint in skeletally immature patients with a diagnosis of an anterior cruciate ligament tear: is there harm in delay of treatment? *Am J Sports Med.* 2011;39:2582-2587.
 41. Levy BA, Dajani KA, Morgan JA, Shah JP, Dahm DL, Stuart MJ. Repair versus reconstruction of the fibular collateral ligament and posterolateral corner in the multiligament-injured knee. *Am J Sports Med.* 2010;38(4):804-809.

42. Levy BA, Dajani KA, Whelan DB, et al. Decision making in the multiligament-injured knee: an evidence-based systematic review. *Arthroscopy*. 2009;25:430-438.
43. Lie MM, Risberg MA, Storheim K, Engebretsen L, Oiestad BE. What's the rate of knee osteoarthritis 10 years after anterior cruciate ligament injury? An updated systematic review. *Br J Sports Med*. 2019;53(18):1162-1167.
44. Lorbach O, Kieb M, Domnick C, et al. Biomechanical evaluation of knee kinematics after anatomic single- and anatomic double-bundle ACL reconstructions with medial meniscal repair. *Knee Surg Sports Traumatol Arthrosc*. 2015;23:2734-2741.
45. Lynch AD, Logerstedt DS, Axe MJ, Snyder-Mackler L. Quadriceps activation failure after anterior cruciate ligament rupture is not mediated by knee joint effusion. *J Orthop Sports Phys Ther*. 2012;42:502-510.
46. Lynch TS, Parker RD, Patel RM, Andrich JT, Spindler KP; MOON Group. The impact of the Multicenter Orthopaedic Outcomes Network (MOON) research on anterior cruciate ligament reconstruction and orthopaedic practice. *J Am Acad Orthop Surg*. 2015;23:154.
47. Magnussen RA, Mansour AA, Carey JL, Spindler KP. Meniscus status at anterior cruciate ligament reconstruction associated with radiographic signs of osteoarthritis at 5- to 10-year follow-up: a systematic review. *J Knee Surg*. 2009;22:347-357.
48. Meuffels DE, Favejee MM, Vissers MM, Heijboer MP, Reijman M, Verhaar JA. Ten year follow-up study comparing conservative versus operative treatment of anterior cruciate ligament ruptures: a matched-pair analysis of high level athletes. *Br J Sports Med*. 2009;43:347-351.
49. Meunier A, Odensten M, Good L. Long-term results after primary repair or non-surgical treatment of anterior cruciate ligament rupture: a randomized study with a 15-year follow-up. *Scand J Med Sci Sports*. 2007;17(3):230-237.
50. Musahl V, Citak M, O'Loughlin PF, Choi D, Bedi A, Pearle AD. The effect of medial versus lateral meniscectomy on the stability of the anterior cruciate ligament-deficient knee. *Am J Sports Med*. 2010;38:1591-1597.
51. Musahl V, Kopf S, Rabuck S, et al. Rotatory knee laxity tests and the pivot shift as tools for ACL treatment algorithm. *Knee Surg Sports Traumatol Arthrosc*. 2012;20:793-800.
52. Musahl V, Rahnama-Azar AA, Costello J, et al. The influence of meniscal and anterolateral capsular injury on knee laxity in patients with anterior cruciate ligament injuries. *Am J Sports Med*. 2016;44:3126-3131.
53. Muthuri SG, McWilliams DF, Doherty M, Zhang W. History of knee injuries and knee osteoarthritis: a meta-analysis of observational studies. *Osteoarthritis Cartilage*. 2011;19:1286-1293.
54. Nebelung W, Wuschech H. Thirty-five years of follow-up of anterior cruciate ligament-deficient knees in high-level athletes. *Arthroscopy*. 2005;21:696-702.
55. Neuman P, Englund M, Kostogiannis I, Friden T, Roos H, Dahlberg LE. Prevalence of tibiofemoral osteoarthritis 15 years after nonoperative treatment of anterior cruciate ligament injury: a prospective cohort study. *Am J Sports Med*. 2008;36:1717-1725.
56. Nippert AH, Smith AM. Psychologic stress related to injury and impact on sport performance. *Phys Med Rehabil Clin N Am*. 2008;19:399-418.
57. Noyes FR, Barber SD, Mangine RE. Abnormal lower limb symmetry determined by function hop tests after anterior cruciate ligament rupture. *Am J Sports Med*. 1991;19:513-518.
58. Noyes FR, Matthews DS, Moar PA, Grood ES. The symptomatic anterior cruciate-deficient knee, part II: the results of rehabilitation, activity modification, and counseling on functional disability. *J Bone Joint Surg Am*. 1983;65:163-174.
59. Noyes FR, Schipplein OD, Andriacchi TP, Saddemi SR, Weise M. The anterior cruciate ligament-deficient knee with varus alignment: an analysis of gait adaptations and dynamic joint loadings. *Am J Sports Med*. 1992;20:707-716.
60. Palmieri-Smith RM, Kreinbrink J, Ashton-Miller JA, Wojtyl EM. Quadriceps inhibition induced by an experimental knee joint effusion affects knee joint mechanics during a single-legged drop landing. *Am J Sports Med*. 2007;35:1269-1275.
61. Paterno MV. Non-operative care of the patient with an ACL-deficient knee. *Curr Rev Musculoskelet Med*. 2017;10:322-327.
62. Pauenberger L, Syre S, Schurz M. "Ligamentization" in hamstring tendon grafts after anterior cruciate ligament reconstruction: a systematic review of the literature and a glimpse into the future. *Arthroscopy*. 2013;29:1712-1721.
63. Peskun CJ, Whelan DB. Outcomes of operative and nonoperative treatment of multiligament knee injuries: an evidence-based review. *Sports Med Arthrosc Rev*. 2011;19:167-173.
64. Pfeiffer TR, Burnham JM, Hughes JD, et al. An increased lateral femoral condyle ratio is a risk factor for anterior cruciate ligament injury. *J Bone Joint Surg Am*. 2018;100:857-864.
65. Pfeiffer TR, Burnham JM, Kanakamedala AC, et al. Distal femur morphology affects rotatory knee instability in patients with anterior cruciate ligament ruptures. *Knee Surg Sports Traumatol Arthrosc*. 2019;27(5):1514-1519.
66. Phillips M, Rönnblad E, Lopez-Rengstig L, et al. Meniscus repair with simultaneous ACL reconstruction demonstrated similar clinical outcomes as isolated ACL repair: a result not seen with meniscus resection. *Knee Surg Sports Traumatol Arthrosc*. 2018;26:2270-2277.
67. Rahnama-Azar AA, Abebe ES, Johnson P, et al. Increased lateral tibial slope predicts high-grade rotatory knee laxity pre-operatively in ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2017;25:1170-1176.
68. Richter M, Bosch U, Wippermann B, Hofmann A, Krettek C. Comparison of surgical repair or reconstruction of the cruciate ligaments versus nonsurgical treatment in patients with traumatic knee dislocations. *Am J Sports Med*. 2002;30:718-727.
69. Sanders TL, Kremers HM, Bryan AJ, et al. Is anterior cruciate ligament reconstruction effective in preventing secondary meniscal tears and osteoarthritis? *Am J Sports Med*. 2016;44:1699-1707.
70. Sarraj M, Coughlin RP, Solow M, et al. Anterior cruciate ligament reconstruction with concomitant meniscal surgery: a systematic review and meta-analysis of outcomes. *Knee Surg Sports Traumatol Arthrosc*. 2019;27(11):3441-3452.
71. Scheffler SU, Unterhauser FN, Weiler A. Graft remodeling and ligamentization after cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2008;16:834-842.
72. Sitler M, Ryan CJ, Hopkinson LW, et al. The efficacy of a prophylactic knee brace to reduce knee injuries in football: a prospective, randomized study at West Point. *Am J Sports Med*. 1990;18:310-315.
73. Sommerfeldt M, Goodine T, Raheem A, Whittaker J, Otto D. Relationship between time to ACL reconstruction and presence of adverse changes in the knee at the time of reconstruction. *Orthop J Sports Med*. 2018;6(12):2325967118813917.
74. Sommerfeldt M, Raheem A, Whittaker J, Hui C, Otto D. Recurrent instability episodes and meniscal or cartilage damage after anterior cruciate ligament injury: a systematic review. *Orthop J Sports Med*. 2018;6(7):2325967118786507.
75. Stannard JP, Brown SL, Farris RC, McGwin G Jr, Volgas DA. The posterolateral corner of the knee: repair versus reconstruction. *Am J Sports Med*. 2005;33(6):881-888.
76. Sundemo D, Blom A, Hoshino Y, et al. Correlation between quantitative pivot shift and generalized joint laxity: a prospective multicenter study of ACL ruptures. *Knee Surg Sports Traumatol Arthrosc*. 2018;26:2362-2370.
77. Thoma LM, Grindem H, Logerstedt D, et al. Coper classification early after anterior cruciate ligament rupture changes with progressive neuromuscular and strength training and is associated with 2-year success: the Delaware-Oslo ACL Cohort Study. *Am J Sports Med*. 2019;47(4):807-814.
78. Troeberg L, Nagase H. Proteases involved in cartilage matrix degradation in osteoarthritis. *Biochim Biophys Acta*. 2012;1824:133-145.
79. van de Pol GJ, Arnold MP, Verdonshot N, van Kampen A. Varus alignment leads to increased forces in the anterior cruciate ligament. *Am J Sports Med*. 2008;37:481-487.
80. van Eck CF, Gravare-Silbernagel K, Samuelsson K, et al. Evidence to support the interpretation and use of the Anatomic Anterior Cruciate

- Ligament Reconstruction Checklist. *J Bone Joint Surg Am.* 2013;95:e153.
81. van Eck CF, Lesniak BP, Schreiber VM, Fu FH. Anatomic single- and double-bundle anterior cruciate ligament reconstruction flowchart. *Arthroscopy.* 2010;26:258-268.
 82. van Yperen DT, Reijman M, van Es EM, Bierma-Zeinstra SMA, Meuffels DE. Twenty-year follow-up study comparing operative versus nonoperative treatment of anterior cruciate ligament ruptures in high-level athletes. *Am J Sports Med.* 2018;46(5):1129-1136.
 83. Vergeer I, Hogg JM. Coaches' decision policies about the participation of injured athletes in competition. *Sport Psychol.* 1999;13:42-56.
 84. Vincent TL. Targeting mechanotransduction pathways in osteoarthritis: a focus on the pericellular matrix. *Curr Opin Pharmacol.* 2013;13(3):449-454.
 85. Voos JE, Suero EM, Citak M, et al. Effect of tibial slope on the stability of the anterior cruciate ligament-deficient knee. *Knee Surg Sports Traumatol Arthrosc.* 2012;20:1626-1631.
 86. Waite JC, Beard DJ, Dodd CAF, Murray DW, Gill HS. In vivo kinematics of the ACL-deficient limb during running and cutting. *Knee Surg Sports Traumatol Arthrosc.* 2005;13:377-384.
 87. Walden M, Hagglund M, Magnusson H, Ekstrand J. Anterior cruciate ligament injury in elite football: a prospective three-cohort study. *Knee Surg Sports Traumatol Arthrosc.* 2011;19:11-19.
 88. Wu WH, Hackett T, Richmond JC. Effects of meniscal and articular surface status on knee stability, function, and symptoms after anterior cruciate ligament reconstruction: a long-term prospective study. *Am J Sports Med.* 2002;30:845-850.
 89. Zaffagnini S, Signorelli C, Grassi A. Anatomic anterior cruciate ligament reconstruction using hamstring tendons restores quantitative pivot shift. *Orthop J Sports Med.* 2018;6(12):2325967118812364.
 90. Zantop T, Herbert M, Raschke MJ, Fu FH, Petersen W. The role of the anteromedial and posterolateral bundles of the anterior cruciate ligament in anterior tibial translation and internal rotation. *Am J Sports Med.* 2007;35:223-227.