



OPEN

DATA DESCRIPTOR

Thirty complete *Streptomyces* genome sequences for mining novel secondary metabolite biosynthetic gene clusters

Namil Lee^{1,6}, Woori Kim^{1,6}, Soonkyu Hwang¹, Yongjae Lee¹, Suhjung Cho¹, Bernhard Palsson ^{3,4,5} & Byung-Kwan Cho ^{1,2,5} ✉

Streptomyces are Gram-positive bacteria of significant industrial importance due to their ability to produce a wide range of antibiotics and bioactive secondary metabolites. Recent advances in genome mining have revealed that *Streptomyces* genomes possess a large number of unexplored silent secondary metabolite biosynthetic gene clusters (smBGCs). This indicates that *Streptomyces* genomes continue to be an invaluable source for new drug discovery. Here, we present high-quality genome sequences of 22 *Streptomyces* species and eight different *Streptomyces venezuelae* strains assembled by a hybrid strategy exploiting both long-read and short-read genome sequencing methods. The assembled genomes have more than 97.4% gene space completeness and total lengths ranging from 6.7 to 10.1 Mbp. Their annotation identified 7,000 protein coding genes, 20 rRNAs, and 68 tRNAs on average. *In silico* prediction of smBGCs identified a total of 922 clusters, including many clusters whose products are unknown. We anticipate that the availability of these genomes will accelerate discovery of novel secondary metabolites from *Streptomyces* and elucidate complex smBGC regulation.

Background & Summary

With the rapid emergence of antibiotic microbial resistance (AMR) to all major classes of antibiotics and the decline in number of potential candidates for new antibiotics, there is a pressing need for the discovery of novel antibacterial compounds¹. *Streptomyces*, soil dwelling gram-positive bacteria, continue to be promising microorganisms for the production of clinically important secondary metabolites, including not only antibiotics, but also antiviral, antifungal, and antiparasitic agents, and antitumorals and immunosuppressant compounds². *Streptomyces* are distinguished by their complex life cycle and high G + C content (often over 70%) in their linear genomes^{3,4}. Traditionally, drug discovery from *Streptomyces* has been based on bioactivity screening followed by mass spectrometry and NMR-based molecular identification⁵. However, recent advances in genomics-based approaches revealed that most of the secondary metabolite biosynthetic gene clusters (smBGCs) of streptomycetes are inactive under laboratory conditions, suggesting that the ability of streptomycetes to produce secondary metabolites has been under-estimated^{5,6}. Each *Streptomyces* species has the genetic potential to produce more than 30 secondary metabolites on average, which are diverse and differ between species^{7,8}. Considering *Streptomyces* is the largest genus of actinobacteria with approximately 900 species characterized so far, streptomycetes are a valuable resource for the discovery of novel secondary metabolites⁹.

SmBGCs, especially polyketide and non-ribosomal peptide synthetase types, are often composed of extraordinarily long genes (>5 kb) encoding multi-modular enzymes with repetitive domain structures. Therefore, accurate gene annotations based on high quality genome sequences are essential for the precise identification of smBGCs¹⁰. Gene annotation with the high quality genome of *S. clavuligerus* revealed that 30% out of a total

¹Department of Biological Sciences and KI for the BioCentury, Korea Advanced Institute of Science and Technology, Daejeon, 34141, Republic of Korea. ²Intelligent Synthetic Biology Center, Daejeon, 34141, Republic of Korea. ³Department of Bioengineering, University of California San Diego, La Jolla, CA, 92093, USA. ⁴Department of Pediatrics, University of California San Diego, La Jolla, CA, 92093, USA. ⁵Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Lyngby, 2800, Denmark. ⁶These authors contributed equally: Namil Lee and Woori Kim. ✉e-mail: bcho@kaist.ac.kr




Fig. 1 Quality of the genome sequencing data. **(a)** Distribution of Illumina reads quality based on Phred score. **(b)** Read quality distribution of PacBio reads. Black line indicates total number of bases in the reads which have greater read quality than the corresponding read quality value on x-axis.

of 7,163 protein coding genes were incorrectly annotated in the previous draft genome of *S. clavuligerus* containing ambiguous and inaccurate nucleotides, indicating the importance of high quality genome sequences¹¹. In addition, high quality genome sequences are essential for multi-omics analysis, which facilitates the understanding of the complex regulation on smBGCs and rational engineering for increasing secondary metabolites production^{11,12}.

Among the 1,614 streptomycetes genomes that have been deposited in the NCBI Assembly database to date (as of 9th December 2019), only 189 and 35 assemblies were designated as complete genome level and chromosome level, respectively. More than 86% of assemblies were draft-quality genome sequences, which contain fragmented multiple contigs or ambiguous sequences^{4,13–15}. One of the main obstacles to obtaining high quality genomic information of streptomycetes is the low fidelity of sequencing techniques when dealing with high G w C genomes and frequently repetitive sequences such as terminal inverted repeats¹³. In addition, since streptomycetes have linear chromosome, it is difficult to confirm the completeness of the assembled chromosome.

In this study, we present the high-quality genome sequences of 30 streptomycetes, increasing the total number of reported complete *Streptomyces* genome by about 10%. The target streptomycetes were 22 *Streptomyces* type strains and eight different *Streptomyces venezuelae* strains, most of which are currently used as industrial strains for producing various bioactive compounds. We applied hybrid assembly strategy with long-read (PacBio) and short-read (Illumina) sequencing techniques to obtain complete genome sequences. PacBio sequencing provides long reads of several kb in length which allows the readthrough of regions with low complexity, enabling the assembly of repetitive regions, which are difficult to assemble by using Illumina sequencing reads, even with the high coverage data¹⁶. However, Illumina sequencing provides reads with a lower error rate compared to the PacBio sequencing, and assembled contigs based on the Illumina sequencing reads are not simply a subset of the contigs from PacBio sequencing reads^{13,17}. Therefore, reconciling PacBio and Illumina sequencing methods enables one to generate more complete genomes by overcoming the shortcomings of each method. During

No.	Species	Final scaffolds (No.)	Scaffold length before correction (bp)	Mapped Illumina reads (%)	Conflict positions (No.)	Added bases (No.)	Deleted bases (No.)	Scaffold length after correction (bp)	G + C contents (%)	Assembly accession number
1	<i>Streptomyces clavuligerus</i>	2	6,748,589 and 1,795,496	71.16 and 14.03	7	4	3	6,748,591 and 1,795,495	72.5	GCA_005519465.1
2	<i>Streptomyces tsukubaensis</i>	1	7,963,727	95.13	15	15	0	7,963,742	71.9	GCA_003932715.1
3	<i>Streptomyces galilaeus</i>	1	7,756,176	90.56	51	34	16	7,756,194	71.4	GCA_008704575.1
4	<i>Streptomyces nitrosporeus</i>	1	7,581,543	93.50	51	35	16	7,581,562	72.2	GCA_008704555.1
5	<i>Streptomyces subrutillus</i>	1	7,604,705	96.41	286	269	0	7,604,974	73.4	GCA_008704535.1
6	<i>Streptomyces viridosporus</i> T7A	1	7,280,447	90.44	90	89	0	7,280,536	72.6	GCA_008704515.1
7	<i>Streptomyces kanamyceticus</i>	1	10,133,525	99.09	376	375	3	10,133,897	71.0	GCA_008704495.1
8	<i>Streptomyces aureofaciens</i>	1	7,757,873	84.86	16	9	5	7,757,877	72.6	GCA_008704475.1
9	<i>Streptomyces prasinus</i>	1	7,646,576	89.70	1,025	1,021	5	7,647,592	72.0	GCA_008704445.1
10	<i>Streptomyces fradiae</i>	1	6,725,574	97.63	5	5	0	6,725,579	74.7	GCA_008704425.1
11	<i>Streptomyces alboniger</i>	1	7,962,594	99.12	193	193	1	7,962,786	71.2	GCA_008704395.1
12	<i>Streptomyces coeruleorubidus</i>	1	9,334,399	99.67	1,297	1,299	0	9,335,698	71.1	GCA_008705135.1
13	<i>Streptomyces cinereoruber</i>	1	7,516,474	99.74	178	178	0	7,516,652	72.9	GCA_009299385.1
14	<i>Streptomyces nodosus</i>	1	7,772,564	99.51	26	25	2	7,772,587	70.9	GCA_008704995.1
15	<i>Streptomyces vinaceus</i>	1	7,673,329	92.46	180	180	0	7,673,509	72.3	GCA_008704935.1
16	<i>Streptomyces platensis</i>	1	8,500,673	99.75	354	352	13	8,501,012	71.1	GCA_008704855.1
17	<i>Streptomyces spectabilis</i>	1	9,806,222	95.30	934	938	0	9,807,160	72.4	GCA_008704795.1
18	<i>Streptomyces chartreusis</i>	1	9,911,637	98.42	461	461	0	9,912,098	71.0	GCA_008704715.1
19	<i>Streptomyces rimosus</i>	1	9,361,132	96.22	22	22	0	9,361,154	72.0	GCA_008704655.1
20	<i>Streptomyces albofaciens</i>	2	4,757,761 and 4,494,336	53.36 and 45.53	504	501	3	4,757,978 and 4,494,617	72.3	GCA_008634025.1
21	<i>Streptomyces filamentosus</i>	2	5,742,252 and 2,129,928	75.22 and 24.28	3,218	3,228	1	5,744,022 and 2,131,385	73.6	GCA_008634015.1
22	<i>Streptomyces venezuelae</i> ATCC 10712	1	8,223,439	99.84	96	81	15	8,223,505	72.5	GCA_008639165.1
23	<i>Streptomyces venezuelae</i> ATCC 21113	1	7,893,622	99.85	173	181	0	7,893,803	72.5	GCA_008639045.1
24	<i>Streptomyces venezuelae</i> ATCC 10595	1	7,871,449	95.50	35	34	3	7,871,480	72.5	GCA_008705255.1
25	<i>Streptomyces venezuelae</i> ATCC 15068	1	8,557,615	99.71	587	587	0	8,558,202	71.9	GCA_008642375.1
26	<i>Streptomyces venezuelae</i> ATCC 14583	1	8,018,461	87.17	29	27	4	8,018,484	71.3	GCA_008642355.1
27	<i>Streptomyces venezuelae</i> ATCC 14584	1	8,941,823	99.00	255	255	0	8,942,078	71.2	GCA_008642315.1
28	<i>Streptomyces venezuelae</i> ATCC 14585	1	8,048,139	82.34	64	41	26	8,048,154	71.3	GCA_008642335.1
29	<i>Streptomyces venezuelae</i> ATCC 21782	1	7,525,235	90.50	87	87	0	7,525,322	71.9	GCA_008642295.1
30	<i>Streptomyces venezuelae</i> ATCC 21018	1	7,746,214	91.61	59	57	4	7,746,267	72.1	GCA_008642275.1

Table 1. The statistics of genome assembly and correction.

the genome assembly using reads from PacBio (0.46–5.18 Gbp) and Illumina (0.5–3.0 Gbp) sequencing, we constructed 6.7 to 10.1 Mbp of streptomycetes genomes, most of which consist of single chromosomes with 72% G + C contents on average. Inaccurate sequences in the assembled genome were corrected using Illumina sequencing reads. The complete streptomycetes genomes have more than 97.4% gene space completeness and on average 7,000 protein coding genes, 20 rRNAs, and 68 tRNAs were annotated. Finally, based on the complete genome sequences and annotations, we predicted a total of 922 smbGCs. The complete genome sequences and newly determined smbGCs in this study should prove to be a fundamental resource for understanding the genetic basis of streptomycetes and for discovering novel secondary metabolites.

Methods

Genomic DNA (gDNA) extraction. Total 30 streptomycetes were purchased from Korean Collection for Type Cultures (KCTC, Korea). A stock of streptomycetes were inoculated to 50 mL of liquid culture medium with 0.16 g mL⁻¹ of glass beads (3 ± 0.3 mm diameter) in 250 mL baffled flask and grown at 30 °C in a 200 rpm orbital shaker. Each streptomycetes was grown in one of four different culture medium, R5(-) medium (25 mM TES (pH 7.2), 103 g L⁻¹ sucrose, 1% (w/v) glucose, 5 g L⁻¹ yeast extract, 10.12 g L⁻¹ MgCl₂·6H₂O, 0.25 g L⁻¹ K₂SO₄, 0.1 g L⁻¹ casamino acids, 0.08 g L⁻¹ ZnCl₂, 0.4 mg L⁻¹ FeCl₃, 0.02 mg L⁻¹ CuCl₂·2H₂O, 0.02 mg L⁻¹ MnCl₂·4H₂O, 0.02 mg L⁻¹ Na₂B₄O₇·10H₂O, and 0.02 mg L⁻¹ (NH₄)₆Mo₇O₂₄·4H₂O), 1 × sporulation medium (3.33 g L⁻¹ glucose,




Fig. 2 Genome assembly of 30 streptomycetes. **(a)** Strategy for genome assembly and corrections. **(b)** Profile of Illumina reads mapped on assembled genomes. Data were visualized using SignalMap (Roche NimbleGen, Inc.). Red line indicates the average Illumina read coverage of all genomic positions.

1 g L⁻¹ yeast extract, 1 g L⁻¹ beef extract, 2 g L⁻¹ tryptose, and 0.006 g L⁻¹ FeSO₄·7H₂O), YEME medium (340 g L⁻¹ sucrose, 10 g L⁻¹ glucose, 3 g L⁻¹ yeast extract, 5 g L⁻¹ bacto peptone, and 3 g L⁻¹ oxioid malt extract), and MYM medium (4 g L⁻¹ maltose, 4 g L⁻¹ yeast extract, 10 g L⁻¹ malt extract). For gDNA extraction, 25 mL cultured cells were harvested at the exponential growth phase and washed twice with same volume of 10 mM EDTA, followed by the lysozyme (10 mg mL⁻¹) treatment at 37 °C for 45 min. gDNA was extracted using a Wizard Genomic DNA Purification Kit (Promega, Madison, WI, USA) according to the manufacturer's instruction. Quality and quantity

No.	Species	Complete and single-copy	Complete and duplicated	Fragmented	Missing	Total	Gene space completeness (%)
1	<i>Streptomyces clavuligerus</i>	343	0	0	9	352	97.4
2	<i>Streptomyces tsukubaensis</i>	350	0	0	2	352	99.4
3	<i>Streptomyces galilaeus</i>	351	0	0	1	352	99.7
4	<i>Streptomyces nitrosporeus</i>	352	0	0	0	352	100.0
5	<i>Streptomyces subrutilus</i>	349	0	0	3	352	99.1
6	<i>Streptomyces viridosporus</i> T7A	351	0	0	1	352	99.7
7	<i>Streptomyces kanamyceticus</i>	352	0	0	0	352	100.0
8	<i>Streptomyces aureofaciens</i>	350	0	0	2	352	99.4
9	<i>Streptomyces prasinus</i>	350	0	0	2	352	99.4
10	<i>Streptomyces fradiae</i>	351	0	0	1	352	99.7
11	<i>Streptomyces alboniger</i>	351	0	0	1	352	99.7
12	<i>Streptomyces coeruleorubidus</i>	351	0	0	1	352	99.7
13	<i>Streptomyces cinereoruber</i>	351	0	0	1	352	99.7
14	<i>Streptomyces nodosus</i>	350	0	1	1	352	99.4
15	<i>Streptomyces vinaceus</i>	349	0	1	2	352	99.1
16	<i>Streptomyces platensis</i>	351	0	0	1	352	99.7
17	<i>Streptomyces spectabilis</i>	350	0	1	1	352	99.4
18	<i>Streptomyces chartreusis</i>	351	0	0	1	352	99.7
19	<i>Streptomyces rimosus</i>	351	0	0	1	352	99.7
20	<i>Streptomyces albofaciens</i>	346	4	0	2	352	99.4
21	<i>Streptomyces filamentosus</i>	351	0	0	1	352	99.7
22	<i>Streptomyces venezuelae</i> ATCC 10712	352	0	0	0	352	100.0
23	<i>Streptomyces venezuelae</i> ATCC 21113	352	0	0	0	352	100.0
24	<i>Streptomyces venezuelae</i> ATCC 10595	352	0	0	0	352	100.0
25	<i>Streptomyces venezuelae</i> ATCC 15068	351	0	0	1	352	99.7
26	<i>Streptomyces venezuelae</i> ATCC 14583	351	0	0	1	352	99.7
27	<i>Streptomyces venezuelae</i> ATCC 14584	351	0	0	1	352	99.7
28	<i>Streptomyces venezuelae</i> ATCC 14585	351	0	0	1	352	99.7
29	<i>Streptomyces venezuelae</i> ATCC 21782	349	0	0	3	352	99.1
30	<i>Streptomyces venezuelae</i> ATCC 21018	350	0	0	2	352	99.4

Table 2. Gene space completeness of completed genomes.

of extracted gDNA samples were evaluated using 1% agarose gel electrophoresis and Nanodrop (Thermo Fisher Scientific, Waltham, MA, USA), respectively.

Short-read (Illumina) genome sequencing. For construction of short-read genome sequencing library, 2.5 µg of gDNA was sheared to approximately 350 bp by a Covaris instrument (Covaris Inc., Woburn, MA, USA) with the following conditions; Power 175, Duty factor 20%, C. burst 200, Time 23 s, 8 times. The library was constructed using a TruSeq DNA PCR-Free LT kit (Illumina Inc., San Diego, CA, USA) following manufacturer's instruction. Briefly, the fragmented DNA samples were cleaned and end-repaired, followed by the adaptor ligation and bead-based size selection ranging from 400 to 500 bp. Quantity of final libraries was measured using Qubit® dsDNA HS Assay Kit (Thermo Fisher Scientific) and the library size was determined using Agilent 2200 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA). Among the constructed sequencing libraries, 29 libraries were sequenced with the HiSeq. 2500 (Illumina Inc.) as 100 bp single-end reads and remaining one library for *S. tsukubaensis* was sequenced with the Miseq v.2 (Illumina Inc.) with 50 bp single-read recipe. Finally, 0.46 to 5.18 Gbp of raw sequence data were obtained and the read qualities were examined by creating sequencing QC reports function of CLC genomic workbench version 6.5.1 (CLC bio, Denmark) (Online-only Table 1 and Fig. 1a).

Long-read (PacBio) genome sequencing. A total of 5 µg gDNA was used as input for PacBio genome sequencing library preparation. The sequencing library was constructed with the PacBio SMRTbell™ Template Prep Kit (Pacific Biosciences, Menlo Park, CA, USA) following manufacturer's instructions. Fragments smaller than 20 kbp were removed using the Blue Pippin Size selection system (Sage Science, Beverly, MA, USA) and the constructed libraries were validated using Agilent 2100 Bioanalyzer (Agilent Technologies). Final SMRTbell libraries were sequenced using one or two SMRT cells with P6-C4-chemistry (DNA Sequencing Reagent 4.0) on the PacBio RS II sequencing platform (Pacific Biosciences). Approximately, 0.5 to 3.0 Gbp of raw sequence data were generated (Online-only Table 1).

Genome assembly. Among the raw PacBio sequencing reads, only the reads with a read quality value greater than 0.75 and a length longer than 50 bp were filtered (Fig. 1b). Post filtered reads were assembled by

No.	Species	CDS (No.)	16s rRNA (No.)	tRNA (No.)	Genome accession number	BioProject accession number
1	<i>Streptomyces clavuligerus</i>	6,880	18	66	CP027858	PRJNA414136
2	<i>Streptomyces tsukubaensis</i>	6,376	18	66	CP020700	PRJNA382016
3	<i>Streptomyces galilaeus</i>	6,725	18	76	CP023703	PRJNA412292
4	<i>Streptomyces nitrosporeus</i>	6,364	18	74	CP023702	PRJNA412292
5	<i>Streptomyces subruttilus</i>	6,431	21	68	CP023701	PRJNA412292
6	<i>Streptomyces viridosporus</i> T7A	6,211	18	70	CP023700	PRJNA412292
7	<i>Streptomyces kanamyceticus</i>	8,384	18	66	CP023699	PRJNA412292
8	<i>Streptomyces aureofaciens</i>	6,453	33	71	CP023698	PRJNA412292
9	<i>Streptomyces prasinus</i>	6,263	18	68	CP023697	PRJNA412292
10	<i>Streptomyces fradiae</i>	5,465	18	65	CP023696	PRJNA412292
11	<i>Streptomyces alboniger</i>	6,613	18	67	CP023695	PRJNA412292
12	<i>Streptomyces coeruleorubidus</i>	8,058	18	67	CP023694	PRJNA412292
13	<i>Streptomyces cinereoruber</i>	6,392	18	69	CP023693	PRJNA412292
14	<i>Streptomyces nodosus</i>	6,491	18	68	CP023747	PRJNA412292
15	<i>Streptomyces vinaceus</i>	6,603	21	68	CP023692	PRJNA412292
16	<i>Streptomyces platensis</i>	7,032	21	67	CP023691	PRJNA412292
17	<i>Streptomyces spectabilis</i>	8,212	18	65	CP023690	PRJNA412292
18	<i>Streptomyces chartreusis</i>	8,396	18	71	CP023689	PRJNA412292
19	<i>Streptomyces rimosus</i>	7,756	21	68	CP023688	PRJNA412292
20	<i>Streptomyces albofaciens</i>	7,520	21	67	PDCM00000000	PRJNA412292
21	<i>Streptomyces filamentosus</i>	6,832	24	70	PDCL00000000	PRJNA412292
22	<i>Streptomyces venezuelae</i> ATCC 10712	7,377	21	67	CP029197	PRJNA454547
23	<i>Streptomyces venezuelae</i> ATCC 21113	6,987	21	67	CP029196	PRJNA454547
24	<i>Streptomyces venezuelae</i> ATCC 10595	6,942	21	67	CP029195	PRJNA454547
25	<i>Streptomyces venezuelae</i> ATCC 15068	7,700	21	69	CP029194	PRJNA454547
26	<i>Streptomyces venezuelae</i> ATCC 14583	7,154	18	66	CP029193	PRJNA454547
27	<i>Streptomyces venezuelae</i> ATCC 14584	7,832	18	65	CP029192	PRJNA454547
28	<i>Streptomyces venezuelae</i> ATCC 14585	7,096	18	66	CP029191	PRJNA454547
29	<i>Streptomyces venezuelae</i> ATCC 21782	6,655	18	69	CP029190	PRJNA454547
30	<i>Streptomyces venezuelae</i> ATCC 21018	6,769	21	71	CP029189	PRJNA454547

Table 3. Summary of genome annotation.

the hierarchical genome assembly process workflow (HGAP, Version 2.3), including consensus polishing with Quiver¹⁸. For each assembled contig, error correction was performed based on their estimated genome size and average coverage. Raw reads from the Illumina sequencing were quality trimmed using CLC genomic workbench version 6.5.1 (ambiguous limit 2 and quality limit 0.05) and assembled using *de novo* assembly function of CLC genomic workbench version 6.5.1 with default parameters. To expand the assembled contigs, all of assembled PacBio and Illumina contigs were aligned using MAUVE 2.4.0¹⁹ and linked using GAP5 program (Staden package)²⁰.

Genome correction. Quality trimmed Illumina sequencing reads were mapped to the assembled genome using CLC genomic workbench version 6.5.1 (mismatch cost 2, insertion cost 3, deletion cost 3, length fraction 0.9, and similarity fraction 0.9). Conflicts showing more than 80% frequency for Illumina reads were corrected as Illumina sequence (Table 1). In addition, percentage of mapped Illumina reads on to the assembled genome represents degree of completeness (Table 1 and Fig. 2b). Completeness of gene space was estimated using the BUSCO v3 (Table 2)²¹.

Genome annotation and secondary metabolite biosynthetic gene cluster prediction. The complete genome sequences of streptomycetes were submitted to the NCBI GenBank database and annotated by the latest updated version of NCBI Prokaryotic Genome Annotation Pipeline (PGAP)²². Using the GenBank formatted files of each genomes as input, secondary metabolite biosynthetic gene clusters were predicted by antiSMASH 4.0²³.

Data Records

Raw reads from short-read (Illumina) and long-read (PacBio) sequencing were deposited in the NCBI Sequence Read Archive (SRA) (Online-only Table 1)^{24,25}. 30 complete genome sequences were deposited in GenBank via the NCBI's submission portal (Table 3)^{26–55}. Detailed information on the predicted 922 smBGCs in 30 streptomycetes genomes has been deposited in FigShare⁵⁶.




Fig. 3 Secondary metabolite biosynthetic gene clusters in 30 complete streptomycetes genomes.

Technical Validation

Streptomyces have drawn considerable attention because of their ability to produce various clinically important secondary metabolites. Total 30 streptomycetes genomes were sequenced by using both PacBio and Illumina sequencing methods to elucidate their biosynthetic potential. After cleaning the reads, on average 98,380 PacBio reads with 11,725 bp length and 18,223,235 Illumina reads with 100 bp length (50 bp for *S. tsukubaensis*) were generated (Fig. 1a,b and Online-only Table 1). Through the assembly of reads from two sequencing platforms using HGAP, CLC workbench, MAUVE, and GAP5 programs, single linear scaffolds ranging from 6.7 to 10.1 Mbp in length with 72% G + C contents were obtained for 27 streptomycetes, whereas two scaffolds were finally constructed for three remaining streptomycetes, *S. clavuligerus* (6.7 and 1.8 Mbp), *S. albofaciens* (4.8 and 4.5 Mbp), and *S. filamentosus* (5.7 and 2.1 Mbp) (Table 1). *S. clavuligerus* has been reported to have a large linear plasmid with a length of 1.8 Mbp, so the genome was correctly assembled into a single chromosome, while the *S. albofaciens* and *S. filamentosus* genomes appear to be assembled into two divided scaffolds^{11,57}. To increase the accuracy of the assembled genome sequences, Illumina sequences showing more than 80% coverage at the conflict sites were taken as the corrected ones (Table 1). Approximately, 96.32% of Illumina sequencing reads were successfully mapped to the corresponding genomes (Table 1 and Fig. 2b). The completeness of the genomes were assessed using the BUSCO approach with a total of 352 orthologue groups from the Actinobacteria Dataset²¹. Results showed that 29 genomes have more than 99.1% gene space completeness and the *S. clavuligerus* genome has 97.4% gene space completeness (Table 2). Following NCBI PGAP, 30 genomes were annotated with 7,000 protein coding genes, 20 rRNAs, and 68 tRNAs on average (Table 3). Finally, based on the annotation, a total of 922 smBGCs were predicted in 30 streptomycetes genomes (Fig. 3). Detailed information, such as genomic positions, types, and putative products of each smBGC are publicly available in Figshare⁵⁶.

Code availability

The version and parameter of all bioinformatics tools used in this work are described in the Methods section.

Received: 11 September 2019; Accepted: 24 January 2020;

Published online: 13 February 2020

References

- Genilloud, O. The re-emerging role of microbial natural products in antibiotic discovery. *Antonie Van Leeuwenhoek* **106**, 173–188 (2014).
- Jones, S. E. & Elliot, M. A. *Streptomyces* exploration: Competition, volatile communication and new bacterial behaviours. *Trends Microbiol* **25**, 522–531 (2017).
- Hopwood, D. A. Soil to genomics: the *Streptomyces* chromosome. *Annu Rev Genet* **40**, 1–23 (2006).
- Lee, N. *et al.* Synthetic biology tools for novel secondary metabolite discovery in *Streptomyces*. *J Microbiol Biotechnol* **29**, 667–686 (2019).
- Ziemert, N., Alanjary, M. & Weber, T. The evolution of genome mining in microbes - a review. *Nat Prod Rep* **33**, 988–1005 (2016).
- Rebets, Y., Brotz, E., Tokovenko, B. & Luzhetskyy, A. Actinomycetes biosynthetic potential: how to bridge *in silico* and *in vivo*? *J Ind Microbiol Biotechnol* **41**, 387–402 (2014).
- Bentley, S. D. *et al.* Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). *Nature* **417**, 141–147 (2002).
- Omura, S. *et al.* Genome sequence of an industrial microorganism *Streptomyces avermitilis*: deducing the ability of producing secondary metabolites. *Proc Natl Acad Sci USA* **98**, 12215–12220 (2001).

9. Nett, M., Ikeda, H. & Moore, B. S. Genomic basis for natural product biosynthetic diversity in the actinomycetes. *Nat Prod Rep* **26**, 1362–1384 (2009).
10. Reva, O. & Tummeler, B. Think big—giant genes in bacteria. *Environ Microbiol* **10**, 768–777 (2008).
11. Hwang, S. *et al.* Primary transcriptome and translome analysis determines transcriptional and translational regulatory elements encoded in the *Streptomyces clavuligerus* genome. *Nucleic Acids Res* **47**, 6114–6129 (2019).
12. Li, Y., Zhang, C., Liu, C., Ju, J. & Ma, J. Genome sequencing of *Streptomyces atratus* SCSIOZH16 and activation production of nocardamine via metabolic engineering. *Front Microbiol* **9**, 1269 (2018).
13. Harrison, J. & Studholme, D. J. Recently published *Streptomyces* genome sequences. *Microb Biotechnol* **7**, 373–380 (2014).
14. Barreiro, C. *et al.* Draft genome of *Streptomyces tsukubaensis* NRRL 18488, the producer of the clinically important immunosuppressant tacrolimus (FK506). *J Bacteriol* **194**, 3756–3757 (2012).
15. Song, J. Y. *et al.* Draft genome sequence of *Streptomyces clavuligerus* NRRL 3585, a producer of diverse secondary metabolites. *J Bacteriol* **192**, 6317–6318 (2010).
16. Koren, S. *et al.* Reducing assembly complexity of microbial genomes with single-molecule sequencing. *Genome Biol* **14**, R101 (2013).
17. Ardui, S., Ameer, A., Vermeesch, J. R. & Hestand, M. S. Single molecule real-time (SMRT) sequencing comes of age: applications and utilities for medical diagnostics. *Nucleic Acids Res* **46**, 2159–2168 (2018).
18. Chin, C. S. *et al.* Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* **10**, 563–569 (2013).
19. Darling, A. C., Mau, B., Blattner, F. R. & Perna, N. T. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* **14**, 1394–1403 (2004).
20. Bonfield, J. K. & Whitwham, A. Gap5—editing the billion fragment sequence assembly. *Bioinformatics* **26**, 1699–1703 (2010).
21. Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210–3212 (2015).
22. Haft, D. H. *et al.* RefSeq: an update on prokaryotic genome annotation and curation. *Nucleic Acids Res* **46**, D851–D860 (2018).
23. Blin, K. *et al.* antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary identification. *Nucleic Acids Res* **45**, W36–W41 (2017).
24. Leinonen, R., Sugawara, H. & Shumway, M. & International Nucleotide Sequence Database, C. The sequence read archive. *Nucleic Acids Res* **39**, D19–21 (2011).
25. NCBI Sequence Read Archive. <https://identifiers.org/ncbi/insdc.sra:SRP200324> (2019).
26. GenBank. <https://identifiers.org/ncbi/insdc:CP020700> (2018).
27. GenBank. <https://identifiers.org/ncbi/insdc:CP023688> (2018).
28. GenBank. <https://identifiers.org/ncbi/insdc:CP023689> (2018).
29. GenBank. <https://identifiers.org/ncbi/insdc:CP023690> (2018).
30. GenBank. <https://identifiers.org/ncbi/insdc:CP023691> (2018).
31. GenBank. <https://identifiers.org/ncbi/insdc:CP023692> (2018).
32. GenBank. <https://identifiers.org/ncbi/insdc:CP023693> (2018).
33. GenBank. <https://identifiers.org/ncbi/insdc:CP023694> (2018).
34. GenBank. <https://identifiers.org/ncbi/insdc:CP023695> (2018).
35. GenBank. <https://identifiers.org/ncbi/insdc:CP023696> (2018).
36. GenBank. <https://identifiers.org/ncbi/insdc:CP023697> (2018).
37. GenBank. <https://identifiers.org/ncbi/insdc:CP023698> (2018).
38. GenBank. <https://identifiers.org/ncbi/insdc:CP023699> (2018).
39. GenBank. <https://identifiers.org/ncbi/insdc:CP023700> (2018).
40. GenBank. <https://identifiers.org/ncbi/insdc:CP023701> (2018).
41. GenBank. <https://identifiers.org/ncbi/insdc:CP023702> (2018).
42. GenBank. <https://identifiers.org/ncbi/insdc:CP023703> (2018).
43. GenBank. <https://identifiers.org/ncbi/insdc:CP023747> (2018).
44. GenBank. <https://identifiers.org/ncbi/insdc:CP029189> (2018).
45. GenBank. <https://identifiers.org/ncbi/insdc:CP029190> (2018).
46. GenBank. <https://identifiers.org/ncbi/insdc:CP029191> (2018).
47. GenBank. <https://identifiers.org/ncbi/insdc:CP029192> (2018).
48. GenBank. <https://identifiers.org/ncbi/insdc:CP029193> (2018).
49. GenBank. <https://identifiers.org/ncbi/insdc:CP029194> (2018).
50. GenBank. <https://identifiers.org/ncbi/insdc:CP029195> (2018).
51. GenBank. <https://identifiers.org/ncbi/insdc:CP029196> (2018).
52. GenBank. <https://identifiers.org/ncbi/insdc:CP029197> (2018).
53. GenBank. <https://identifiers.org/ncbi/insdc:PDCL00000000> (2018).
54. GenBank. <https://identifiers.org/ncbi/insdc:PDCM00000000> (2018).
55. GenBank. <https://identifiers.org/ncbi/insdc:CP027858> (2019).
56. Lee, N. *et al.* Thirty complete *Streptomyces* genome sequences for mining novel secondary metabolite biosynthetic gene clusters. <https://doi.org/10.6084/m9.figshare.c.4823394> (2020).
57. Medema, M. H. *et al.* The sequence of a 1.8-Mb bacterial linear plasmid reveals a rich evolutionary reservoir of secondary metabolic pathways. *Genome Biol Evol* **2**, 212–224 (2010).

Acknowledgements

This work was supported by a grant from the Novo Nordisk Foundation (grant number NNF10CC1016517). This work was also supported by the Bio & Medical Technology Development Program (2018M3A9F3079664 to B.-K.C.) through the National Research Foundation (NRF) funded by the Ministry of Science and ICT (MSIT).

Author contributions

B.-K.C. conceived and supervised the study. N.L. and B.-K.C. designed the experiments. N.L., W.K., S.H. and Y.L. performed the experiments. N.L., W.K., S.H., Y.L., S.C., B.P. and B.-K.C. analyzed the data. N.L., W.K., S.C., B.P. and B.-K.C. wrote the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to B.-K.C.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

The Creative Commons Public Domain Dedication waiver <http://creativecommons.org/publicdomain/zero/1.0/> applies to the metadata files associated with this article.

© The Author(s) 2020