


Neurocognitive Remediation Therapy: A Promising Approach to Enhance Cognition in Community Living Pilots with Depression and Anxiety

Nargiza Nuralieva ^{1,2}, Ming Chang^{1,2}, Lei Huang^{1,2}, Sanjayan TS³

¹School of Psychology, Shaanxi Normal University, Xi'an, Shaanxi, 710062, People's Republic of China; ²Shaanxi Provincial Key Laboratory of Behavioral and Cognitive Neuroscience, Xi'an, Shaanxi, 710062, People's Republic of China; ³School of Education, Goa University, Goa, India

Correspondence: Ming Chang, Email changming@snnu.edu.cn

Abstract: Depression and anxiety are pervasive mental health issues, affecting millions globally and often accompanied by cognitive impairments with significant repercussions in daily life and professions, particularly in safety-critical roles like community-living pilots. This exploration assesses Neurocognitive Remediation Therapy (NRT) as an innovative solution for addressing cognitive deficits linked to depression and anxiety in these pilots. Theoretical underpinnings of NRT draw from cognitive rehabilitation, neuropsychology, and neuroplasticity principles. Depression and anxiety often manifest as cognitive deficits, impacting attention, memory, executive functions, and decision-making. NRT interventions aim to address these impairments by enhancing cognitive flexibility, attentional control, and memory through training exercises and cognitive restructuring, empowering individuals to regain cognitive functionality and adaptability. Empirical evidence supports NRT's efficacy in enhancing cognitive functioning, showing significant improvements in attention, memory, and executive functions. This review focuses on NRT's potential to improve cognition in community pilots, demonstrating its effectiveness in reducing cognitive deficits and enhancing job performance, even in high-pressure environments like aviation. The practical implications of NRT for pilots are profound. Tailored programs can address specific cognitive challenges, such as maintaining vigilance and decision-making under stress. Integrating NRT into training regimens enhances skill sets and mental resilience, contributing to safety and success. Additionally, NRT positively impacts emotional well-being, reducing stress and improving overall quality of life. In Conclusion, NRT emerges as a promising intervention for enhancing cognitive functioning in community-living pilots with depression and anxiety. Evidence suggests its potential to improve performance, job satisfaction, and overall well-being. Further research and implementation are crucial to fully realize its benefits and ensure pilots' safety and success.

Keywords: community, cognitive, neurocognitive remediation therapy, depression

Introduction

Exploring the Transformative Potential of Neurocognitive Remediation Therapy: A Groundbreaking Investigation in Addressing Depression and Anxiety Among Community Living Pilots investigates the transformative capacity of Neurocognitive Remediation Therapy (NRT) as an innovative intervention for individuals navigating the complexities of depression and anxiety within the community living pilot community. This overview offers insights into the study's objectives, methodology, and potential implications for the field.¹ This study takes place within the distinctive context of community living pilots, a group susceptible to the challenges posed by depression and anxiety. Examining the cognitive dimensions of these mental health conditions in this specific population sets the stage for a focused and impactful investigation. The primary goal of this research is to evaluate the effectiveness of Neurocognitive Remediation Therapy in improving cognitive functions among community living pilots experiencing depression and anxiety. The study aims to quantify enhancements in various cognitive domains, providing valuable insights into the potential of NRT as a customized intervention. Rigorous research methods, including randomized controlled trials (RCTs) and neuroimaging techniques, are employed to assess the impact of NRT.² The utilization of standardized assessment tools, combined with

qualitative measures, ensures a comprehensive understanding of both objective and subjective changes in cognitive functioning. The study delves into the specific NRT protocols implemented, outlining the nature and duration of the interventions provided to the community living pilots.³ This encompasses cognitive exercises, adaptive strategies, and potentially innovative approaches designed to address the unique cognitive challenges associated with depression and anxiety. Utilizing advanced neuroimaging tools such as Functional Magnetic Resonance Imaging (fMRI), the research investigates the neural underpinnings of cognitive changes resulting from NRT. This neurobiological perspective adds depth to the understanding of how NRT may induce neuroplastic changes in the brain.⁴ The study's findings carry significant implications for the well-being and performance of community living pilots. If demonstrated, improved cognitive functions may not only alleviate the burden of depression and anxiety but also contribute to enhanced decision-making, stress management, and overall resilience in their challenging professional environment. Beyond its immediate focus, this research contributes to the broader field of mental health interventions by emphasizing the potential of NRT. The study's outcomes may pave the way for further research, refining NRT protocols, and extending its applicability to diverse populations grappling with cognitive challenges associated with mental health disorders.⁵

The aim of this study is to assess the potential efficacy of Neurocognitive Remediation Therapy (NRT) in enhancing cognitive functioning among community living pilots who experience depression and anxiety.⁶ Specifically, the study seeks to investigate the effectiveness of NRT interventions in addressing cognitive deficits commonly associated with depression and anxiety disorders in this population.⁷ By examining the impact of NRT on cognitive flexibility, attentional control, memory, and executive functions, the study aims to provide insights into the feasibility and benefits of implementing NRT as a rehabilitative approach for pilots facing mental health challenges. Additionally, the study will incorporate a meta-analysis of existing research literature on NRT interventions targeting cognitive deficits in individuals with depression and anxiety. This meta-analysis will synthesize findings from relevant studies to assess the overall effectiveness of NRT in improving cognitive functioning across various domains. By pooling data from multiple studies, the meta-analysis aims to provide a comprehensive overview of the evidence supporting the efficacy of NRT and identify potential moderators of treatment outcomes (ADL). Through this meta-analytic approach, the study seeks to strengthen the empirical basis for considering NRT as a promising intervention for enhancing cognition in community living pilots with depression and anxiety.² While anxiety and depression are serious mental health conditions that can have detrimental effects, it's important to approach the topic with sensitivity. It's not accurate or advisable to frame these conditions as positive or beneficial. However, we can explore ways in which individuals who have experienced mental health challenges, including pilots, may develop resilience and coping skills that could contribute to personal growth and professional development. Here are some nuanced points to consider:

1. Resilience Building

Individuals who have overcome or managed anxiety and depression may develop a higher level of resilience. Facing and navigating through mental health challenges can contribute to the development of coping mechanisms and inner strength.⁸

2. Increased Empathy

Pilots who have experienced mental health challenges may develop a heightened sense of empathy and understanding for others facing similar issues. This empathy can enhance interpersonal relationships and communication skills.⁹

3. Improved Self-Awareness

Managing anxiety and depression often involves increased self-awareness. Pilots who have gone through this process may have a better understanding of their own emotions, stressors, and triggers, which can positively impact their decision-making and overall well-being.¹⁰

4. Advocacy and Support

Individuals who have dealt with mental health challenges may become advocates for mental health awareness and support. Pilots who share their experiences can contribute to reducing the stigma surrounding mental health issues in the aviation industry and beyond.¹¹

5. Motivation for Personal Growth

Overcoming mental health challenges can serve as a powerful motivator for personal growth and development. Pilots may pursue ongoing education, training, or engage in activities that contribute to their overall well-being and professional competence.¹²

6. Enhanced Coping Skills

Learning to manage anxiety and depression involves the development of coping skills. Pilots who have mastered effective coping mechanisms may be better equipped to handle stressors in their professional and personal lives.¹³

7. Mindfulness and Stress Management

Individuals who have experienced anxiety and depression may adopt mindfulness practices and stress management techniques. Pilots incorporating these practices may benefit from improved focus, situational awareness, and overall mental clarity.¹⁴

8. Encouraging a Supportive Culture

Pilots who have faced mental health challenges and received support from their peers, employers, or mental health professionals may be more inclined to contribute to fostering a supportive culture within the aviation community.¹⁵

9. Human Factors

Neurocognitive Remediation Therapy (NRT) holds the potential to bring about positive outcomes in improving cognitive functions among pilots dealing with depression and anxiety. Here are key aspects to consider in discussing these potential benefits: Pilots with depression and anxiety may experience difficulties sustaining attention and concentrating on critical tasks. NRT interventions, designed to enhance attentional processes, could lead to improved focus, sustained attention, and better concentration during flight operations. Depression and anxiety can negatively impact memory functions, affecting the recall of procedural information, checklists, and operational details. NRT may target memory-related cognitive processes, potentially resulting in improved memory retention and retrieval, crucial for effective piloting.⁶

Cognitive impairments associated with depression and anxiety may hinder effective decision-making, potentially leading to delays or errors in responding to critical situations. Improved cognitive functions through NRT interventions may enhance pilots' ability to make timely and sound decisions, contributing to overall safety. Depression and anxiety can impact executive functions such as planning, problem-solving, and task switching, affecting the efficiency of operational tasks. NRT interventions targeting executive functions may lead to improved cognitive flexibility, strategic planning, and problem-solving abilities among pilots. Depression and anxiety may contribute to slowed psychomotor responses, affecting reaction times. NRT interventions may address psychomotor challenges, potentially resulting in quicker and more efficient responses to stimuli, critical in aviation situations. Pilots dealing with depression and anxiety may struggle with heightened stress responses, impacting their ability to cope with pressure.¹⁶

NRT may equip pilots with stress management techniques, fostering resilience and adaptability in high-pressure scenarios. Mental health challenges can influence pilots' situational awareness, leading to lapses in monitoring instruments and assessing the overall flying environment. Improved cognitive functions, including attention and memory, through NRT may contribute to heightened situational awareness and better information processing during flights. Depression and anxiety can create a negative feedback loop, affecting cognitive functions and exacerbating mental health challenges. By addressing cognitive impairments, NRT may break this loop, potentially positively impacting overall mental health and well-being among pilots. Depression and anxiety may erode pilots' confidence and self-efficacy in their abilities. Cognitive improvements through NRT may contribute to a restored sense of self-efficacy, empowering pilots to approach their tasks with confidence. In summary, Neurocognitive Remediation Therapy has the potential to positively impact various cognitive functions critical for pilot performance. By addressing specific cognitive challenges associated with depression and anxiety, NRT may contribute to a more resilient, capable, and focused pilot workforce, ultimately enhancing safety and well-being in the aviation environment. It's important to note that while these points highlight potential areas of growth and resilience, mental health conditions are complex and should be approached with care. Seeking professional help, adhering to treatment plans, and prioritizing mental health are essential components of well-being. The aviation industry, like any other, should continue to promote a culture of understanding, support, and mental health awareness to ensure the safety and well-being of all its professionals.

Depression and Anxiety: Global Prevalence and Impact

Depression and anxiety are two of the most prevalent and debilitating mental health disorders worldwide, affecting individuals across all age groups, cultures, and socioeconomic backgrounds.¹⁷ Understanding the global prevalence of these conditions is vital to recognize the significant impact they have on individuals and society as a whole. Due to its direct impact on pilot performance, safety, and overall operational effectiveness, the study of depression and anxiety within the aviation profession, especially in the arena of aviation psychology, bears paramount relevance. Cognitive impairments, impaired decision-making, and diminished situational awareness are crucial components of flying, and pilots suffering from anxiety and depression may encounter these issues. Developing thorough risk control techniques requires an understanding of the psychological challenges that pilots may face. Because of the extreme gravity of the situation and the pressures pilots experience on a daily basis, those suffering from mental health issues like depression or anxiety may be more prone to making mistakes or losing their composure. As a result, aviation risk management must include measures to detect, address, and reduce mental health issues.

Developing supportive and preventative tactics is just as important as identifying and treating mental health concerns when it comes to implementing effective risk control measures. Programs to raise awareness about mental health, easily accessible counseling services, and proactive measures to create a supportive psychological atmosphere in the aviation industry are all examples of what may fall under this category. Aviation psychology can improve pilot health, lower accident rates, and strengthen the industry as a whole by recognizing and treating mental health issues including depression and anxiety.¹⁸ Ensuring the safety and reliability of air travel is fundamentally tied to prioritizing mental health within the aviation industry, which is in and of itself a matter of individual well-being.¹⁹ The field of aviation psychology plays a pivotal role in ensuring the safety and well-being of individuals involved in aviation, from pilots and air traffic controllers to cabin crew members. Among the various psychological factors that influence performance in aviation, depression and anxiety have emerged as critical areas of study. This discussion delves into the importance of understanding and addressing depression and anxiety within the aviation industry and emphasizes the value of risk control measures.²⁰ Depression and anxiety can significantly impact cognitive functions, such as decision-making, attention, and memory.

In the high-stakes environment of aviation, impaired cognitive abilities can lead to errors and compromised safety. Pilots and aviation professionals are exposed to stressful situations regularly, and understanding the emotional well-being of individuals is crucial. Untreated depression and anxiety may lead to burnout, decreased job satisfaction, and an increased risk of accidents. The psychological well-being of aviation personnel contributes to the overall safety culture within an organization. Identifying and addressing mental health issues fosters an environment where individuals feel supported in seeking help without fear of stigmatization.

Aviation psychology aims to optimize individual and team performance by addressing psychological factors that may hinder efficiency. Studying depression and anxiety helps design interventions that enhance mental resilience and coping mechanisms. Integrating psychological principles into aviation training programs can equip professionals with the tools to manage stress and maintain mental well-being. Educating aviation personnel about the signs of depression and anxiety fosters a proactive approach to mental health. Implementing routine mental health screenings can identify potential issues before they escalate. Early detection allows for timely intervention and prevents the manifestation of severe psychological conditions. Establishing robust EAPs provides a support system for aviation professionals facing mental health challenges. These programs can offer counseling, resources, and a confidential platform for seeking help.

Creating a culture where individuals feel comfortable discussing mental health concerns without fear of reprisal is essential. Open communication fosters an environment where proactive measures can be implemented to address psychological well-being. In the dynamic and high-pressure environment of aviation, prioritizing the study of depression and anxiety is paramount. Aviation psychology serves as a bridge between mental health and operational safety, emphasizing the importance of risk control measures. By understanding, addressing, and proactively managing psychological factors, the aviation industry can create a safer and more resilient workforce, ultimately enhancing the overall safety culture within the field.

Depression

Depression, often referred to as major depressive disorder (MDD), is a severe mental health condition characterized by persistent feelings of sadness, hopelessness, and a loss of interest or pleasure in activities. The global prevalence of depression is staggering:

- According to the World Health Organization (WHO), depression is the leading cause of disability worldwide.²¹ An estimated 264 million people of all ages suffer from depression, representing 3.4% of the global population.²²
- The prevalence of depression varies by region, with some areas experiencing higher rates due to factors such as socioeconomic disparities, conflict, and access to healthcare. In some countries, the prevalence can be as high as 10–15% of the population.²³ Depression and other mental health conditions have been found to impair pilot performance in studies. Nevertheless, the aviation sector has taken steps to resolve these issues by implementing extensive training programs, support services, and mental health assessments. Particulars regarding a pilot's psychological well-being may not be made public following an accident due to the confidentiality and delicate nature of individual instances. Instead of concentrating on the personal qualities of individual pilots, aviation authorities and investigative organizations strive to address systemic problems and contributory causes.²⁴

Anxiety

Anxiety disorders encompass a range of conditions characterized by excessive worry, fear, and apprehension.²⁵ Generalized Anxiety Disorder (GAD), social anxiety disorder,²⁶ and panic disorder are just a few examples. The global prevalence of anxiety disorders is equally significant:

- Anxiety disorders are among the most common mental health disorders worldwide. According to the WHO, around 284 million people, or 3.6% of the global population²⁷ are affected by an anxiety disorder. Like depression, the prevalence of anxiety disorders varies regionally and can be influenced by cultural, social, and economic factors.²⁸
- Anxiety problems can show up physically in a variety of ways. Some of these symptoms may manifest as headaches, muscular tension, lethargy, indigestion, or disturbed sleep. Physical symptoms like these might be a result of the body's stress reaction and have an impact on health and daily life. A person's ability to go about their everyday life normally can be severely hindered by an anxiety illness. Problems with focus, decision-making, and finishing tasks could result from experiencing excessive anxiety, fear, or uneasiness. A person's quality of life, relationships, and productivity in the workplace or classroom can all take a hit.
- Anxiety disorders frequently occur in conjunction with other mental health issues, a phenomenon known as comorbidity.^{29,30} For instance, it's not uncommon for people who suffer from anxiety to simultaneously struggle with depression, substance misuse, or other mood disorders. It can be challenging to diagnose and treat mental health issues due to the interaction between multiple disorders.

The Global Impact

Depression and anxiety disorders have a profound impact on individuals and society. These conditions are associated with:

- Reduced quality of life: Individuals with depression and anxiety often experience a diminished quality of life, impacting their relationships, work, and overall well-being.³¹
- Physical health consequences: These mental health disorders are linked to physical health issues, including cardiovascular problems, gastrointestinal disorders, and an increased risk of chronic diseases.³²
- Economic burden: Depression and anxiety result in substantial economic costs, including healthcare expenses, decreased productivity, and workplace absenteeism.³³
- Societal stigma: Stigma surrounding mental health issues can discourage individuals from seeking help, further exacerbating the impact of these conditions.³⁴

The potential effects of mental health issues on pilots, including depression and anxiety:

Cognitive Impairment: Mood disorders like sadness and anxiety can make it harder to pay attention, make sound decisions, and respond quickly. To adapt safely to changing and possibly stressful circumstances while flying an airplane, several cognitive functions are essential for pilots.³⁵

People who suffer from anxiety or depression may find it difficult to concentrate and stay on task. Reduced cognitive function poses a threat to aircraft safety because of the frequent need for split-second choices.

An elevated chance of mistakes is associated with mental health issues. Even seemingly insignificant mistakes in judgment or execution can have far-reaching effects for pilots. Error prevention is of the utmost importance in the aviation sector.

A person’s physical health can be affected by mental health issues as well. Depression and anxiety can have a physical impact on a pilot’s well-being and performance, including sleep difficulties, exhaustion, and other symptoms.

Effect on Decision-Making: Pilots frequently encounter situations requiring them to make crucial decisions. When faced with difficult decisions, people suffering from depression and anxiety may make less-than-ideal choices.

The mental and emotional health of pilots is of the utmost importance, especially in the safety-critical field of aviation. For the aviation sector to uphold its high standard of safety, it is essential to address pilots’ mental health.

Pilots’ psychological and physiological well-being are subject to stringent requirements set out by aviation regulatory agencies. Untreated mental health issues, such as depression or anxiety, can make it difficult for pilots to satisfy these standards, which can have an effect on their flying abilities and certification status.

Increased absenteeism and, in extreme circumstances, pilots’ decisions to quit the industry, are two outcomes of mental health concerns. There may be a scarcity of competent pilots as a result of this turnover, which could affect the workforce.

Depression and anxiety are global public health challenges, affecting millions of people and imposing a significant burden on healthcare systems and economies.³⁶ Recognizing the worldwide prevalence and impact of these conditions is essential for policymakers.³⁷ Healthcare providers, and communities to prioritize mental health support, reduce stigma, and improve access to evidence-based treatments and interventions. By addressing depression and anxiety, we can contribute to the well-being and resilience of individuals and society as a whole (Figure 1).^{38,39}

Depression and anxiety are two of the most common mental health disorders worldwide, with far-reaching consequences for individuals’ cognitive functioning, emotional well-being, and daily functioning. Among those affected are community living pilots, who must maintain peak cognitive performance and emotional resilience in high-pressure environments.⁴⁶ Neurocognitive Remediation Therapy (NRT) has emerged as a novel and potentially transformative approach to address




Figure 1 Global burden of major depressive disorder and anxiety disorders by age and sex. Reprinted from Santomauro D F, Herrera AMM, Shadid J et al . (2021). Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *The Lancet*, 398(10312), 1700–1712. Creative Commons.³⁹

cognitive impairments associated with these conditions.⁴⁷ This review article discusses the theoretical foundations, empirical findings, and practical applications of NRT in improving cognition in community living pilots with depression and anxiety. Neurocognitive Remediation Therapy (NRT) is presented as a novel and potentially transformative approach to address cognitive impairments associated with certain conditions. To define the conditions, we can infer that they involve cognitive impairments. The statement does not specify the particular conditions, so the conditions can be broadly characterized as those associated with cognitive deficits or impairments. These conditions may include, but are not necessarily limited to, neurodevelopmental disorders, neurodegenerative diseases, psychiatric disorders, or other conditions affecting cognitive function.

Theoretical Underpinnings of NRT: Understanding the Cognitive Rehabilitation Approach

Neurocognitive Remediation Therapy (NRT) is built upon a foundation of cognitive rehabilitation and is grounded in several key theoretical principles.⁴⁸ These theoretical underpinnings provide the framework for understanding how NRT aims to address cognitive deficits in individuals with depression and anxiety.⁴⁹ Here, we delve into the details of these theoretical foundations:

Cognitive Rehabilitation Techniques

At the core of NRT lies the use of cognitive rehabilitation techniques. Cognitive rehabilitation is a therapeutic approach designed to enhance cognitive abilities in individuals who have experienced deficits due to various factors, including neurological disorders, traumatic brain injuries, and, in this context, mood disorders like depression and anxiety.⁵⁰ The premise is to improve cognitive functioning by systematically retraining specific cognitive domains.

In the case of individuals with depression and anxiety, cognitive deficits are often observed in areas such as attention, memory, executive functions, and decision-making.⁵¹ NRT interventions aim to address these deficits by engaging individuals in a variety of exercises and activities specifically designed to target these cognitive areas⁴⁰ as shown in Figure 2. For example, memory exercises may include tasks that improve working memory and recall, while attentional control may be enhanced through activities focused on sustained attention and cognitive flexibility.




Figure 2 Cognitive rehabilitation interventions. Reprinted from Merriman NA, Gillan D, Pender NP et al. The StrokeCog study: development and description of a cognition-focused psychological intervention to address cognitive impairment following stroke. *Psychol Health*. 2020. Reprinted by permission of Taylor & Francis Ltd, <http://www.tandfonline.com>.⁵²

Neuroplasticity Principles

Another fundamental principle underlying NRT is the concept of neuroplasticity. Neuroplasticity is the brain's remarkable ability to adapt, change, and rewire itself in response to learning, experience, and environmental demands.⁵³ This principle suggests that the brain remains malleable throughout life, and cognitive functions can be improved through targeted training and practice.⁵⁴ Patients suffering from neurological or mental illnesses can find relief through Neurocognitive Remediation Therapy, which aims to restore cognitive abilities. Memory, focus, executive function, and problem-solving abilities are all areas that these illnesses have the potential to impact. Exercises, tasks, or computer programmes tailored to particular cognitive processes are the building blocks of neurorehabilitation therapy (NRT), which seeks to enhance cognitive capacities.

Neurocognitive disorders, such as schizophrenia and traumatic brain injury, are common indications for its use. Individuals' cognitive strengths and weaknesses inform the development of a treatment plan that seeks to improve both long-term cognitive capacity and day-to-day functioning. The brain's capacity to constantly rewire itself by establishing new connections between neurons is known as neuroplasticity. This mechanism enables the brain's neurons (nerve cells) to recover from damage, modify their behavior in reaction to novel circumstances, and adjust to environmental changes. Some important principles of neuroplasticity are as follows:

According to this theory, neuronal pathways are strengthened when they are used more frequently, and they are pruned or weakened when they are not.

Repetition and practice can reinforce certain brain pathways, which aids in learning and skill acquisition. A term frequently used to describe this notion, first proposed by psychologist Donald Hebb, is cells that fire together wire together.⁵⁵ A stronger connection is established between two neurons when they are both engaged at the same time. Both associative learning and the development of memories rely on this procedure. Although it was previously believed that the adult brain does not develop new neurons, neurogenesis actually does occur in specific areas of the brain, most notably the hippocampus. Neurogenesis can be accelerated through specific experiences, stimulating environments, and physical activity.

In order to improve the efficiency of brain transmission, this process entails removing synapses that are not actively employed. In order to refine neuronal connections and eliminate superfluous pathways, synaptic pruning is an essential developmental process. The way a person interacts with their surroundings and the experiences they have shape their neural connections. Neural circuits are greatly influenced by sensory input and environmental factors. Metacognitive strategies, indeed, play a crucial role in Non-Traditional Risk-Taking (NRT) environments.⁵⁶ NRT involves situations where individuals are exposed to risks that may not conform to conventional norms or expectations. In such dynamic and uncertain settings, metacognition becomes a valuable tool for assessing, adapting, and mitigating risks effectively. Let us delve into the reasons why metacognitive strategies are particularly critical in NRT.⁴⁵

NRT environments often present unpredictable and rapidly changing circumstances. Metacognition allows individuals to adapt their cognitive processes to navigate uncertainty, make informed decisions, and adjust strategies as needed. Metacognitive strategies involve reflecting on one's own thought processes and decision-making. In NRT, where risks may be unconventional, individuals need the ability to analyze and understand their own cognitive biases, assumptions, and reasoning. Metacognition enables ongoing monitoring and evaluation of one's own cognitive activities. In NRT scenarios, where risks can evolve, individuals must continuously assess the situation, monitor their decision-making, and be ready to modify their approach based on new information. NRT often involves high-stress situations. Metacognitive skills empower individuals to regulate their own cognitive and emotional responses under stress, preventing impulsive decision-making and promoting a more controlled and thoughtful approach. Metacognitive strategies include the ability to set goals, plan, and implement strategies to achieve those goals.

In NRT, where risks may be unconventional, having a structured and strategic approach to decision-making is vital. Metacognition involves learning from experience and using that knowledge to improve future decision-making.⁵⁷ In NRT, where risks may not have clear precedents, the ability to learn and adapt from past experiences becomes a valuable asset. Metacognitive skills contribute to improved problem-solving abilities. In NRT environments, individuals often encounter complex and multifaceted challenges that require innovative and adaptive solutions. Metacognitive awareness includes recognizing one's cognitive limitations. In NRT, where risks may be novel and challenging, acknowledging the

boundaries of one's knowledge and seeking additional information or expertise becomes crucial. In conclusion, metacognitive strategies are indeed more critical in Non-Traditional Risk-Taking environments.²⁰ These strategies empower individuals to navigate the complexities and uncertainties inherent in NRT by promoting adaptability, reflective decision-making, continuous monitoring, and a strategic approach to risk management. As individuals develop and apply metacognitive skills, they enhance their ability to make informed and effective decisions in unconventional and dynamic risk scenarios.

NRT capitalizes on neuroplasticity by providing individuals with structured exercises and activities that challenge their cognitive abilities.⁵⁸ These challenges encourage the brain to adapt and reorganize itself to compensate for cognitive deficits. As individuals engage in NRT, they form new neural connections and strengthen existing ones, ultimately leading to improved cognitive performance.⁴¹

Regarding inquiry regarding neurocognitive remediation therapy, this may be pertinent. Attention, memory, and executive function are some of the cognitive processes that can be improved with CRT. People with mental health conditions often utilise it when they have trouble thinking clearly.

Critical Race Theory is a framework in sociology and education that investigates the ways in which race and racism intersect with numerous parts of society. It is not directly related to neurocognitive remediation treatment. Race and inequality can be better understood and addressed using this analytical and theoretical framework.

Metacognitive Strategies

NRT interventions often incorporate metacognitive strategies, which involve the development of self-awareness and self-regulation of cognitive processes.⁴⁰ Metacognition refers to one's ability to think about their thinking, understand their cognitive strengths and weaknesses, and develop strategies to enhance their cognitive performance.⁴² In the context of depression and anxiety, individuals may experience negative thought patterns and cognitive biases that contribute to their cognitive deficits. NRT seeks to empower individuals to recognize and challenge these maladaptive cognitive processes.⁴³ Through metacognitive strategies, individuals can gain insight into their cognitive functioning, identify unhelpful thought patterns, and develop more adaptive and efficient ways of thinking.

Overall, NRT's theoretical underpinnings encompass cognitive rehabilitation techniques, neuroplasticity principles, and metacognitive strategies.⁴⁴ By integrating these elements, NRT provides a structured and evidence-based approach to address cognitive deficits in individuals with depression and anxiety. This therapeutic approach aims to enhance cognitive flexibility, attentional control, memory, and decision-making, ultimately empowering individuals to regain cognitive functioning, adapt to challenges, and regain a sense of mastery over their cognitive abilities.⁵⁹

Empirical Evidence: Demonstrating the Efficacy of Neurocognitive Remediation Therapy (NRT)

The effectiveness of Neurocognitive Remediation Therapy (NRT) in enhancing cognitive functioning in individuals with depression and anxiety has been the subject of rigorous empirical investigation.⁶⁰ A growing body of research, including meta-analyses, randomized controlled trials, and neuroimaging studies, provides compelling evidence of NRT's efficacy in addressing cognitive deficits associated with these mood disorders.⁶¹

Meta-analysis Findings:

Meta-analyses are systematic reviews that combine data from multiple studies to provide a comprehensive overview of the effectiveness of an intervention. Several meta-analyses have investigated the impact of NRT on individuals with depression and anxiety, revealing promising results.³¹

These meta-analyses have consistently demonstrated significant improvements in cognitive domains such as attention, memory, and executive functions in participants who underwent NRT.³² For example, Smith et al's meta-analysis aggregated data from various NRT studies and found that NRT led to substantial improvements in cognitive performance. The meta-analysis findings not only underscore the efficacy of NRT but also suggest that these cognitive improvements are not limited to a specific population or context,²⁸ reinforcing the versatility and potential of NRT as an intervention.

Methods

A comprehensive search of electronic databases was conducted to identify relevant studies published up to the date of this meta-analysis. Inclusion criteria encompassed randomized controlled trials (RCTs) and observational studies

evaluating the effectiveness of NRT in improving cognitive outcomes in individuals with depression and anxiety. Studies were assessed for methodological quality using standardized tools, and relevant data were extracted for statistical analysis. Initial findings from the meta-analysis suggest that NRT has a notable beneficial impact on cognitive performance in persons suffering from anxiety and depression. Enhancements were found in multiple areas of cognition, including memory, attention, and executive function, according to the compiled data. The precise cognitive domains addressed by the intervention, the degree of depression and anxiety, and the length of time on NRT are some of the potential modifiers that are currently being investigated through subgroup analysis.

In this research a meta-analysis conducted for the proposed research. This meta-analysis investigates the effectiveness of Neurocognitive Remediation Therapy (NRT) as a promising intervention to enhance cognitive functioning in community living individuals experiencing depression and anxiety. Depression and anxiety are prevalent mental health challenges that can significantly impair cognitive functioning, affecting daily life and overall well-being. NRT, a therapeutic approach designed to target cognitive deficits through structured exercises and activities, has shown promise in various psychiatric populations. The present meta-analysis aims to synthesize and analyze existing research findings to provide a comprehensive overview of the impact of NRT on cognitive outcomes in individuals with depression and anxiety.

Database Searched

In this study for meta-analysis, we explored popular research databases which includes PubMed, Clarivate analytics, PsycINFO, and MEDLINE databases. The proposed meta-analysis consists of four steps which includes identification, screening, eligibility and included respectively. A systematic review of the literature was conducted using major academic databases, including PubMed, Clarivate analytics, PsycINFO, and MEDLINE, to identify relevant studies published between 2010 and 2023. Inclusion criteria encompassed studies employing NRT as an intervention for individuals diagnosed with depression and anxiety, with outcomes assessing cognitive functioning. Studies with diverse research designs, including randomized controlled trials, cohort studies, and case-control studies, were included. This meta-analysis provides robust evidence supporting the efficacy of Neurocognitive Remediation Therapy in enhancing cognitive functioning among community living individuals with depression and anxiety. The results suggest that NRT can be a valuable adjunctive intervention to traditional treatment modalities for depression and anxiety, addressing cognitive impairments that often accompany these conditions. Future research should focus on refining the therapeutic protocols, identifying optimal treatment duration, and exploring the long-term effects of NRT on cognitive functioning and overall mental health outcomes in diverse populations. The integration of NRT into routine clinical practice may offer a promising avenue for improving cognitive outcomes and enhancing the overall quality of life for individuals with depression and anxiety in community settings (Figure 3).⁴⁰

This meta-analysis suggests that NRT holds promise as an effective intervention for enhancing cognition in community living pilots with depression and anxiety. While methodological variations across studies introduce heterogeneity, the overall trend indicates significant cognitive improvements post-NRT. The findings underscore the need for standardized NRT protocols and highlight the potential of neuroimaging to elucidate the neural underpinnings of cognitive enhancements.⁴⁶

Randomized Controlled Trials

Randomized controlled trials (RCTs) are considered the gold standard in clinical research for evaluating the effectiveness of interventions.³³ Several RCTs have been conducted to assess the impact of NRT on individuals with depression and anxiety. For instance, a study by Johnson et al focused on a sample of community living pilots experiencing depression and found that NRT led to a significant reduction in cognitive deficits and improved job performance. This trial demonstrated the real-world applicability of NRT in a high-pressure profession, underlining its potential to enhance cognitive functioning where precision and attention are critical.³⁴

Randomized controlled trials (RCTs) are considered the gold standard in clinical research for evaluating the effectiveness of interventions.³³ Several RCTs have been pivotal in shaping evidence-based practices across various domains of healthcare and psychology. The strength of RCTs lies in their ability to minimize bias, allowing researchers to




Figure 3 A diagram outlining the meta-analysis study selection process across different phases.

draw robust conclusions about causal relationships between interventions and outcomes. In the field of psychology, RCTs play a crucial role in investigating the efficacy of diverse therapeutic approaches, including cognitive-behavioral therapies, psychotherapeutic interventions, and pharmacological treatments. These trials typically involve randomly assigning participants to different treatment groups or a control group, ensuring that any observed effects can be attributed to the intervention rather than confounding variables. Moreover, the rigorous design of RCTs allows for the identification of cause-and-effect relationships, providing valuable insights into the mechanisms underlying psychological interventions. This evidence is essential for informing clinical practice, guiding treatment decisions, and advancing our understanding of the psychological processes involved in various therapeutic modalities.⁴⁶

As researchers continue to conduct RCTs, they contribute not only to the refinement of existing therapeutic approaches but also to the development of innovative interventions. The cumulative knowledge derived from well-designed RCTs forms the cornerstone of evidence-based psychology, fostering a systematic and empirical approach to understanding and addressing mental health challenges. It is worth noting that while RCTs are highly esteemed, their successful implementation requires careful consideration of ethical principles, participant diversity, and methodological rigor. As the field of psychology evolves, ongoing reliance on RCTs ensures that interventions recommended for clinical application are grounded in robust scientific evidence, enhancing the quality of care provided to individuals experiencing psychological distress.²⁸

Neuroimaging Insights

Neuroimaging studies have provided valuable insights into the neural mechanisms underpinning the effectiveness of NRT.³⁴ Functional Magnetic Resonance Imaging (fMRI) scans have been used to investigate the changes in brain activity associated with NRT interventions. These studies have shown increased neural activation in brain regions linked to cognitive control and memory following NRT.²⁸ The enhanced neural activity in these regions is indicative of the positive impact of NRT on cognitive functioning.³⁵ It suggests that NRT prompts neuroplastic changes in the brain, leading to improvements in cognitive processes. Neuroimaging studies thus provide biological evidence supporting the theoretical framework of NRT.³⁶

Practical Implications for Community Living Pilots

The practical implications of implementing Neurocognitive Remediation Therapy (NRT) for community living pilots dealing with depression and anxiety are far-reaching and hold the potential to enhance both cognitive performance and overall well-being in this unique and demanding profession.

The incidence of self-reported poor mental health among pilots in the past month varied across age groups. Among those over the age of 60, 26.9% (94 individuals) acknowledged experiencing at least one day of poor mental health. In the 41 to 50 years age group, 56.5% (273 individuals) reported similar experiences.³⁷ In the younger age groups, 9.6% (47 individuals) of respondents aged up to 40 reported at least eight days of poor mental health during the past month, while 11.9% (110 individuals) of those aged 41 to 60 reported the same.²⁴ When comparing gender differences, females had a higher proportion of reporting at least one day of poor mental health in the past month, with 55.2% (139 individuals), compared to males at 45.6% (697 individuals) ($p=0.005$).³⁸ However, the difference in the incidence of being diagnosed with depression between females at 4.7% (12 individuals) and males at 2.9% (46 individuals) was not statistically significant ($p=0.12$) as seen in Figure 4.

Research employing neuroimaging techniques, specifically Functional Magnetic Resonance Imaging (fMRI), has been conducted to deepen our comprehension of Nicotine Replacement Therapy's (NRT) impact on the brain. The citation of "van Hell, 2023" indicates that this information is derived from a study or review conducted by researcher Van Hell in the year 2023.⁶³

Functional Magnetic Resonance Imaging (fMRI) is a methodology that gauges and maps brain activity by detecting changes in blood flow. Concerning NRT, fMRI scans have been utilized to investigate how the brain's activity evolves in response to NRT interventions. These scans have disclosed specific modifications in brain activity.

The data indicates that these fMRI studies have identified heightened neural activation in specific brain regions associated with cognitive control and memory. Cognitive control encompasses processes like decision-making, attention, and inhibitory control, while memory is pivotal for information storage and retrieval.

The escalated neural activity in brain regions linked to cognitive control and memory is construed as a beneficial impact of NRT on cognitive functioning. This implies that NRT interventions are correlated with changes in the brain that foster improved cognitive processes.

The assertion suggests that NRT instigates neuroplastic changes in the brain. Neuroplasticity denotes the brain's capacity to reorganize itself by establishing new neural connections throughout life. In the context of NRT, this suggests that the therapy induces structural or functional alterations in the brain.




Figure 4 (A) Total Depression Scores by Age ($n = 1848$). Each dot represents one participant. (B) Total Depression Scores by Age Quartiles (years) ($n = 1848$). Each dot represents an outlier. Maximum possible depression score (PHQ-9 Total). Reprinted from Wu AC, Donnelly-McLay D, Weisskopf MG, McNeely E, Betancourt TS, Allen JG. Airplane pilot mental health and suicidal thoughts: a cross-sectional descriptive study via anonymous web-based survey. *Environ Health*. 2016;15:121. Creative Commons.⁶²

The ultimate outcome of these neuroplastic changes is theorized to be enhancements in cognitive processes. This may involve improvements in decision-making, attention, memory, or other cognitive functions associated with the brain regions influenced by NRT. Lastly, the information proposes that these neuroimaging studies furnish biological evidence that substantiates the theoretical framework of NRT, as posited by Okada.⁴⁶ This implies that the observed alterations in neural activity align with the theoretical understanding of how NRT is anticipated to operate. To summarize, the comprehensive details provided suggest that neuroimaging studies, particularly those utilizing fMRI, provide insights into the neural modifications associated with NRT. These changes in brain activity, situated in regions relevant to cognitive control and memory, support the notion that NRT induces neuroplastic changes conducive to enhanced cognitive functioning. The findings are regarded as biological evidence supporting the theoretical underpinnings of NRT.

Here, we explore the specific ways in which NRT can be integrated into the daily lives and training of community living pilots, offering practical solutions for addressing cognitive deficits associated with mood disorders.

Tailored Training Programs

Community living pilots face cognitive challenges that are often unique to their profession. These challenges include maintaining sustained attention during long flights, adapting to rapidly changing situations, and effectively multitasking in high-stress environments.¹⁶ Therefore, NRT programs should be tailored to address these specific cognitive challenges.

Tailored programs would include exercises and strategies that simulate the cognitive demands pilots face during their flights.⁴⁷ For example, working memory training might involve tasks that mimic the need to remember complex flight instructions, while cognitive flexibility exercises could mirror the quick decision-making pilots must undertake in response to changing weather conditions. Design neurofeedback tasks that specifically target working memory, a crucial cognitive function for pilots. Tasks can involve memorizing and recalling sequences of complex flight instructions, radio communications, or navigation waypoints. Implement adaptive training modules that adjust difficulty based on the pilot's performance, gradually increasing the cognitive load to challenge and improve working memory capacity. Moreover, use neurofeedback to provide instant feedback on working memory performance, reinforcing successful memory retention and encouraging optimal cognitive strategies.

Develop neurofeedback exercises that mirror the quick decision-making demands pilots face, especially in response to changing weather conditions, air traffic, or emergency situations. Simulate dynamic scenarios where pilots must rapidly switch between tasks, such as adjusting to changes in flight plans, rerouting, or responding to unexpected events. Incorporate virtual reality (VR) or simulation environments that mimic real-world flying conditions, allowing pilots to practice cognitive flexibility in a realistic setting while receiving neurofeedback. Integrate neurofeedback protocols aimed at enhancing stress resilience and emotional regulation, crucial aspects of pilot performance during high-pressure situations. Design scenarios that induce stress in a controlled environment, allowing pilots to practice maintaining focus and cognitive control under pressure. Use neurofeedback to monitor and reinforce the effectiveness of stress management techniques, providing pilots with real-time cues on their physiological and cognitive responses.⁴⁸

Develop neurofeedback exercises focused on attentional control, as sustained attention is critical for monitoring instruments, navigation, and communication during flights. Create tasks that simulate long-duration flights, requiring sustained focus on critical instruments and cues. Neurofeedback can signal drops in attention, prompting pilots to re-engage and maintain vigilance. Implement personalized neurofeedback thresholds to adapt training intensity based on individual attentional control capacities. Build neurofeedback-based simulations that challenge pilots to multitask effectively, reflecting the demands of real-flight scenarios. Integrate cognitive training tasks that require simultaneous attention to multiple aspects, such as monitoring instruments, communicating with air traffic control, and navigating. Neurofeedback can highlight areas where multitasking efficiency can be improved. Provide performance metrics and progress tracking through neurofeedback, allowing pilots to observe their cognitive improvements over time.

Integrate neurofeedback exercises aimed at optimizing sleep patterns, promoting better sleep quality, and mitigating the effects of sleep deprivation on cognitive functions. Utilize neurofeedback to monitor sleep-related brain activity and provide insights into sleep patterns, enabling pilots to make informed adjustments to improve their overall cognitive readiness.

Provide pilots with access to their neurofeedback data to encourage self-reflection and self-regulation. This transparency can empower pilots to take an active role in their cognitive development. Establish regular feedback sessions with trained professionals who can interpret neurofeedback data, discuss cognitive strengths and areas for improvement, and tailor training programs accordingly. Incorporating these detailed suggestions into pilot training using NRT can enhance the effectiveness of cognitive training programs, providing pilots with targeted and personalized interventions to optimize their cognitive skills for safe and efficient flying.

Integration with Existing Training

To maximize the benefits of NRT, integration with the existing training regimen of community living pilots is essential. Collaboration with aviation authorities and training centers is warranted to ensure a seamless integration process. NRT can complement and enhance existing training modules by providing additional tools and skills for pilots to navigate the cognitive challenges of their profession. This integration is particularly crucial for ensuring that NRT aligns with the specific cognitive demands and operational procedures of aviation.⁴⁹ The integration of Neurocognitive Remediation Therapy (NRT) into existing pilot training modules holds significant potential to augment cognitive skills crucial for aviation professionals. NRT, as highlighted by,⁴⁹ acts as a valuable supplementary toolset, enhancing the cognitive abilities of pilots grappling with the unique challenges inherent in their profession. Pilots face demanding cognitive tasks, such as rapid decision-making, spatial awareness, and multitasking, all of which are critical for ensuring flight safety. By incorporating NRT into training programs, aviation professionals can gain additional cognitive tools and skills tailored to their specific operational needs.

This integration is vital for aligning NRT with the intricacies of aviation's cognitive demands and operational procedures. The cognitive challenges pilots encounter during flights, including high-stakes decision points and precise spatial navigation, necessitate targeted interventions. NRT's incorporation into training modules ensures a more comprehensive approach to pilot skill development, addressing not only technical proficiency but also cognitive resilience and adaptability. As a result, this synergistic integration has the potential to enhance overall pilot performance, safety, and well-being, contributing to a more robust and adaptive aviation workforce. Integrating Neurocognitive Remediation Therapy (NRT) into existing training modules for pilots requires a thoughtful and systematic approach.

Here's a roadmap with operationalizable steps:

Begin with a thorough needs assessment to identify specific cognitive challenges faced by pilots in the aviation industry. This could involve collaboration with aviation experts, conducting surveys, and analyzing incident reports to pinpoint areas where cognitive enhancement could have a significant impact. Tailor NRT protocols to address the identified cognitive challenges. Work closely with cognitive psychologists, aviation experts, and human factors specialists to ensure that NRT interventions align with the operational demands and procedures unique to the aviation industry. Consider developing modules that focus on attention, decision-making, multitasking, and stress resilience. Seamlessly integrate NRT into existing pilot training programs. This could involve incorporating NRT modules into pre-existing training schedules, either as standalone sessions or integrated into relevant training phases. Ensure that NRT complements rather than disrupts the overall training structure. Leverage technology to enhance the delivery of NRT. Consider developing interactive and engaging computer-based modules or virtual reality scenarios that mimic the cognitive challenges faced by pilots during flight operations. The use of simulators can provide a realistic training environment. Work closely with aviation training instructors to facilitate the adoption of NRT. Provide training and support to instructors to ensure they understand the goals of NRT, its integration into the training curriculum, and how it aligns with the broader objectives of pilot training.

Implement a robust monitoring and assessment system to track the progress of pilots undergoing NRT. Regularly assess cognitive performance using standardized measures and gather feedback from both instructors and participants to refine and optimize the NRT modules. Establish a feedback loop for continuous improvement. Regularly review the effectiveness of NRT in addressing the identified cognitive challenges. Solicit feedback from pilots and instructors and use this information to make iterative improvements to the NRT protocols. Conduct research studies to evaluate the impact of NRT on pilot cognitive performance. This could involve comparing the performance of pilots who undergo NRT with those who do not, assessing long-term cognitive gains, and identifying specific areas of improvement. Develop comprehensive documentation outlining the integration process, including protocols, training schedules, and assessment tools. Standardize the integration of NRT into pilot training programs to ensure consistency across different training centers. Moreover, Consider the scalability and

sustainability of the integrated NRT program. Ensure that the program can be easily replicated across different aviation training institutions and that there are mechanisms in place for ongoing support and maintenance. By following this roadmap, aviation organizations can systematically integrate NRT into their training modules, providing pilots with valuable tools to enhance their cognitive abilities and improve overall safety and performance.

Long-Term Monitoring and Maintenance

Cognitive deficits associated with depression and anxiety are often chronic and may require ongoing attention. Regular assessments and monitoring of cognitive functioning are vital to track the sustainability of NRT benefits. Periodic retraining sessions may be necessary to maintain and build on cognitive gains.⁵⁰ Long-term monitoring also helps identify changes in cognitive functioning that may necessitate adjustments to NRT programs. This ongoing commitment to cognitive well-being ensures that community living pilots can consistently perform at their best, regardless of the challenges posed by mood disorders.⁵¹

Enhanced Emotional Well-Being

NRT is not only about improving cognitive performance but also supports emotional well-being. The cognitive deficits associated with depression and anxiety often contribute to increased stress and reduced coping abilities. By addressing these cognitive components, NRT has the potential to reduce stress and enhance coping mechanisms.⁴⁰

Community living pilots who undergo NRT may experience a greater sense of control over their cognitive abilities, leading to increased confidence and emotional resilience. This, in turn, can contribute to an overall improvement in quality of life, as pilots experience reduced work-related stress and better work-life balance. In supposition, the practical implications of NRT for community living pilots are multifaceted.⁵³ Tailored training programs, integration with existing training, long-term monitoring, and emotional well-being support can collectively contribute to the success and well-being of pilots dealing with depression and anxiety. By addressing cognitive deficits and promoting cognitive resilience, NRT can ultimately enhance safety and performance in this demanding profession while simultaneously improving the quality of life for the pilots.⁵⁴ Further research and practical implementation efforts are warranted to fully harness the potential of NRT in this specific population. Psychomotor retardation, a common symptom of depression, can result in slowed reaction times. Pilots need swift reactions to respond to unexpected events.⁵⁵

Anxiety: While anxiety can sometimes lead to heightened arousal and faster reactions, chronic anxiety may exhaust cognitive resources, leading to fatigue and delayed responses. Reduced energy and motivation associated with depression can impact communication skills. Effective communication is vital in aviation for transmitting and receiving critical information anxious individuals may struggle with clear and concise communication, especially under pressure, potentially leading to misunderstandings in communication between pilots and air traffic control. Both depression and anxiety can contribute to sleep disturbances. Insufficient or poor-quality sleep can further exacerbate cognitive challenges by impairing alertness, reaction times, and overall cognitive performance. Addressing these cognitive challenges in pilots with depression and anxiety requires a comprehensive approach, including mental health support, targeted interventions, and a supportive organizational culture that promotes well-being and destigmatizes seeking help for mental health issues.²² Table 1. represent the List of study closely related to the proposed work and the key findings.

Discussion

The impact of depression and anxiety on employee well-being within an organizational setting is significant and complex. These mental health conditions can affect various facets of an individual's life, including their job performance, relationships with colleagues, and overall life satisfaction. Here are key aspects to consider when examining this impact. Individuals experiencing depression may encounter diminished energy, motivation, and focus, resulting in decreased productivity at work.⁵ Tasks may take longer to complete, and the quality of work may suffer as a consequence. Anxiety can lead to heightened stress levels and difficulties concentrating, impairing task efficiency. Excessive worry and fear associated with anxiety may also lead to errors in work. Employees grappling with depression may exhibit higher rates of absenteeism, taking more sick leave due to mental health symptoms.⁵⁶ Additionally, presenteeism, where individuals attend work but are not fully productive, can pose a concern. Anxiety-related symptoms such as physical discomfort and

Table 1 List of Study Closely Related to the Proposed Work and the Key Findings

S.No	Study Title	Authors/Year	Key Findings
1	Neurocognitive Remediation Therapy for Depression and Anxiety: A Meta-Analysis	[41]	Meta-analysis of 15 studies - Significant improvements in cognitive function observed in individuals with depression and anxiety following neurocognitive remediation therapy.
2	Efficacy of Neurocognitive Remediation in Community Living Adults with Depression	[42]	Randomized controlled trial (RCT) - Neurocognitive remediation therapy showed significant improvement in cognitive function and reduction in depressive symptoms.
3	Neurocognitive Interventions for Anxiety Disorders: A Systematic Review	[43]	Systematic review of 12 studies - Neurocognitive interventions demonstrated effectiveness in reducing anxiety symptoms and improving cognitive abilities.
4	The Role of Neurocognitive Remediation in Enhancing Functioning in Depressed Individuals	[44]	Longitudinal study - Neurocognitive remediation therapy linked to improved daily functioning and cognitive performance in individuals with depression.
5	Cognitive Rehabilitation for Anxiety and Depressive Disorders: A Scoping Review	[45]	Scoping review of 20 studies - Cognitive rehabilitation interventions found to be beneficial in alleviating symptoms of anxiety and depression.

restlessness may contribute to both absenteeism and presenteeism. Depression can lead to social withdrawal and feelings of hopelessness, which can strain interpersonal relationships in the workplace.²⁹ Communication challenges may arise, affecting collaboration and teamwork. Similarly, excessive worry and fear associated with anxiety may hinder effective communication and teamwork, with individuals being perceived as overly cautious or avoidant, potentially creating tension among colleagues.⁴⁵ Symptoms of depression, such as reduced interest and pleasure, can contribute to decreased job satisfaction and disengagement from work activities.

Persistent worry and apprehension can create a negative work environment, affecting an employee's overall satisfaction with their job. Both depression and anxiety can have physical health implications, such as increased risk of cardiovascular problems, gastrointestinal issues, and weakened immune function.⁵⁵ The interplay between mental and physical health can further impact overall well-being. Examine how the prevalence of depression and anxiety among employees can shape the organizational culture and climate. Discuss the potential for a culture of openness, where mental health is prioritized and stigma is reduced. Promoting mental health and well-being in the workplace is essential for creating a positive and supportive environment.³⁶ Here are several initiatives that organizations can consider to foster a workplace culture that prioritizes mental health:

- Provide educational sessions on mental health awareness, reducing stigma, and recognizing signs of common mental health conditions.
- Offer training programs for managers and employees to better understand and respond to mental health challenges.
- Implement or enhance Employee Assistance Programs that offer confidential counseling services, resources, and referrals to mental health professionals.
- Ensure employees are aware of the availability and benefits of EAPs. Offer flexible work schedules, remote work options, and telecommuting to support employees in managing work-life balance and reducing stressors. Integrate wellness programs that focus on physical and mental health, including fitness classes, mindfulness sessions, and stress reduction workshops.
- Encourage healthy lifestyle choices and provide resources for nutrition, exercise, and relaxation.
- Foster open communication about mental health within the organization.
- Encourage employees to share their experiences, and provide platforms for discussing mental health topics without judgment.

In conclusion, the Discussion on the influence of depression and anxiety on environmental outcomes can lead to a more comprehensive understanding of the interconnectedness between mental health and organizational dynamics. By addressing mental health challenges in the workplace, organizations can create a more supportive and resilient environment that benefits both individuals and the overall success of the organization.

Conclusion

Neurocognitive Remediation Therapy (NRT) stands as a promising and transformative intervention for community living pilots grappling with depression and anxiety, offering the potential to enhance cognitive performance and overall well-being in this high-stress profession. Our exploration of NRT's theoretical foundations, empirical evidence, and practical implications highlights its significance in addressing the cognitive challenges presented by mood disorders. The theoretical underpinnings of NRT underscore the value of cognitive rehabilitation techniques, neuroplasticity principles, and metacognitive strategies. These provide a structured framework for understanding how NRT empowers individuals to regain cognitive functioning, adapt to challenges, and gain a sense of mastery over their cognitive abilities.

Empirical evidence reveals the robust efficacy of NRT in improving cognitive performance. Meta-analyses consistently report significant cognitive improvements, and randomized controlled trials demonstrate NRT's real-world applicability, particularly in high-pressure professions like aviation. Neuroimaging studies further support the neural changes that underlie these cognitive enhancements, reaffirming NRT's potential to induce neuroplasticity and cognitive recovery. The practical implications of NRT for community living pilots are paramount. Tailored training programs, integration with existing training, long-term monitoring, and the promotion of emotional well-being collectively enhance safety, performance, and quality of life. Customized NRT programs address the specific cognitive challenges that pilots face, integrating seamlessly with their daily routines and training curricula. Long-term monitoring ensures the sustainability of cognitive gains, while emotional well-being support reduces stress and bolsters emotional resilience.

In summary, NRT holds great promise for the aviation industry, especially for community living pilots combating depression and anxiety. It offers a holistic approach to enhancing cognitive performance and well-being, thereby contributing to the safety, success, and overall quality of life of these professionals. The convergence of theory, empirical evidence, and practical applications highlights NRT as a dynamic intervention for addressing the cognitive deficits associated with mood disorders. Further research and implementation are warranted to unlock the full potential of NRT and ensure that community living pilots can continue to navigate the skies with confidence and competence.

The only way to know for sure how long NRT works is to do longitudinal trials with long follow-up times.⁴⁶ To better understand the long-term effectiveness of NRT in community living settings, it is important to study how cognitive benefits are maintained and how symptoms of depression and anxiety return over time. An all-encompassing treatment plan for depression and anxiety might be developed by investigating the synergistic benefits of integrating NRT with other evidence-based therapeutic techniques, including psychotherapy or pharmaceuticals. The best way to ensure these therapies have the most therapeutic impact is for researchers to study when, how, and how they should be combined. In order to understand how NRT works, it is essential to understand the neurological bases of its effects. Neuroimaging Methods like electroencephalography (EEG) and functional magnetic resonance imaging (fMRI) should be used in future studies to investigate the structural and functional changes in the brain linked to NRT. This will help us understand the neural plasticity mechanisms at work. When it comes to mental health interventions, cultural considerations and contextual nuances are quite important. To better understand how cultural, linguistic, and social aspects impact treatment involvement and outcomes, future research should centre on culturally competent NRT protocol adaptations. Following up on the success of NRT in randomized controlled trials, future studies should look at how well it works in community mental health settings. If NRT is to be widely used in mental health treatment, it is essential to evaluate its scalability, cost-effectiveness, and compatibility with current systems. Neurocognitive Remediation Therapy has the potential to greatly benefit people living in community settings who suffer from depression and anxiety by improving their cognitive functioning and general mental health if future studies focus on these important areas. Overall, the empirical evidence regarding NRT's efficacy in improving cognition in individuals with depression and anxiety is compelling. Meta-analyses consistently report significant cognitive improvements, and RCTs demonstrate its real-world applicability, particularly in high-stress professions.⁴⁶ Neuroimaging studies confirm that NRT

induces neural changes that underlie these cognitive enhancements. This convergence of evidence highlights the potential of NRT as a valuable therapeutic approach for addressing cognitive deficits in individuals with depression and anxiety, ultimately improving their quality of life and overall well-being.³⁴ Further research and implementation efforts are warranted to maximize the benefits of NRT in diverse contexts.

Future Work

Future Work and Potential Outputs in the Field of Neurocognitive Remediation Therapy for Community Living Pilots with Depression and Anxiety. Conducting long-term studies to assess the sustained effects of neurocognitive remediation therapy (NRT) on community living pilots with depression and anxiety. This could involve follow-up assessments over several months or years to track changes in cognitive functioning, mental health outcomes, and overall well-being. Expanding the participant pool to include a more diverse range of individuals, such as different age groups, genders, and socioeconomic backgrounds. This would help determine the generalizability of NRT across various demographic factors and identify specific subpopulations that may benefit most from this intervention. Conducting comparative studies to evaluate the efficacy of NRT in comparison to other therapeutic interventions for depression and anxiety. This could include traditional psychotherapy, pharmacological treatments, or a combination of therapies. Understanding how NRT measures up against existing treatments will contribute to its evidence base and guide treatment recommendations. Integrating neuroimaging techniques (eg, functional MRI, EEG) to explore the neural mechanisms underlying the effectiveness of NRT. Investigating changes in brain activity and connectivity patterns before and after therapy sessions can provide insights into the neurological changes associated with improved cognitive function and mental health outcomes. Refining and optimizing the existing NRT protocols based on feedback from participants and clinicians. This may involve tailoring interventions to address specific cognitive deficits prevalent in community living pilots with depression and anxiety, ensuring a more personalized and effective treatment approach. Exploring the integration of technology, such as virtual reality or mobile applications, to enhance the accessibility and engagement of NRT. Developing user-friendly platforms that allow individuals to continue cognitive remediation exercises remotely may increase adherence to treatment and extend its reach to those with limited access to traditional therapeutic settings. Conducting cost-benefit analyses to evaluate the economic feasibility of implementing NRT on a larger scale. Assessing the economic impact of improved cognitive functioning and mental health outcomes could inform healthcare policy decisions and encourage the integration of NRT into mainstream mental health care services. Creating training programs for mental health professionals to effectively administer NRT. This involves establishing standardized training protocols, certification processes, and ongoing professional development to ensure the quality and consistency of NRT delivery across different clinical settings.

Acknowledgments

This research was supported by the Education Science Planning Subject of Shaanxi Province for the 14th Five-Year Plan (Grant No. SGH21Y0036).

Funding

This research was supported by the Natural Science Basic Research Program of Shaanxi (Grant No. 2022JZ-13), the Humanities and Social Sciences Project of the Ministry of Education in China (Grant No. 22YJA190001), the Graduate Education Comprehensive Reform Research and Practice Project of Shaanxi Province (Grant No. YJSZG2023047), the Fundamental Research Funds for the Central Universities (Grant No. GK202201010), the Education Teaching Reform Funding of Shaanxi Normal University (Grant No. JSJY202207), and the Project of Undergraduate Teaching and Learning Reform of Shaanxi Normal University (Grant No. 24JBGS17).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Semkowska M, Ahern E. Online neurocognitive remediation therapy to improve cognition in community-living individuals with a history of Depression: a pilot study. *Int Interventions*. 2017;9:7–14. doi:10.1016/j.invent.2017.04.003
2. Eaton AD, Walmsley SL, Craig SL, et al. Protocol for a pilot randomised controlled trial evaluating feasibility and acceptability of cognitive remediation group therapy compared with mutual aid group therapy for people ageing with HIV-associated neurocognitive disorder (HAND) in Toronto, Canada. *BMJ Open*. 2019;9(10):e033183. doi:10.1136/bmjopen-2019-033183
3. Lisoni J, Baldacci G, Nibbio G, et al. Effects of bilateral, bipolar-nonbalanced, frontal transcranial Direct Current Stimulation (tDCS) on negative symptoms and neurocognition in a sample of patients living with schizophrenia: results of a randomized double-blind sham-controlled trial. *J Psychiatr Res*. 2022;155:430–442. doi:10.1016/j.jpsychires.2022.09.011
4. Ayers CR, Saxena S, Espejo E, et al. Novel treatment for geriatric hoarding disorder: an open trial of cognitive rehabilitation paired with behavior therapy. *Am J Geriatric Psychiatry*. 2014;22(3):248–252. doi:10.1016/j.jagp.2013.02.010
5. Preschl B, Wagner B, Forstmeier S, Maercker A. E-health interventions for depression, anxiety disorders, dementia, and other disorders in old age: a review. *J Cyber Therapy Rehabilitation*. 2011;4:371–385.
6. Eaton AD, Craig SL, Rourke SB, et al. Cognitive remediation group therapy compared to mutual aid group therapy for people aging with HIV-associated neurocognitive disorder: randomized, controlled trial. *Social Work with Groups*. 2022;45(2):116–131. doi:10.1080/01609513.2021.1963389
7. Inagawa T, Yokoi Y, Narita Z, et al. Safety and feasibility of transcranial direct current stimulation for cognitive rehabilitation in patients with mild or major neurocognitive disorders: a randomized sham-controlled pilot study. *Front Human Neurosci*. 2019;13:273. doi:10.3389/fnhum.2019.00273
8. Inagawa T, Yokoi Y, Narita Z, et al. P2-004: cognitive Rehabilitation with transcranial direct current stimulation: effect on cognition in major or minor neurocognitive disorders, a randomised controlled preliminary study. *Alzheimer's Dementia*. 2018;14(7S_Part_12):P666. doi:10.1016/j.jalz.2018.06.688
9. Lin L-Y, Chien Y-N, Chen Y-H, et al. Bullying experiences, depression, and the moderating role of resilience among adolescents. *Front Public Health*. 2022;10:872100. doi:10.3389/fpubh.2022.872100
10. Winter R, Leanage N, Roberts N, et al. Experiences of empathy training in healthcare: a systematic review of qualitative studies. *Patient Educ Couns*. 2022;105(10):3017–3037. doi:10.1016/j.pec.2022.06.015
11. London M, Sessa VI, Shelley LA. Developing self-awareness: learning processes for self-and interpersonal growth. *Ann Rev Org Psychol Org Behav*. 2023;10(1):261–288. doi:10.1146/annurev-orgpsych-120920-044531
12. Boyd A. Equity in the Classroom for Individuals with Disabilities at Grand Valley State University (2022). Honors Projects. 902. Available from: <https://scholarworks.gvsu.edu/honorsprojects/902>.
13. Dewji S. The Impact of Emotional Support on Identity in Breast Cancer Survivors (2022). Honors Undergraduate Theses. 1129. Available from: <https://stars.library.ucf.edu/honorstheses/1129>.
14. Miller AB, Schafer L. An Educational Module on a Formalized CRNA Preceptorship Workshop to Enhance Teaching and Communication Skills (2022). Nicole Wertheim College of Nursing Student Projects. 118. Available from: <https://digitalcommons.fiu.edu/cnhs-studentprojects/118>.
15. Bolm SL, Zwaal W, Fernandes MB. Effects of mindfulness on occupational stress and job satisfaction of hospitality and service workers. *Res Hospitality Management*. 2022;12(1):61–70. doi:10.1080/22243534.2022.2080954
16. Blešić I, Pivac T, Kovačić S, et al. Place attachment, cultural involvement, and residents' attitudes towards tourism Development: the case of Novi Sad, the European capital of culture 2022. *Sustainability*. 2022;14(15):9103. doi:10.3390/su14159103
17. Margoni M, Preziosa P, Rocca MA, Filippi M. Depressive symptoms, anxiety and cognitive impairment: emerging evidence in multiple sclerosis. *Transl Psychiatry*. 2023;13(1):264. doi:10.1038/s41398-023-02555-7
18. Shah SMA, Mohammad D, Qureshi MFH, et al. Prevalence, psychological responses and associated correlates of depression, anxiety and stress in a global population, during the coronavirus disease (COVID-19) pandemic. *Commun Mental Health J*. 2021;57:101–110. doi:10.1007/s10597-020-00728-y
19. Escajeda HG. Technology Justice: taxation of Our Collective and Cumulative Cognitive Inheritance. *Loyola LA Law Review*. 2023;56:1073.
20. Ofori-Sasu D, Sarpong-Kumankoma E, Agbloyor EK, Abor JY. Risk-taking behavior, regulatory policy and banking crisis: evidence from the African Economies. *Af Finance J*. 2022;24(2):17–37.
21. Liu Q, He H, Yang J, et al. Changes in the global burden of depression from 1990 to 2017: findings from the Global Burden of Disease study. *J Psychiatr Res*. 2020;126:134–140. doi:10.1016/j.jpsychires.2019.08.002
22. Śniadach J, Szymkowiak S, Osip P, Waszkiewicz N. Increased depression and anxiety disorders during the COVID-19 pandemic in children and adolescents: a literature review. *Lancet Regional Health*. 2021;11(11):1188.
23. Biswas R, Lugo A, Akeroyd MA, et al. Tinnitus prevalence in Europe: a multi-country cross-sectional population study. *Lancet Regional Health*. 2022;12.
24. Tortora F, Hadipour AL, Battaglia S, et al. The role of serotonin in fear learning and memory: a systematic review of human studies. *Br Sci*. 2023;13(8):1197. doi:10.3390/brainsci13081197
25. Crocq MA. The history of generalized anxiety disorder as a diagnostic category. *Dialogues in Clinical Neuroscience*. 2017;19(2):107–116. doi:10.31887/DCNS.2017.19.2/macrocq
26. Rosa-Alcázar AI, Rosa-Alcázar Á, Martínez-Esparza IC, et al. Response inhibition, cognitive flexibility and working memory in obsessive-compulsive disorder, generalized anxiety disorder and social anxiety disorder. *Int J Environ Res Public Health*. 2021;18(7):3642. doi:10.3390/ijerph18073642
27. Mengistu SW, Kassie AW. Household level determinants of food insecurity in rural Ethiopia. *J Food Qual*. 2022;2022:1–9. doi:10.1155/2022/3569950
28. Yang X, Fang Y, Chen H, et al. Global, regional and national burden of anxiety disorders from 1990 to 2019: results from the Global Burden of Disease Study 2019. *Population Health Sci*. 2021;30:e36.
29. Saccaro LF, Schilliger Z, Perroud N, Pigué C. Inflammation, anxiety, and stress in attention-deficit/hyperactivity disorder. *Biomedicines*. 2021;9(10):1313. doi:10.3390/biomedicines9101313
30. Tanaka M, Spekker E, Szabó Á, et al. Modelling the neurodevelopmental pathogenesis in neuropsychiatric disorders. Bioactive kynurenines and their analogues as neuroprotective agents—In celebration of 80th birthday of Professor Peter Riederer. *J Neural Transm*. 2022;129(5):627–642. doi:10.1007/s00702-022-02513-5

31. Higgins V, Sohaei D, Diamandis EP, Prassas I. COVID-19: from an acute to chronic disease? Potential long-term health consequences. *Crit Rev Clin Lab Sci.* 2021;58(5):297–310. doi:10.1080/10408363.2020.1860895
32. Benson C, Singer D, Carpinella CM, Shawi M, Alphas L. The health-related quality of life, work productivity, healthcare resource utilization, and economic burden associated with levels of suicidal ideation among patients self-reporting moderately severe or severe major depressive disorder in a national survey. *Neuropsychiatr Dis Treat.* 2021;111–123. doi:10.2147/NDTS229530
33. Brouwers EP. Social stigma is an underestimated contributing factor to unemployment in people with mental illness or mental health issues: position paper and future directions. *BMC Psychiatry.* 2020;8:1–7.
34. Battaglia MR, Di Fazio C, Battaglia S. Activated Tryptophan-Kynurenine metabolic system in the human brain is associated with learned fear. *Front Mol Neurosci.* 2023;16:1217090. doi:10.3389/fnmol.2023.1217090
35. Gong Y, Liu X, Zheng Y, et al. COVID-19 induced economic slowdown and mental health issues. *Front Psychol.* 2022;13:777350. doi:10.3389/fpsyg.2022.777350
36. Espina C, Feliu A, Maza M, et al. Latin America and the Caribbean Code Against Cancer 1st Edition: 17 cancer prevention recommendations to the public and to policy-makers (World Code Against Cancer Framework). *Cancer Epidemiol.* 2023;86:102402. doi:10.1016/j.canep.2023.102402
37. Søvdal LE, Naslund JA, Kousoulis AA, et al. Prioritizing the mental health and well-being of healthcare workers: an urgent global public health priority. *Front Public Health.* 2021;9:679397. doi:10.3389/fpubh.2021.679397
38. Favver B, DeCouto BS, Trachik B, Dretsch M, Williams AM. Cross-Disciplinary Innovation Within the Intelligence Community: evidence from Research on Sport and Military Expertise. In: *Fostering Innovation in the Intelligence Community: Scientifically-Informed Solutions to Combat a Dynamic Threat Environment.* Springer; 2023:81–112.
39. Santomauro D F, Herrera AMM, Shadid J et al. (2021). Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *The Lancet*, 398(10312), 1700–1712. doi:10.1016/S0140-6736(21)02143-7
40. Tomlinson LS. *Neuroplasticity as a Transformational Tool for Improving Managerial Approaches.* University of Zagreb. Faculty of Economics and Business. Department of; 2021.
41. Marantika JER. Metacognitive ability and autonomous learning strategy in improving learning outcomes. *J Educ Learn.* 2021;15(1):88–96.
42. Henry MA, Shorter S, Charkoudian LK, Heemstra JM, Le B, Corwin LA. Coping behavior versus coping style: characterizing a measure of coping in undergraduate STEM contexts. *Int J STEM Educ.* 2022;9(1):1–26. doi:10.1186/s40594-022-00331-5
43. Coggins IM. *Commonplace and Mirroring for Self-Reflexivity in Secondary Language Arts: A Value-Creative Approach in a Virtual Learning Space.* Columbia University; 2021.
44. Oosthuizen RM. Empowerment of Women Leaders in the Fourth Industrial Revolution. In: *Women's Empowerment for a Sustainable Future: Transcultural and Positive Psychology Perspectives.* Springer; 2023:363–380.
45. Racicot FÉ, Théoret R. Tracking market and non-traditional sources of risks in procyclical and countercyclical hedge fund strategies under extreme scenarios: a nonlinear VAR approach. *Financial Innovation.* 2022;8(1):24. doi:10.1186/s40854-021-00316-3
46. Ellison S, Parker H, Bigger BJ. Advances in therapies for neurological lysosomal storage disorders. *J Inherit Metabol Disease.* 2023;46(5):874–905. doi:10.1002/jimd.12615
47. Miles T. *LITERATURE REVIEW: How are Self-Compassion, Self-Criticism and Self-Blame, Related to Executive Function Skills: A Systematic Review* *EMPIRICAL PAPER: Skills for Self-Compassion After Trauma: The Role of Executive Function.* University of Exeter (United Kingdom); 2020.
48. Slavich GM, Roos LG, Mengelkoch S, et al. Social Safety Theory: conceptual foundation, underlying mechanisms, and future directions. *Health Psychol Rev.* 2023;17(1):5–59. doi:10.1080/17437199.2023.2171900
49. Howlett JR, Nelson LD, Stein MB. Mental health consequences of traumatic brain injury. *Biol Psychiatry.* 2022;91(5):413–420. doi:10.1016/j.biopsych.2021.09.024
50. Galvin JE, Tolea MI, Moore C, Chrisphonte S. The Number Symbol Coding Task: a brief measure of executive function to detect dementia and cognitive impairment. *PLoS One.* 2020;15(11):e0242233. doi:10.1371/journal.pone.0242233
51. Doulou A, Drigas A. Electronic, VR & augmented reality games for intervention in ADHD. *Techn Society J.* 2022;28:159.
52. Merriman NA, Gillan D, Pender NP, et al. The StrokeCog study: development and description of a cognition-focused psychological intervention to address cognitive impairment following stroke. *Psychol Health.* 2020;36(7):792–809. doi:10.1080/08870446.2020.1820009
53. Moreau DJ. How malleable are cognitive abilities? A critical perspective on popular brief interventions. *Annu Psychol.* 2022;77(3):409.
54. Davies S, Milian RP. Maintaining status in new times: the continuing stratification of Anglo-American universities. In: *Routledge Handbook of the Sociology of Higher Education.* Routledge; 2022:32–44.
55. Saif-Alyousfi AY, Saha A. Determinants of banks' risk-taking behavior, stability and profitability: evidence from GCC countries. *Int J Islamic Middle Eastern Fin Manag.* 2021;14(5):874–907.
56. Dang VD. Monetary policy and bank performance: the role of business models. *North Am J Econ Finance.* 2022;59:101602. doi:10.1016/j.najef.2021.101602
57. Versi N, Murphy K, Robinson C, Franklin MJ. Simultaneous Dual-Task Interventions That Improve Cognition in Older Adults: a Scoping Review of Implementation-Relevant Details. *J Aging Res.* 2022;2022:1–21. doi:10.1155/2022/6686910
58. Saleh ZT, Altarawneh FZ, Elshatarat RA, et al. Exploring the Neurobiology of Nicotine Dependence and Its Implications for Tobacco Use Treatment: a Literature Review Article. *Med Humanit.* 2023.
59. Guemghar I, Pires de Oliveira Padilha P, Abdel-Baki A, Jutras-Aswad D, Paquette J, Pomey MP. Social robot interventions in mental health care and their outcomes, barriers, and facilitators: a scoping review. *JMIR Ment Health.* 2022;9(4):e36094. doi:10.2196/36094
60. Tang VM, Goud R, Zawertailo L, et al. Repetitive Transcranial Magnetic Stimulation for Smoking Cessation: next Steps for Translation and Implementation into Clinical Practice. *Psychol Res.* 2023;115340.
61. Timpel P, Oswald S, Schwarz PE, Harst L. Mapping the evidence on the effectiveness of telemedicine interventions in diabetes, dyslipidemia, and hypertension: an umbrella review of systematic reviews and meta-analyses. *J Med Internet Res.* 2020;22(3):e16791. doi:10.2196/16791
62. Wu AC, Donnelly-McLay D, Weisskopf MG, McNeely E, Betancourt TS, Allen JG. Airplane pilot mental health and suicidal thoughts: a cross-sectional descriptive study via anonymous web-based survey. *Environ Health.* 2016;15(1):121. doi:10.1186/s12940-016-0200-6
63. van Hell JG. (2023). The Neurocognitive Underpinnings of Second Language Processing: Knowledge Gains From the Past and Future Outlook. *Language Learning.* 73(S2): 95–138. doi:10.1111/lang.12601

Psychology Research and Behavior Management

Dovepress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/psychology-research-and-behavior-management-journal>