











Correction

Correction: Sugimoto et al. Trends in the Prevalence and Progression of Diabetic Retinopathy Associated with Hyperglycemic Disorders during Pregnancy in Japan. *J. Clin. Med.* 2022, 11, 165

Masahiko Sugimoto ^{1,*}, Kohei Sampa ¹, Hideyuki Tsukitome ¹, Kumiko Kato ¹, Hisashi Matsubara ¹, Shin Asami ², Kaori Sekimoto ³, Shigehiko Kitano ³, Shigeo Yoshida ⁴, Yoshihiro Takamura ⁵, Takao Hirano ⁶, Toshinori Murata ⁶, Miho Shimizu ⁷, Takamasa Kinoshita ⁷, Sentaro Kusuhara ⁸, Osamu Sawada ⁹, Masahito Ohji ⁹, Rina Yoshikawa ¹⁰, Kazuhiro Kimura ¹⁰, Hiroto Ishikawa ¹¹, Fumi Gomi ¹¹, Hiroto Terasaki ¹², Mineo Kondo ¹, Tomoaki Ikeda ¹³ and on behalf of the Writing Committee of Japan-Clinical Retina Study Group (J-CREST) [†]



Citation: Sugimoto, M.; Sampa, K.; Tsukitome, H.; Kato, K.; Matsubara, H.; Asami, S.; Sekimoto, K.; Kitano, S.; Yoshida, S.; Takamura, Y.; et al. Correction: Sugimoto et al. Trends in the Prevalence and Progression of Diabetic Retinopathy Associated with Hyperglycemic Disorders during Pregnancy in Japan. *J. Clin. Med.* 2022, 11, 165. *J. Clin. Med.* 2022, 11, 2789. <https://doi.org/10.3390/jcm11102789>

Received: 28 April 2022

Accepted: 28 April 2022

Published: 16 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

- ¹ Department of Ophthalmology, Graduate School of Medicine, Mie University, Tsu 514-8507, Japan; k.sampa0525@gmail.com (K.S.); dyrkn618@yahoo.co.jp (H.T.); cqw14171jp@yahoo.co.jp (K.K.); hmatsu@clin.medic.mie-u.ac.jp (H.M.); mineo@clin.medic.mie-u.ac.jp (M.K.)
 - ² Graduate School of Medicine, Faculty of Medicine, Mie University, Tsu 514-8507, Japan; 318002@m.mie-u.ac.jp
 - ³ Department of Ophthalmology, Diabetes Center, Tokyo Women's Medical University, Tokyo 162-8666, Japan; sekimoto.dmc@twmu.ac.jp (K.S.); ge2s-ktn@asahi-net.or.jp (S.K.)
 - ⁴ Department of Ophthalmology, Graduate School of Medicine, Kurume University, Kurume 830-0011, Japan; usyosi@gmail.com
 - ⁵ Department of Ophthalmology, School of Medical Sciences, University of Fukui, Yoshida 910-1193, Japan; ytakamura@hotmail.com
 - ⁶ Department of Ophthalmology, School of Medicine, Shinshu University, Matsumoto 390-8621, Japan; takaoh@shinshu-u.ac.jp (T.H.); murata@shinshu-u.ac.jp (T.M.)
 - ⁷ Department of Ophthalmology, Sapporo City General Hospital, Sapporo 060-8604, Japan; miho.shimizu@doc.city.sapporo.jp (M.S.); knst129@gmail.com (T.K.)
 - ⁸ Division of Ophthalmology, Department of Surgery, Graduate School of Medicine, Kobe University, Kobe 650-0017, Japan; kusu@med.kobe-u.ac.jp
 - ⁹ Department of Ophthalmology, Shiga University of Medical Science, Otsu 520-2192, Japan; eye.sawada@gmail.com (O.S.); ohji@belle.shiga-med.ac.jp (M.O.)
 - ¹⁰ Department of Ophthalmology, Graduate School of Medicine, Yamaguchi University, Ube 755-8505, Japan; rinayskw@yamaguchi-u.ac.jp (R.Y.); k.kimura@yamaguchi-u.ac.jp (K.K.)
 - ¹¹ Department of Ophthalmology, Hyogo College of Medicine, Nishinomiya 663-8501, Japan; ohmyeye@gmail.com (H.I.); fgomi@hyo-med.ac.jp (F.G.)
 - ¹² Department of Ophthalmology, Graduate School of Medical and Dental Sciences, Kagoshima University, Kagoshima 890-8520, Japan; teracchi16@yahoo.co.jp
 - ¹³ Department of Obstetrics and Gynecology, Graduate School of Medicine, Mie University, Tsu 514-8507, Japan; t-ikeda@clin.medic.mie-u.ac.jp
- * Correspondence: sugmochi@clin.medic.mie-u.ac.jp; Tel.: +81-59-231-5027
- † Writing Committee of Japan-Clinical Retina Study group (J-CREST) are listed in the data Supplementary Materials.

Error in Table

In the original publication [1], due to an editorial office error, there were mistakes in: “Table 1. The prevalence of DR from the Japanese claim database cohort; Table 2. Complications during pregnancy from Japanese claim database cohort; Table 3. The distribution of DR stage from the multicenter cohort; Table 4. DR progression ratio from multicenter cohort.” as published. “The columns of the tables were divided by mistake”. The corrected “Table 1. The prevalence of DR from the Japanese claim database cohort; Table 2. Complications during pregnancy from Japanese claim database cohort; Table 3. The distribution of DR stage from the multicenter cohort; Table 4. DR progression ratio from multicenter

cohort.” appear below. The authors apologize for any inconvenience caused and state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Table 1. The prevalence of DR from the Japanese claim database cohort.

	Total		DR (+)		DR (−)		Odds Ratio for DR between the Groups	p-Value
	N	Age (Years)	N (%)	Age (Years)	N (%)	Age (Years)		
pexD	1139	34.2 ± 4.6	73 (6.4)	34.4 ± 4.6	1066 (93.6)	34.2 ± 4.6		
GDM	4129	33.7 ± 4.5	61 (1.5)	35.4 ± 4.5	4068 (98.5)	33.7 ± 4.5		
Total	5268	33.8 ± 4.5	134 (2.5)	34.8 ± 4.5	5134 (97.5)	33.8 ± 4.5	4.6 (3.2–6.5) **	1.44 × 10 ^{−15}

DR: diabetic retinopathy, GDM: gestational diabetes mellitus, pexD: pre-existing diabetes mellitus. Values are presented as the mean ± standard deviation. ** *p* < 0.01.

Table 2. Complications during pregnancy from Japanese claim database cohort.

	pexD		GDM		Odds Ratio (95% CI)	p-Value
	DR (+)	DR (−)	DR (+)	DR (−)		
<General complications>						
Hypertension	13 (11.7%)	98 (88.3%)	8 (2.1%)	367 (97.9%)	6.0 (2.5–15.0) **	9.78 × 10 ^{−5}
Renal failure	9 (19.2)	38 (80.9)	1 (1.1)	95 (99.0)	22.5 (2.8–183.7) **	0.003
Eclampsia	0 (0)	0 (0)	0 (0)	13 (100)	Not available	Not available
Caesarean section	25 (11.7)	189 (88.3)	12 (1.5)	812 (98.5)	9.0 (4.4–18.1) **	1.19 × 10 ^{−9}
Induced labor	32 (9.88)	292 (90.1)	6 (1.7)	1537 (98.3)	6.5 (3.8–11.0) **	6.05 × 10 ^{−12}
<Ophthalmic complications>						
Cataract	1 (100%)	0 (0%)	3 (27.3%)	8 (72.7%)	7.3 (0.2–225.9)	0.26
Glaucoma	2 (20.0)	8 (80.0)	4 (13.8)	25 (86.2)	1.6 (0.2–10.2)	0.64
Other	10 (12.8)	68 (87.2)	28 (8.4)	304 (91.6)	1.6 (0.7–3.4)	0.23

DR: diabetic retinopathy, GDM: gestational diabetes mellitus, pexD: pre-existing diabetes mellitus. Values are presented as the mean ± standard deviation. ** *p* < 0.01.

Table 3. The distribution of DR stage from the multicenter cohort.

	N	Age (Years)	NDR	Mild	Moderate	Severe	PDR	Total DR
pexD	119	33.4 ± 4.9	88	17	6	4	4	31 (26.1%) *
GDM	96	34.3 ± 5.3	94	2	0	0	0	2 (2.1%)
ODM	10	30.9 ± 4.6	7	2	1	0	0	3 (30.0%)
Total	225	33.6 ± 5.1	189	21	7	4	4	36 (16.0%)

DR: diabetic retinopathy, GDM: gestational diabetes mellitus, NDR: no diabetic retinopathy, ODM: overt diabetes mellitus, pexD: pre-existing diabetes mellitus, PDR: proliferative diabetic retinopathy. Analysis of covariance (ANCOVA) with Tukey-type multiple comparison as a post hoc test was performed. * *p* < 0.05.

Table 4. DR progression ratio from multicenter cohort.

	N	Age (Years)	Duration (Years)	DR Progression (%)
pexD	102	33.5 ± 4.6	12.9 ± 7.9	10 (9.8)
GDM	42	34.1 ± 5.4		0 (0)
ODM	5	31.6 ± 3.2		2 (40.0)
Total	149	33.6 ± 4.8	12.9 ± 7.9	12 (8.1)

DR: diabetic retinopathy, GDM: gestational diabetes mellitus, ODM: overt diabetes mellitus, pexD: pre-existing diabetes mellitus.

Reference

1. Sugimoto, M.; Sampa, K.; Tsukitome, H.; Kato, K.; Matsubara, H.; Asami, S.; Sekimoto, K.; Kitano, S.; Yoshida, S.; Takamura, Y.; et al. Trends in the Prevalence and Progression of Diabetic Retinopathy Associated with Hyperglycemic Disorders during Pregnancy in Japan. *J. Clin. Med.* **2022**, *11*, 165. [[CrossRef](#)] [[PubMed](#)]