

Implementing hamstring injury prevention programmes remotely: a randomised proof of concept trial

Fearghal P Behan ^{1,2}, Nicol van Dyk ^{3,4,5}, Lance Rane,¹ Emma Thorne,⁶ Alexa Banaghan,⁶ Kevin Gilshan,⁶ Enda Francis Whyte⁶

To cite: Behan FP, van Dyk N, Rane L, *et al.* Implementing hamstring injury prevention programmes remotely: a randomised proof of concept trial. *BMJ Open Sport & Exercise Medicine* 2024;**10**:e001728. doi:10.1136/bmjsem-2023-001728

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjsem-2023-001728>).

Accepted 6 January 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Bioengineering, Imperial College London, London, UK

²Discipline of Physiotherapy, Trinity College Dublin, Dublin, Ireland

³High Performance Unit, Irish Rugby Football Union, Dublin, Ireland

⁴Faculty of Health Science, University of Pretoria, Section Sports Medicine, Pretoria, Gauteng, South Africa

⁵School of Public Health, Physiotherapy and Sports Science, University College Dublin, Dublin, Ireland

⁶School of Health and Human Performance, Dublin City University, Dublin, Ireland

Correspondence to

Dr Fearghal P Behan;
F.Behan@imperial.ac.uk

ABSTRACT

Objectives This study aimed to (1) compare the effectiveness of a Nordic hamstring exercise (NHE) versus single-leg Romanian deadlift (SLRDL) exercise programme on a hamstring injury risk surrogate; (2) compare the muscle soreness experienced by both exercise programmes; and (3) assess compliance to remote injury prevention exercise protocols through video software.

Methods Twenty participants (10 women and 10 men: 21.45±1.6 years; 176±23 cm; 70±10 kg) were randomised into an NHE or SLRDL programme for 6 weeks. Single-leg hamstring bridge (SLHB), a hamstring injury surrogate, was the primary outcome for exercise efficacy. Muscle soreness and exercise adherence were also assessed. Significance was set at $p < 0.05$.

Results Both exercises increased SLHB performance resulting in an overall effect ($p = 0.013$) with no effect for group ($p = 0.470$) and no interaction effect ($p = 0.709$), indicating both groups improved but there was no difference in improvement between interventions. There was no difference in muscle soreness between groups ($p = 0.087$). Finally, both groups had 100% adherence to the programme.

Conclusions Both the NHE and SLRDL are equally effective in increasing SLHB performance and demonstrate a similar level of muscle soreness. This suggests that SLRDL may be a viable option as a preventative exercise to mitigate the risk of hamstring injury. Finally, implementing injury prevention programmes remotely has the potential to enhance adherence.

INTRODUCTION

Hamstring injuries continue to be a persistent problem during sporting activities, with little change in incidence rates over time.¹ These injuries account for 24% of all injuries in elite European men's football,² with similar hamstring injury rates in females and males in field sports.³

The Nordic hamstring exercise (NHE) can reduce hamstring injury incidence by up to 51%.⁴ However, this exercise is poorly adopted and adhered to across elite football,⁵ as it is deemed to cause muscle soreness.⁶ Therefore, the pursuit of potential alternative exercises seems prudent.

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ The Nordic hamstring exercise (NHE) reduces the risk of hamstring strain injury but adherence to this exercise has been poor.

WHAT THIS STUDY ADDS

⇒ Conducting an NHE programme remotely resulted in 100% adherence. Furthermore, an alternative exercise, the single-leg Romanian deadlift, also achieved 100% adherence. Additionally, both exercises improved performance on a hamstring strain injury surrogate measure.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ This evidence suggests that conducting injury prevention remotely over web-enabled video technology may be a viable alternative to on-site programmes.

The deadlift is a staple of hamstring strengthening routines, and of its variations, the single-leg Romanian deadlift (SLRDL) offers specific benefits. As a single-leg exercise, the SLRDL adds an extra balance requirement by reducing the base of support⁷ while also increasing eccentric biceps femoris activation over its bilateral equivalent.⁸ Additionally, the SLRDL targets the hamstrings more specifically, as evidenced by a higher ratio of biceps femoris and semitendinosus to erector spinae activation, compared with other deadlift variations.⁹ The SLRDL may present an alternative to the NHE.

The progression in internet-based video technology facilitates a potential alternative medium¹⁰ to implement and monitor injury prevention exercises. Routines involving the SLRDL and the NHE are conducive to remote delivery, and the remote setting has the advantage of wider and less costly implementation. It also allows delivery to athletes who live away from their team's location due to education/work or within elite populations, due to international duty. The difficulty of assessing on-site injury-related outcomes

may be overcome by using surrogate measures with a relationship to injury that can be assessed remotely to act as a substitute for measuring prospective hamstring strain injuries directly when this is not feasible. An appropriate surrogate measure may be the single-leg hamstring bridge (SLHB) test as it is possible to implement remotely and has been shown to have predictive validity for hamstring strain injury.¹¹

Therefore, this study aimed to (1) compare the effectiveness of an NHE versus SLRDL programme on a hamstring injury risk surrogate; (2) compare muscle soreness experienced by both exercise programmes; and (3) assess compliance to remote injury prevention protocols.

METHODS

Participants

Twenty individuals (10 women and 10 men) were recruited for this study. We recruited through email and word of mouth. Inclusion criteria consisted of healthy, active men and women, aged between 18 and 25 years. Exclusion criteria comprised a history of hamstring or lower limb injury in the last year, any strength training contraindications or participation in any concurrent lower limb resistance training.

Study design

All assessments and interventions in this randomised interventional trial were conducted remotely through web-enabled video software (Zoom Video

Communication, San Jose, California, USA). Participants were located at a variety of diverse geographical locations throughout the island of Ireland. Participants underwent an initial SLHB test prior to allocation to either NHE or SLRDL, with 50:50 block randomisation used to maintain gender balance. Prior to the SLHB test, participants were familiarised with the exercise technique. Following group allocation, participants were familiarised with their specific exercise prior to commencing their intervention sessions. Participants completed two weekly sessions of their allocated exercise for 6 weeks with increasing volumes to maintain progressive overload (online supplemental material 1).^{12 13} The investigators remotely supervised all sessions. Three separate investigators supervised in a crossover fashion, with each supervisor alternating their intervention group on a weekly basis. Delayed-onset muscle soreness (DOMS) values were assessed during each session using a verbal Numerical Rating Scale of 0–10, with the peak weekly value used for analysis.¹⁴ After 6 weeks, participants completed a final SLHB test.

SLHB test

This test was chosen as it has demonstrated predictive validity for hamstring injury risk,¹¹ can be monitored remotely and requires minimal equipment.

Participants lay supine with the heel of their foot on a stable surface 60 cm in height. Participants were asked to measure the height of the equipment they would place their heel on (eg, table, desk) to ensure the height was as close to 60 cm as possible. They repeated their post-test using the same equipment. With the test leg in 20° of knee flexion, they crossed their arms over their chest and pushed through their heel to raise the pelvis off the floor and return to the starting position.¹¹ This was repeated until the subject was unable to continue, and the number of complete repetitions was counted. During the test, verbal feedback was given to ensure that the athlete touched the floor with their pelvis without resting it on the ground, before extending their hip to 0°. Further feedback was given to ensure controlled and consistent duration of eccentric and concentric phases for each repetition in all participants. Both legs were tested, with starting legs alternated between participants. As the investigator completing the assessment has supervised each group due to crossover supervision of the intervention, blinded assessment was not possible.

Nordic hamstring exercise

Participants knelt on a soft surface and kept ankles stabilised either through partner assistance or by securing the ankles under a heavy object (eg, large furniture). Participants crossed their arms over their chest with their hips extended and lowered their body, controlling the descent as far as possible until reaching the floor.¹⁵ Those able to control the movement in the final 10–20° were required to hold a weight plate to the chest to ensure intensity.^{15 16}




Figure 1 Single-leg hamstring bridge (SLHB) box plot results for the Nordic hamstring exercise (NHE) and single-leg Romanian deadlift (SLRDL) groups. Pre-intervention results are in hatched grey, while post-intervention results are plain white boxes. Dotted lines represent individual change across time points.

Single-leg Romanian deadlift

Participants stood erect with their legs–shoulder width apart while holding a weight plate. The non-working leg was lifted slightly above the ground and the stance leg knee flexed to approximately 15°. Participants lowered the plate through hip flexion of the weight-bearing leg, while the non-working leg extended backwards. The weight was lowered until the plate reached the distal tibia¹⁷; the hips were then extended to return to the starting position. Failure was deemed as a point in the repetition range where participants were unable to maintain technical control or unable to return to the starting position after the lowering phase of the exercise. The resistance was adjusted weekly to ensure failure at the predetermined final repetition. Participants supplied their own weights to ensure consistent overload.

Statistical analysis

Analysis was conducted using Stata V.16 (StataCorp, Texas, USA). Normal data distribution was determined by the Shapiro-Wilk test. The primary dependent variable was number of completed SLHB repetitions; as there was no statistical difference between limbs, the average score of both legs was used. A two-way (2×2) mixed-model analysis of variance was used to examine the effect of the intervention group (NHE vs SLRDL), time (pre-intervention vs post-intervention) and interaction. Within-group comparisons were analysed using a paired-sample t-test. The secondary dependent variable was DOMS score. A Kruskal-Wallis test assessed the differences between interventional exercises on DOMS. Compliance was assessed through the number of missed supervised training sessions as a percentage. Statistical significance was set at $p < 0.05$. Calculations of sample size were performed a priori. A sample size of 20 participants was deemed sufficient based on an effect size of 1.2, power set at 80% and an alpha level of < 0.050 .¹⁵

RESULTS

Group characteristics

Twenty individuals (10 women and 10 men: 21.45 ± 1.6 years; 176 ± 23 cm; 70 ± 10 kg) completed this study. All participants exercised recreationally two to three times per week, with 14 of these participating in regional-level competitive sports (Gaelic football, camogie, hurling, soccer). None had any history of hamstring strain injury or any lower limb injury in the previous year. Participants were not participating in any concurrent lower limb resistance training, but otherwise were not asked to alter their normal activity.

SLHB test

Both groups increased their SLHB score post-intervention (NHE (mean±SD): pre: 26 ± 12 (95% CI: 18 to 35) vs post: 34 ± 10 (95% CI: 27 to 41), Cohen's $d = 0.67$; SLRDL: pre: 28 ± 9 (95% CI: 21 to 34) vs post: 38 ± 11 (95% CI: 30 to 45), Cohen's $d = 1.00$, figure 1) resulting in an overall effect for time (95% CI for SLHB difference by time: 2




Figure 2 Weekly delayed-onset muscle soreness (DOMS) results (mean±SD) for the Nordic hamstring exercise (NHE, black) and single-leg Romanian deadlift (SLRDL, grey) groups.

to 16, $p = 0.013$). There was no effect for group (95% CI for SLHB difference by group: -5 to 10 , $p = 0.470$) and no interaction effect ($p = 0.709$), indicating that both groups improved but there was no difference in improvement between interventions.

Delayed-onset muscle soreness

There was no difference between groups for DOMS ($p = 0.087$, figure 2). DOMS decreased weekly in both groups ($p < 0.001$).

External load

The median external load required for the NHE group was 0 kg (range: 0–5 kg), with only one participant requiring external load, while the median external load for the SLRDL group was 7.5 kg (range: 0–15 kg), with all participants requiring external load.

Compliance

Neither group had any participant miss a supervised interventional session, resulting in 100% compliance in both groups.

DISCUSSION

Both the NHE and SLRDL increased hamstring bridge performance, with neither exercise found to be superior. Conducting these exercises remotely resulted in 100% compliance for both interventions.

An improvement in SLHB performance is of significance due to its predictive validity for hamstring injury risk.¹¹ It has previously been demonstrated that athletes with a lower SLHB score were more likely to sustain a hamstring injury in the following season.¹¹ The current study represents, to the authors' knowledge, the first evaluation of SLRDL effects on indirect hamstring injury indicators, such as the SLHB. Similar exercises have been compared with NHE previously with no differences being found,¹⁸ in keeping with our results. However, neither the SLRDL or previous alternatives (45° hip extension)¹⁹ have been proven to have direct effects on preventing hamstring injuries.^{4,12,13} Considering the relatively higher biceps femoris long head force production during deadlifts compared with the NHE,¹⁸ future trials assessing

the preventive effects of these alternative exercises on hamstring injury risk appear warranted.

Despite the repeated demonstration of positive outcomes resulting from NHE programmes, compliance in elite sport remains poor,^{5,6} an issue of pertinence given the importance of compliance to the effectiveness of injury prevention.²⁰ Poor compliance may be linked to a high repetition number prescribed by the seminal randomised control trials in the area,^{12,13} but our study adhered to a high-volume prescription and maintained a 100% adherence rate. A second proposed factor causing poor compliance is muscle soreness.⁶ However, we observed minimal muscle soreness during both interventions, with no significant difference between them. DOMS experienced during the NHE programme in the current study appears lower than what was reported in previous work with similar dosage^{14,19}; whether this is a consequence of the remote delivery remains speculative. The lack of difference in DOMS values suggests that if DOMS is the main reason for avoidance, SLRDL at the current dosage may not result in enhanced compliance over NHE. However, issues of compliance are not restricted to NHE.²¹ For all interventions, remote delivery offers benefits including time flexibility and convenience, and may be more cost-effective, notwithstanding possible additional time required of supervising staff, as group delivery of intervention may be precluded by the remote format.^{12,13}

Limitations

Despite the success of the interventions on SLHB and full compliance achieved, the intervention has limitations. Other potentially relevant hamstring injury measures, such as muscle architecture,²² were not assessed due to the remote delivery of the trial. There were no prospective follow-ups to assess whether intervention improved future hamstring injuries. Comparing ratings of perceived exertion would have allowed a comparison of the intensity of the interventions between groups. Furthermore, blinded assessment would have reduced potential bias. Finally, no performance measures were assessed, despite both exercises previously demonstrating a potential benefit on athletic attributes.^{23,24} In elite sport, it may be the ergogenic rather than preventive benefits of an exercise that ultimately result in its inclusion among an athlete's programme.

Clinical implications

Both the NHE and SLRDL are equally effective in increasing SLHB performance and demonstrate a similar level of muscle soreness. This suggests that SLRDL may be a viable option as a preventative exercise to mitigate the risk of hamstring injury. Finally, implementing injury prevention programmes remotely has the potential to enhance compliance.

Twitter Nicol van Dyk @nicolvandyk

Acknowledgements The authors wish to thank the participants for their time and effort.

Contributors FPB, EFW, AB, KG and ET conceived of the study. FPB, AB, KG and ET—data analysis and initial draft. LR, NvD and EFW—additional analysis and manuscript revisions. All authors—final manuscript approval.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Ethics approval This study involves human participants and was approved by the Dublin City University research ethics board (2020_09_FB_ATT). Digital written informed consent was provided prior to the study.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Fearghal P Behan <http://orcid.org/0000-0001-8714-3116>

Nicol van Dyk <http://orcid.org/0000-0002-0724-5997>

REFERENCES

- Maniar N, Carmichael DS, Hickey JT, *et al*. Incidence and prevalence of hamstring injuries in field-based team sports: a systematic review and meta-analysis of 5952 injuries from over 7 million exposure hours. *Br J Sports Med* 2023;57:109–16.
- Ekstrand J, Bengtsson H, Waldén M, *et al*. Hamstring injury rates have increased during recent seasons and now constitute 24% of all injuries in men's professional football: the UEFA Elite Club Injury Study from 2001/02 to 2021/22. *Br J Sports Med* 2022;57:292–8.
- Mullins K, Mac Colgáin D, Carton P. Incidence and severity of hamstring injuries in female athletes who play field sports: a systematic review with meta-analysis of prospective studies. *J Orthop Sports Phys Ther* 2022;52:740–A5.
- van Dyk N, Behan FP, Whiteley R. Including the nordic hamstring exercise in injury prevention programmes halves the rate of hamstring injuries: a systematic review and meta-analysis of 8459 athletes. *Br J Sports Med* 2019;53:1362–70.
- Ekstrand J, Bengtsson H, Waldén M, *et al*. Still poorly adopted in male professional football: but teams that used the nordic hamstring exercise in team training had fewer hamstring injuries - a retrospective survey of 17 teams of the UEFA Elite Club Injury Study during the 2020-2021 season. *BMJ Open Sport Exerc Med* 2022;8:e001368.
- Bahr R, Thorborg K, Ekstrand J. Evidence-based hamstring injury prevention is not adopted by the majority of champions league or norwegian premier league football teams: the nordic hamstring survey. *Br J Sports Med* 2015;49:1466–71.
- Marcori AJ, Monteiro PHM, Oliveira JA, *et al*. Single leg balance training: a systematic review. *Percept Mot Skills* 2022;129:232–52.
- Bourne MN, Williams MD, Opar DA, *et al*. Impact of exercise selection on hamstring muscle activation. *Br J Sports Med* 2017;51:1021–8.
- Martín-Fuentes I, Oliva-Lozano JM, Muyor JM. Electromyographic activity in deadlift exercise and its variants. A systematic review. *PLoS One* 2020;15:e0229507.
- Stoessl AJ, Bhatia KP, Merello M. Movement disorders in the world of COVID-19. *Mov Disord Clin Pract* 2020;7:355–6.
- Freckleton G, Cook J, Pizzari T. The predictive validity of a single leg bridge test for hamstring injuries in Australian rules football players. *Br J Sports Med* 2014;48:713–7.
- Petersen J, Thorborg K, Nielsen MB, *et al*. Preventive effect of eccentric training on acute hamstring injuries in men's

- soccer: a cluster-randomized controlled trial. *Am J Sports Med* 2011;39:2296–303.
- 13 van der Horst N, Smits D-W, Petersen J, *et al.* The preventive effect of the nordic hamstring exercise on hamstring injuries in amateur soccer players. *Am J Sports Med* 2015;43:1316–23.
 - 14 Behan FP, Opar DA, Vermeulen R, *et al.* The dose-response of pain throughout a nordic hamstring exercise intervention. *Scand J Med Sci Sports* 2023;33:542–6.
 - 15 Presland JD, Timmins RG, Bourne MN, *et al.* The effect of nordic hamstring exercise training volume on biceps femoris long head architectural adaptation. *Scand J Med Sci Sports* 2018;28:1775–83.
 - 16 Behan FP, Vermeulen R, Whiteley R, *et al.* The dose-response of the nordic hamstring exercise on biceps femoris architecture and eccentric knee flexor strength: a randomized interventional trial. *Int J Sports Physiol Perform* 2022;17:646–54.
 - 17 Lee S, Schultz J, Timgren J, *et al.* An electromyographic and kinetic comparison of conventional and romanian deadlifts. *J Exerc Sci Fit* 2018;16:87–93.
 - 18 Van Hooren B, Vanwanseele B, van Rossom S, *et al.* Muscle forces and fascicle behavior during three hamstring exercises. *Scand J Med Sci Sports* 2022;32:997–1012.
 - 19 Bourne MN, Duhig SJ, Timmins RG, *et al.* Impact of the nordic hamstring and hip extension exercises on hamstring architecture and morphology: implications for injury prevention. *Br J Sports Med* 2017;51:469–77.
 - 20 Ripley NJ, Cuthbert M, Ross S, *et al.* The effect of exercise compliance on risk reduction for hamstring strain injury: a systematic review and meta-analyses. *Int J Environ Res Public Health* 2021;18:21.
 - 21 van Reijen M, Vriend I, van Mechelen W, *et al.* Compliance with sport injury prevention interventions in randomised controlled trials: a systematic review. *Sports Med* 2016;46:1125–39.
 - 22 Timmins RG, Bourne MN, Shield AJ, *et al.* Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): a prospective cohort study. *Br J Sports Med* 2016;50:1524–35.
 - 23 Ishoi L, Hölmich P, Aagaard P, *et al.* Effects of the nordic hamstring exercise on sprint capacity in male football players: a randomized controlled trial. *J Sports Sci* 2018;36:1663–72.
 - 24 Morencos E, González-Frutos P, Rivera C, *et al.* Effects of six weeks of flywheel single-leg romanian deadlift training on speed, jumping and change of direction performance. *Int J Environ Res Public Health* 2022;19:1200.