

## ARTICLE

# Odds of Acute Kidney Injury in Patients Receiving Dipeptidyl Peptidase 4 Inhibitors: A National Cohort Study Within the Department of Veterans Affairs

S. Scott Sutton<sup>1,2,\*</sup> , Joseph Magagnoli<sup>1,2</sup>, Tammy H. Cummings<sup>2</sup> and James W. Hardin<sup>2,3</sup>

Preclinical and clinical data of dipeptidyl peptidase 4 (DPP-4) inhibitors have demonstrated discordant data regarding acute kidney injury (AKI). Therefore, we aimed to evaluate the association between DPP-4 use and AKI. This cohort study utilized data from the Department of Veterans Affairs evaluating patients diagnosed with type 2 (T2) diabetes with a DPP-4 inhibitor and compared with nondiabetic and diabetic patients. The primary end point is the development of AKI, and statistical analyses were performed to examine the association. DPP-4 use is associated with a lower odds of AKI compared with diabetics (adjusted odds ratio (OR) = 0.39; 95% confidence interval (CI) = 0.32–0.48) and nondiabetics (OR = 0.64; 95% CI = 0.52–0.79). DPP-4 use in patients with T2 diabetes mellitus is associated with lower odds of AKI within 120 days compared with nondiabetic and diabetic controls when adjusting for study covariates.

## Study Highlights

### WHAT IS THE CURRENT KNOWLEDGE ON THE TOPIC?

✓ Because the dipeptidyl peptidase 4 (DPP-4) enzyme is expressed on non-glucagon-like peptide-1 cell types, nondiabetes medication effects of the DPP-4 inhibitors are possible. Preclinical and clinical data for DPP-4 inhibitors have demonstrated a nephroprotective effect; however, clinical data have also demonstrated that DPP-4 inhibitor use was associated with a higher risk of hospitalization for acute kidney injury (AKI).

### WHAT QUESTION DID THIS STUDY ADDRESS?

✓ The purpose of this study is to evaluate patients with type 2 diabetes mellitus with a prescription for a DPP-4 inhibitor vs. diabetic and nondiabetic patients and the associated odds for developing AKI. Research question: What are the odds of AKI in patients with diabetes receiving DPP-4 inhibitors compared

with diabetics not receiving DPP-4 inhibitors and nondiabetics?

### WHAT DOES THIS STUDY ADD TO OUR KNOWLEDGE?

✓ This study used a large nationwide cohort of US Veterans to examine the association of AKI between DPP-4 inhibitors and matched control patients. This study adds additional literature to previous published discordant results.

### HOW MIGHT THIS CHANGE CLINICAL PHARMACOLOGY OR TRANSLATIONAL SCIENCE?

✓ This study can assist prescribers and payers in understanding real-world experience with this medication. There were lower odds of AKI in the DPP-4 cohort compared with the diabetic and nondiabetic controls when using adjusted models. However, future research should be considered in understanding the risks and benefits of DPP-4 use and the occurrence of AKI.

Diabetes affects more than 29 million people in the United States, and the majority of diabetes cases are type 2 diabetes mellitus (T2DM).<sup>1</sup> Diabetes is a risk factor for acute kidney injury (AKI), and AKI episodes may lead to electrolyte abnormalities, pulmonary edema, chronic kidney disease, and cardiovascular complications.<sup>2–4</sup> Dipeptidyl peptidase 4 (DPP-4) inhibitors are a class of oral hypoglycemic medications that are used to manage patients with T2DM.<sup>5–8</sup> DPP-4 is an enzyme expressed on the surface of most cell types that deactivates bioactive peptides, including

glucagon-like peptide-1 (GLP-1). DPP-4 inhibitors affect glucose control through enhancement of glucose-dependent insulin secretion, slowed gastric emptying, and reduction of postprandial glucagon and of food intake. DPP inhibitors are utilized as adjunctive drug therapy or monotherapy for patients inadequately controlled or who cannot take/tolerate other oral diabetes medications. Because the DPP-4 enzyme is also expressed on non-GLP-1 cell types, nondiabetes medication effects of the DPP-4 inhibitors are possible. For example, DPP-4 is also bound on the kidney

<sup>1</sup>College of Pharmacy, Department of Clinical Pharmacy and Outcomes Sciences, University of South Carolina, Columbia, South Carolina, USA; <sup>2</sup>Dorn Research Institute, WJB Dorn Veterans Affairs Medical Center, Columbia, South Carolina, USA; <sup>3</sup>Department of Epidemiology & Biostatistics, University of South Carolina, Columbia, South Carolina, USA. \*Correspondence: S. Scott Sutton (Sutton@cop.sc.edu)

Received: January 30, 2019; accepted: June 10, 2019. doi:10.1111/cts.12676

proximal tubular cell, and microvesicle-bound DPP-4 secreted from tubular epithelial cells is found in the urine and may be an early marker of kidney damage.<sup>9</sup> Additionally, renal effects of DPP-4 inhibitors may be mediated by increased GLP-1 levels.<sup>10</sup> In addition to the pancreas, the GLP-1 receptor is expressed in glomerular endothelial cells, and renal effects of DPP-4 inhibitors are, in part, mediated by GLP-1 levels.<sup>11</sup> However, the exact role and importance of the mechanism of kidney effects relating to the DPP-4 enzyme or DPP-4 inhibitors are not completely understood. Preclinical and clinical data for DPP-4 inhibitors have demonstrated a nephroprotective effect.<sup>12–15</sup> However, there have also been clinical data that demonstrated DPP-4 inhibitor use was associated with a higher risk of hospitalization for AKI.<sup>16</sup> The discordant study results warrant the need for more research evaluating DPP-4 inhibitors and AKI. Therefore, the purpose of this study is to evaluate patients with T2DM with a prescription for a DPP-4 inhibitor vs. diabetic and nondiabetic patients and the associated odds for developing AKI within the Department of Veterans Affairs (VA) Veterans Health Administration system.

## METHODS

### Data

This retrospective cohort study was conducted using data from the Department of Veterans Affairs. The Veterans Affairs Informatics and Computing Infrastructure was utilized to obtain individual-level information on demographics, administrative claims, and pharmacy dispensation. The completeness, utility, accuracy, validity, and access methods are described on the VA website, <http://www.virec.research.va.gov>. The study was conducted in compliance with the Department of Veterans Affairs' requirements and received institutional review board and research and development approval. The study utilized diagnosis International Classification of Disease 9th revision (ICD-9-CM) data from October 1999 to September 2015, laboratory data from January 2001 to June 2016, and pharmacy data from December 1999 to June 2016.

### Cohort selection

Cases were defined as patients who were diagnosed with T2DM (during October 1999 to September 2015) and had a prescription for a DPP-4 inhibitor (including sitagliptin, saxagliptin, or linagliptin) between the dates June 2005 and February 2016. The study index period for the DPP-4 cases was the earliest date the patient was on a DPP-4 and had an estimated glomerular filtration rate (eGFR) laboratory measure. The eGFR is a calculated laboratory from the patients' medical record and was captured via claims data by Logical Observation Identifiers Names and Codes code 33914. Nondiabetic controls were defined as patients who never had an ICD-9-CM diagnosis of T2DM and never had an A1c laboratory result. Uncontrolled blood sugar was defined as an A1c value  $\geq 7$  for the nondiabetic controls. Diabetic non-DPP-4 controls were defined as those who had a diagnosis of T2DM but never had a prescription for a DPP-4 inhibitor. To ensure adequate medical surveillance and baseline kidney function, patients who never had an eGFR measure while on a DPP-4 were excluded.

Furthermore, controls who never had an eGFR laboratory result during the study, June 2005 to February 2016, were excluded.

### Matching

A matching algorithm was utilized that matched DPP-4 cases to controls. A variable number of matched controls per case was utilized, with one control being the minimum and seven controls as a maximum. Controls were exact matched on age at index, race, sex, and constrained controls to be within 10 mL/minute of baseline eGFR. Control baseline eGFR measures were also required to be in the same month and year as their matched case.

### AKI outcome

Cases and controls were followed from index to 120 days or AKI incident, whichever occurred first. Specifically, AKI was identified from an outpatient acute kidney diagnosis using ICD-9-CM: values were 584 (acute kidney failure), 584.5 (acute kidney failure with lesion of tubular necrosis), 584.6 (acute kidney failure with lesion of renal cortical necrosis), 584.7 (acute kidney failure with lesion of renal medullary necrosis), 584.8 (acute kidney failure with other specified pathological lesion in kidney), and 584.9 (acute kidney failure, unspecified).

### Study variables

In addition to our primary predictor (DPP-4 inhibitor use), covariates were utilized to account for possible differences in demographic, disease burden, pharmacy, and laboratory results between the cohorts. Demographic variables included age, sex, and race coded as white, black, other/unknown. The Charlson comorbidity index was utilized to account for differences in disease burden. The Charlson index utilizes all outpatient claims for the 1 year preceding the study index. Additionally, other pharmacy medications were evaluated and included metformin use during the follow-up period, nonsteroidal anti-inflammatory drugs (NSAIDs), and proton pump inhibitors (PPIs).

### Statistical analysis

The statistical analysis occurred in two steps. The initial step utilized statistical tests to compare baseline characteristics between the DPP-4 and control cohorts. For continuous variables, such as age, eGFR, and the Charlson comorbidity index, we utilized the *F*-test from the analysis of variance results. All categorical variables were analyzed using the  $\chi^2$  test. In the second step, we estimated logistic regression models to examine the association between DPP-4 use and AKI. We estimate three models: (i) unadjusted, with only the treatment indicator (diabetic DPP-4, diabetic non-DPP-4, and nondiabetic); (ii) a second adjusted for the Charlson index; and (iii) a model with all covariates (which included treatment indicator, Charlson index, baseline eGFR, uncontrolled A1c, sex, race, age, PPI use, NSAID use, metformin use, and index year). Unadjusted odds of AKI within 120 days for comparisons between cohort groups using a Bonferroni multiplicity adjusted interval were also tested for each model in this step. All data management and matching was done with SAS enterprise guide 7.1 (SAS Institute Cary,

NC), and all statistical analysis was done using R (R Core Team 2016, Vienna, Austria).

**RESULTS**

A total of 363,873 patients met all the study criteria. Of these, 250,269 patients made up the nondiabetic control group, 77,722 the diabetic non-DPP-4 group, and 35,882 the DPP-4 inhibitor cohort. **Table 1** displays the summary statistics for the three cohorts. The DPP-4 and control cohorts are clinically similar in terms of age (66.7 (10.1) vs. 66.8 (9.9) vs. 66.5 (10.2)), race (68% white), and sex (96% men). The DPP-4 cohort, on average, has a lower eGFR measure at index (69.5 (24.4) vs. 69.8 (23.3) vs. 72.2 (22.9) mL/minute). The diabetic non-DPP-4 control group, on average, had a higher Charlson index (2.8 (2.3) vs. 2.5 (1.9) vs. 0.8 (1.5)) compared with the diabetic DPP-4 cases and nondiabetic controls. Approximately 56% of the DPP-4 cohort had an uncontrolled A1C (defined as A1c ≥ 7) at baseline, and 55% were on metformin during the follow-up period. Additionally, 14% of the DPP-4 and diabetic non-DPP-4 cohorts and 13% of the nondiabetic

control cohort had a prescription for NSAIDs, whereas 32% of both DPP-4 and diabetic non-DPP-4 cohorts and 21% of the nondiabetic controls had a prescription for PPIs. Of the patients on a DPP-4 inhibitor, 61.8% of patients had a prescription for saxagliptin, 44.8% had a prescription for sitagliptin, and 12.9% had a prescription for linagliptin. Patients may have had a prescription for more than one DPP-1, and as a result, the percentages do not sum to 100.

**Table 2** presents the occurrence of AKI within 120 days for each cohort. The DPP-4 cohort had 274 incidents of AKI equaling 7.63 (95% confidence interval (CI) = 6.74–8.52) per 1,000 people, the diabetic non-DPP-4 cohort had 21,784 incidents of AKI at 22.95 (95% CI = 21.98–24.04) incidents per 1,000 people, and the nondiabetic cohort had 1,786 incidents of AKI equaling 7.13 (95% CI = 6.80–7.45) incidents per 1,000 people.

The unadjusted logistic regression (**Table 3**, model 1) demonstrates that patients with diabetes not on a DPP-4 inhibitor had a 3.05-fold increased odds of AKI compared with diabetics on a DPP-4 within 120 days (odds ratio (OR) = 3.05; 95% CI = 2.69–3.47), whereas there was no

**Table 1 Patient characteristics**

	Diabetic non-DPP-4	Diabetic DPP-4	Nondiabetic	P value
Variable	N = 77,722	N = 35,882	N = 250,269	
Age, mean (SD)	66.8 (9.9)	66.7 (10.1)	66.5 (10.2)	< 0.001
Race				
Black	13,451 (17.3%)	6,009 (16.7%)	41,390 (16.5%)	< 0.001
Other/unknown	11,274 (14.5%)	5,401 (15.1%)	38,386 (15.3%)	
White	52,997 (68.2%)	24,472 (68.2%)	170,493 (68.1%)	
Sex				
Male	75,220 (96.8%)	34,484 (96.1%)	240,436 (96.1%)	< 0.001
Baseline eGFR, mean (SD)	69.8 (23.3)	69.5 (24.4)	72.2 (22.9)	< 0.001
Charlson index, mean (SD)	2.8 (2.3)	2.5 (1.9)	0.8 (1.5)	< 0.001
Uncontrolled A1c	23,326 (30.0%)	20,194 (56.3%)	0 (0%)	< 0.001
NSAID prescription	11,013 (14.2%)	5,064 (14.1%)	33,682 (13.5%)	< 0.001
PPI prescription	25,164 (32.4%)	11,325 (31.6%)	52,263 (20.8%)	< 0.001
Metformin prescription	28,222 (36.3%)	19,742 (55.0%)	0 (0%)	< 0.001
Index year				
2005	15 (0.02%)	7 (0.02%)	89 (0.03%)	0.24
2006	27 (0.035%)	13 (0.036%)	159 (0.064%)	
2007	348 (0.448%)	161 (0.449%)	1,162 (0.464%)	
2008	1,850 (2.38%)	864 (2.41%)	6,042 (2.41%)	
2009	2,958 (3.81%)	1,367 (3.81%)	9,522 (3.805%)	
2010	3,976 (5.12%)	1,840 (5.13%)	12,852 (5.14%)	
2011	5,901 (7.59%)	2,700 (7.53%)	18,660 (7.46%)	
2012	8,503 (10.94%)	3,926 (10.94%)	27,086 (10.82%)	
2013	8,377 (10.78%)	3,864 (10.78%)	26,553 (10.61%)	
2014	14,938 (19.22%)	6,897 (19.22%)	48,003 (19.18%)	
2015	25,814 (33.21%)	11,923 (33.23%)	83,993 (33.56%)	
2016	5,015 (6.45%)	2,320 (6.47%)	16,148 (6.45%)	
DPP-4 medication <sup>a</sup>				
Saxagliptin	–	22,207 (61.8%)	–	
Sitagliptin	–	16,092 (44.8%)	–	
Linagliptin	–	4,664 (12.9%)	–	

DPP-4, dipeptidyl peptidase 4; eGFR, estimated glomerular filtration rate; NSAID, nonsteroidal anti-inflammatory drugs; PPI, proton pump inhibitor.  
<sup>a</sup>Percentages do not equal 100 because select patients may have received different DPP inhibitors.

statistically significant difference between the nondiabetic control group and the diabetic DPP-4 users (OR = 0.93; 95% CI = 0.82–1.06). However, the second logistic regression model was adjusted for the Charlson comorbidity index as a covariate and demonstrated that both diabetics not on a DPP-4 inhibitors and nondiabetic controls were associated with a higher odds of AKI within 120 days compared with the diabetics on a DPP-4 (OR = 2.55; 95% CI = 2.24–2.9 and OR = 1.7; 95% CI = 1.49–1.94, respectively). This model also revealed that as the Charlson index increases, the odds of AKI also increase (OR = 1.38; 95% CI = 1.37–1.40). Additionally, a third logistic regression model was utilized that was adjusted for all study covariates and demonstrates that diabetic non-DPP-4 inhibitors and nondiabetics are associated with a higher odds of AKI within 120 days (OR = 2.78; 95% CI = 2.43–3.17 and OR = 1.32; 95% CI = 1.14–1.53, respectively). Model 3 also demonstrates that (i) higher values of the Charlson comorbidity index were associated with an increase in the odds of AKI (OR = 1.26; 95% CI = 1.24–1.27); (ii) higher baseline eGFR values are associated with a lower odds of AKI (OR = 0.96; 95% CI = 0.96–0.96); and (iii) compared with black patients, both white (OR = 0.76;

95% CI = 0.70–0.83) and other/unknown (OR = 0.60; 95% CI = 0.53–0.68) have a lower odds of AKI.

**Table 4** presents the results from the unadjusted odds of AKI within 120 days for comparisons between cohort groups using the Bonferroni multiplicity adjusted interval. When comparing DPP-4 cohort with diabetic non-DPP-4 controls, we found that DPP-4 use was associated with a lower odds AKI (unadjusted OR = 0.33; 95% CI = 0.27–0.4; adjusted OR = 0.34; 95% CI = 0.28–0.42). Comparing the DPP-4 cohort with the nondiabetic controls, we found that the odds of AKI in the DPP-4 cohort was not statistically significantly different in the unadjusted model (OR = 1.07, 95% CI = 0.88–1.3), yet was statistically significant with a lower odds of AKI within 120 days in the adjusted models (OR = 0.59, 95% CI = 0.48–0.71 and OR = 0.64, 95% CI = 0.52–0.79). Finally, in comparing the diabetic non-DPP-4 controls to the nondiabetic cohort, we found that the diabetic non-DPP-4 cohort had higher odds of AKI in both unadjusted (OR = 3.27; 95% CI = 2.96–3.61) and adjusted models (OR = 1.5; 95% CI = 1.34–1.67; OR = 1.86; 95% CI = 1.65–2.1).

**DISCUSSION**

This retrospective analysis of US veterans compared a national cohort of patients with T2DM and a DPP-4 inhibitor prescription with matched controls of nondiabetic patients. Patients were followed for an episode of AKI within 120 days. The goal of the study was to evaluate the association of DPP-4 inhibitor on the odds of AKI. This study found that of the cohort groups, the diabetic non-DPP-4 control cohort (22.95 (95% CI = 21.98–24.04) per 1,000 patients) had the highest rate of AKI compared with the DPP-4 cohort (7.63 (95% CI = 6.74–8.52) per 1,000 patients) and the nondiabetic control cohort (7.13 (95% CI = 6.80–7.45) per

**Table 2 Occurrence of AKI within 120 days**

Cohort	Total AKI	Per 1,000 people (95% CI <sup>a</sup> )
Diabetic non-DPP-4	1,784	22.95 (21.98–24.04)
Diabetic DPP-4	274	7.63 (6.74–8.52)
Nondiabetic	1,786	7.13 (6.80–7.45)

AKI, acute kidney injury; CI, confidence interval; DPP-4, dipeptidyl peptidase 4.

<sup>a</sup>Bootstrapped 95% CI.

**Table 3 Odds of AKI within 120 days**

Variable	Model 1	Model 2	Model 3
	OR (95% CI)	OR (95% CI)	OR (95% CI)
(Intercept)	0.01 (0.01–0.01)	0.00 (0.00–0.00)	0.38 (0.26–0.55)
Diabetic DPP-4 (baseline)			
Diabetic non-DPP-4	3.05 (2.69–3.47)	2.55 (2.24–2.9)	2.78 (2.43–3.17)
Nondiabetic	0.93 (0.82–1.06)	1.7 (1.49–1.94)	1.32 (1.14–1.53)
Charlson index		1.38 (1.37–1.40)	1.26 (1.24–1.27)
Baseline eGFR			0.96 (0.96–0.96)
Uncontrolled A1c			0.92 (0.84–1.02)
Sex (female baseline)			
Male			1.19 (0.95–1.48)
Race (black baseline)			
Other/unknown			0.6 (0.53–0.68)
White			0.76 (0.7–0.83)
Age			0.98 (0.98–0.99)
PPI use			0.71 (0.66–0.77)
NSAID use			0.58 (0.49–0.68)
Metformin use			0.41 (0.35–0.48)
Index year			0.92 (0.91–0.94)

AKI, acute kidney injury; CI, confidence interval; DPP-4, dipeptidyl peptidase 4; eGFR, estimated glomerular filtration rate; NSAID, nonsteroidal anti-inflammatory drugs; PPI, proton pump inhibitor.

**Table 4 Odds of AKI within 120 days: Comparisons between cohort groups**

Contrast	Model 1	Model 2	Model 3
	OR (95% CI) <sup>a</sup>	OR (95% CI) <sup>a</sup>	OR (95% CI) <sup>a</sup>
Diabetic DPP-4 vs. diabetic non-DPP-4	0.33 (0.27–0.4)	0.39 (0.32–0.48)	0.34 (0.28–0.42)
Diabetic DPP-4 vs. nondiabetic	1.07 (0.88–1.3)	0.59 (0.48–0.71)	0.64 (0.52–0.79)
Diabetic non-DPP-4 vs. nondiabetic	3.27 (2.96–3.61)	1.5 (1.34–1.67)	1.86 (1.65–2.1)

AKI, acute kidney injury; CI, confidence interval; DPP-4, dipeptidyl peptidase 4; OR, odds ratio.

<sup>a</sup>Bonferroni multiplicity adjusted interval.

1,000 patients). Diabetic non-DPP-4 controls had higher odds of AKI within 120 days compared with both the DPP-4 cohort and nondiabetic controls in all models. Additionally, the unadjusted logistic regression model demonstrates that DPP-4 inhibitors are not statistically different compared with nondiabetic controls, and a decreased odds of AKI compared with diabetic non-DPP-4 controls. In addition, in the unadjusted model, diabetic non-DPP-4 cohort had an over threefold increased odds of AKI within 120 days compared with the nondiabetic cohort. Furthermore, the fully adjusted model 3 demonstrated that the DPP-4 cohort had 66% and 36% lower odds of AKI within 120 days when compared with both control groups (diabetic non-DPP-4 and nondiabetics, respectively). Further, the diabetic non-DPP-4 cohort had an almost two fold increase in odds of AKI within 120 days compared with the nondiabetic cohort.

Systematic reviews and meta-analyses of the safety, efficacy, and tolerance of DPP-4 inhibitors on patients with T2DM calls for continued research and investigations on the effects of DPP-4 inhibitor use.<sup>17–20</sup> In a pooled analysis of randomized clinical studies of patients with T2DM on the DPP-4 inhibitor sitagliptin, the researchers concluded that the drug was tolerable up to 76 weeks and 2 years.<sup>21,22</sup> However, preclinical and clinical studies have focused on the association of AKI related to DPP-4 inhibitors and have found discordant results. Preclinical animal experimental studies have demonstrated a nephroprotective effect of DPP-4 inhibitors. Rats given an oral dose of DPP-4 sitagliptin (200 mg/kg/day) for 8 weeks had a renoprotective effect on their remnant kidneys.<sup>23</sup> Diabetic rats compared with nondiabetic rats treated with a DPP-4 inhibitor (vildagliptin) had lower body weight, plasma insulin levels, and serum creatinine levels. Additionally, the existence of interstitial expansion, glomerulosclerosis, and the thickening of the glomerular basement membrane attenuated renal injury in diabetic rats administered DPP-4 inhibitors.<sup>13</sup> DPP-4 inhibition resulted in a dose-dependent protective effect on kidney function in a rat model, as shown by reduced serum creatinine levels after renal ischemia–reperfusion injury. Additionally, the reduced serum creatinine levels were accompanied by morphological protection indicated by a decrease in tubular necrosis.<sup>15</sup>

Select clinical evaluations have demonstrated consistent results of the preclinical animal studies on the association

of AKI with DPP-4 inhibitors. The DPP-4 inhibitor linagliptin was not associated with increased kidney disease risk compared with placebo in individual patient-level data of pooled analysis of 13 phase II or III clinical trials.<sup>24</sup> DPP-4 inhibitors were associated with reduced risk of mild and severe forms of AKI among patients with incident diabetes in a nationally representative cohort in Taiwan.<sup>25</sup> Additionally, a retrospective cohort study of a large medical and pharmacy claims database demonstrated that there was not an association between the use of sitagliptin and AKI.<sup>26</sup> In contrast, there are reports of an increased association of AKI and DPP-4 inhibitors. The US Food and Drug Administration has reported episodes of acute worsening of renal function following exposure to inappropriate doses of sitagliptin in patients with renal insufficiency where there has been inappropriate dose adjustment.<sup>27</sup> Additionally, a nested case-control study conducted using Taiwan’s National Health Insurance Research Database demonstrated that those using DPP-4 inhibitors had an increased risk of AKI compared with nonusers.<sup>16</sup>

Our study results are consistent with the preclinical animal models and clinical studies that demonstrate a protective effect of DPP-4 inhibitor use on AKI. Our study provides value to the current literature regarding DPP-4 inhibitors and AKI because of previous discordant results. We utilized a different patient population to evaluate the external validity and generalizability of previous studies. Our large sample size utilizing matched controls provides additional evidence that DPP-4 inhibitors may have another role in addition to managing patients’ blood glucose levels. Although our study has a significant number of strengths, there are several limitations that are common to claims database studies of like methodology. Because this study was retrospective and observational, unmeasured confounders may have affected the results, despite matching of key covariates between cohorts. For example, there is a significant difference in the utilization of metformin between the DPP4 and non-DPP4 cohorts. Therefore, it is possible the patients on DPP4/metformin could have received more stringent care, and this could have influenced the outcome. The use of antihypertensive medications was not evaluated, and the outcome of blood pressure control through an antihypertensive may have had an influence on renal outcomes. Our study variable was the utilization of a DPP inhibitor and not the dose. However, the dose and/or existing renal function of the patient could have impacted the results. Specifically, the US Food and Drug Administration has reported episodes of acute worsening of renal function following exposure to inappropriate doses of sitagliptin in patients with renal insufficiency where there has been inappropriate dose adjustment.<sup>27</sup> Additionally, the study population was predominantly middle-to older-aged white men; therefore, our findings may not be generalizable to patients of different age groups or sexes. Additionally, only the association of DPP-4 inhibitors and AKI were studied; therefore, causality or mechanisms could not be investigated. However, our association data are consistent with previously conducted preclinical studies evaluating mechanisms. Preclinical studies suggest a glomerular and/or tubular mechanism of kidney injury; however, the exact role

and importance of the mechanism of kidney effects relating to the DPP-4 enzyme or DPP-4 inhibitors are not completely understood. Further research is needed to continue the evaluation of AKI in patients receiving DPP4 inhibitors to further our understanding of the association, mechanisms, and clinical application of AKI episodes.

**Acknowledgments.** This paper represents original research conducted using data from the Department of Veterans Affairs. This material is the result of work supported with resources and the use of facilities at the Dorn Research Institute, WJB Dorn Veterans Affairs Medical Center, Columbia, South Carolina. The contents do not represent the views of the US Department of Veterans Affairs or the US Government.

**Funding.** No funding was received for this work.

**Conflict of Interest.** The authors declared no competing interests for this work.

**Author Contributions.** S.S., J.M., T.C., and J.H. wrote the manuscript. S.S. and J.M. designed the research. S.S. and J.M. performed the research. S.S., J.M., T.C., and J.H. analyzed the data.

- Centers for Disease Control (CDC). National Center for Chronic Disease Prevention and Health Promotion – At A Glance 2016 Diabetes: Working to Reverse the US Epidemic. <<https://www.cdc.gov/chronicdisease/resources/publications/aag/pdf/2016/diabetes-aag.pdf>>. Published 2016. Accessed June 12, 2017.
- Jain, A., McDonald, H.I., Nitsch, D., Tomlinson, L. & Thomas, S.L. Risk factors for developing acute kidney injury in older people with diabetes and community-acquired pneumonia: a population-based UK cohort study. *BMC Nephrol.* **18**, 142 (2017).
- Branan, T.N. & Hawkins, W.A. Acute kidney injury. In McGraw-Hill's NAPLEX® Review Guide 3rd edn (ed. Sutton, S.S.) 324–333 (McGraw-Hill, New York, NY, 2015). <<https://accesspharmacy.mhmedical.com/content.aspx?bookx=ml:id=2512&sectionx=ml:id=201033811>>.
- Patschan, D. & Muller, G.A. Acute kidney injury in diabetes mellitus. *Int. J. Nephrol.* **2016**, 6232909 (2016).
- Dicker, D. DPP-4 inhibitors. *Diabetes Care* **34**(suppl. 2), S276–S278 (2011).
- Karagiannis, T., Paschos, P., Paletas, K., Matthews, D.R. & Tsapas, A. Dipeptidyl peptidase-4 inhibitors for treatment of type 2 diabetes mellitus in the clinical setting: systematic review and meta-analysis. *BMJ* **344**, e1369 (2012).
- Sakura, H. et al. Effect of sitagliptin on blood glucose control in patients with type 2 diabetes mellitus who are treatment naive or poorly responsive to existing antidiabetic drugs: the JAMP study. *BMC Endocr. Disord.* **16**, 70 (2016).
- Wang, W. et al. Efficacy and safety of linagliptin in Asian patients with type 2 diabetes mellitus inadequately controlled by metformin: a multinational 24-week, randomized clinical trial. *J. Diabetes* **8**, 229–237 (2016).
- Sun, A.-L. et al. Dipeptidyl peptidase-IV is a potential molecular biomarker in diabetic kidney disease. *Diabetes Vasc. Dis. Res.* **9**, 301–308 (2012).
- Mima, A. et al. Protective effects of GLP-1 on glomerular endothelium and its inhibition by PKC $\beta$  activation in diabetes. *Diabetes* **61**, 2967–2979 (2012).
- Górriz, J.L., Nieto, J., Navarro-González, J.F., Molina, P., Martínez-Castelao, A. & Pallardó, L.M. Nephroprotection by hypoglycemic agents: do we have supporting data? *J. Clin. Med.* **4**, 1866–1889 (2015).
- Mega, C. et al. Diabetic nephropathy amelioration by a low-dose sitagliptin in an animal model of type 2 diabetes (Zucker diabetic fatty rat). *Exp. Diabetes Res.* **2011**, 162092 (2011).
- Liu, W.J. et al. Dipeptidyl peptidase IV inhibitor attenuates kidney injury in streptozotocin-induced diabetic rats. *J. Pharmacol. Exp. Ther.* **340**, 248–255 (2012).
- Alter, M.L. et al. DPP-4 inhibition on top of angiotensin receptor blockade offers a new therapeutic approach for diabetic nephropathy. *Kidney Blood Press. Res.* **36**, 119–130 (2012).
- Glorie, L.L.F. et al. DPP4 inhibition improves functional outcome after renal ischemia-reperfusion injury. *Am. J. Physiol. Renal Physiol.* **303**, F681–F688 (2012).
- Shih, C.-J., Lee, Y.-J., Lo, Y.-H., Kuo, S.-C., Ou, S.-M. & Chen, Y.-T. Association between use of dipeptidyl peptidase-4 inhibitors and the risk of acute kidney injury: a nested case-control study. *Mayo Clin. Proc.* **91**, 867–872 (2016).
- Cox, M.E., Rowell, J., Corsino, L., Green, J.B. Dipeptidyl peptidase-4 inhibitors in the management of type 2 diabetes: safety, tolerability, and efficacy. *Drug Healthc. Patient Saf.* **2**, 7–19 (2010).
- Liu, J. et al. Incretin based treatments and mortality in patients with type 2 diabetes: systematic review and meta-analysis. *BMJ* **357**, j2499 (2017).
- Unger, J. Incretins: clinical perspectives, relevance, and applications for the primary care physician in the treatment of patients with type 2 diabetes mellitus. *Mayo Clin. Proc.* **85**(suppl. 12), S38–S49 (2010).
- Scheen, A.J. DPP-4 inhibitors in the management of type 2 diabetes: a critical review of head-to-head trials. *Diabetes Metab.* **38**, 89–101 (2012).
- Williams-Herman, D. et al. Safety and tolerability of sitagliptin in clinical studies: a pooled analysis of data from 10,246 patients with type 2 diabetes. *BMC Endocr. Disord.* **10**, 7 (2010).
- Frederich, R., McNeill, R., Berglind, N., Fleming, D. & Chen, R. The efficacy and safety of the dipeptidyl peptidase-4 inhibitor saxagliptin in treatment-naïve patients with type 2 diabetes mellitus: a randomized controlled trial. *Diabetol. Metab. Syndr.* **4**, 36 (2012).
- Joo, K.W. et al. Dipeptidyl peptidase IV inhibitor attenuates kidney injury in rat remnant kidney. *BMC Nephrol.* **14**, 98 (2013).
- Cooper, M.E. et al. Kidney disease end points in a pooled analysis of individual patient-level data from a large clinical trials program of the dipeptidyl peptidase 4 inhibitor linagliptin in type 2 diabetes. *Am. J. Kidney Dis.* **66**, 441–449 (2015).
- Chao, C.-T., Wang, J., Wu, H.-Y., Chien, K.-L. & Hung, K.-Y. Dipeptidyl peptidase 4 inhibitor use is associated with a lower risk of incident acute kidney injury in patients with diabetes. *Oncotarget* **8**, 53028–53040 (2017).
- Pendergrass, M., Fenton, C., Haffner, S.M. & Chen, W. Exenatide and sitagliptin are not associated with increased risk of acute renal failure: a retrospective claims analysis. *Diabetes Obes. Metab.* **14**, 596–600 (2012).
- Hoher, B., Reichetzedler, C. & Alter, M.L. Renal and cardiac effects of DPP-4 inhibitors - from preclinical development to clinical research. *Kidney Blood Press. Res.* **36**, 65–84 (2012).

**Published 2019. This article is a U.S. Government work and is in the public domain in the USA. *Clinical and Translational Science* published by Wiley Periodicals, Inc. on behalf of the American Society for Clinical Pharmacology and Therapeutics. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.**