

RESEARCH ARTICLE

Nasopharyngeal Microbiome Diversity Changes over Time in Children with Asthma

Marcos Pérez-Losada^{1,2,3*}, Lamia Alamri², Keith A. Crandall¹, Robert J. Freishtat²

1 Computational Biology Institute, George Washington University, Ashburn, Virginia, United States of America, **2** Division of Emergency Medicine, Children's National Medical Center, Washington DC, United States of America, **3** CIBIO-InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Campus Agrário de Vairão, Vairão, Portugal

* mlosada@gwu.edu



Abstract

Background

The nasopharynx is a reservoir for pathogens associated with respiratory illnesses such as asthma. Next-generation sequencing (NGS) has been used to characterize the nasopharyngeal microbiome of infants and adults during health and disease; less is known, however, about the composition and temporal dynamics (i.e., longitudinal variation) of microbiotas from children and adolescents. Here we use NGS technology to characterize the nasopharyngeal microbiomes of asthmatic children and adolescents (6 to 18 years) and determine their stability over time.

Methods

Two nasopharyngeal washes collected 5.5 to 6.5 months apart were taken from 40 children and adolescents with asthma living in the Washington D.C. area. Sequence data from the 16S-V4 rRNA gene region (~250 bp) were collected from the samples using the MiSeq platform. Raw data were processed in mothur (SILVA123 reference database) and Operational Taxonomic Units (OTU)-based alpha- and beta-diversity metrics were estimated. Relatedness among samples was assessed using PCoA ordination and Procrustes analyses. Differences in microbial diversity and taxon mean relative proportions were assessed using linear mixed effects models. Core microbiome analyses were also performed to identify stable and consistent microbes of the nasopharynx.

Results and Discussion

A total of 2,096,584 clean 16S sequences corresponding to an average of 167 OTUs per sample were generated. Representatives of *Moraxella**, *Staphylococcus**, *Dolosigranulum*, *Corynebacterium*, *Prevotella*, *Streptococcus**, *Haemophilus**, *Fusobacterium** and a Neisseriaceae genus accounted for 86% of the total reads. These nine genera have been previously found in the nasopharynxes of both infants and adults, but in different proportions. OTUs from the five genera highlighted (*) above defined the nasopharyngeal core microbiome at the 95% level. No significant differences in alpha- and beta-diversity were

OPEN ACCESS

Citation: Pérez-Losada M, Alamri L, Crandall KA, Freishtat RJ (2017) Nasopharyngeal Microbiome Diversity Changes over Time in Children with Asthma. PLoS ONE 12(1): e0170543. doi:10.1371/journal.pone.0170543

Editor: Paul J Planet, Columbia University, UNITED STATES

Received: January 29, 2016

Accepted: January 6, 2017

Published: January 20, 2017

Copyright: © 2017 Pérez-Losada et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Sequence data have been uploaded to the GenBank under SRA accession number SRP069020.

Funding: MP-L was funded in part by a K12 Career Development Program K12HL119994 award and by Award Numbers UL1TR001876 and KL2TR001877 from the NIH National Center for Advancing Translational Sciences. The AsthMaP2 project was supported by Award Numbers R01MD007075 to RJF and UL1TR000075 from the NIH National Center for Advancing Translational Sciences (RJF, KAC), in addition to funding to RJF

from the Clark Charitable Foundation. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the National Center for Advancing Translational Sciences or the National Institutes of Health.

Competing Interests: The authors have declared that no competing interests exist.

observed between seasons, but bacterial mean relative proportions of *Haemophilus*, *Moraxella*, *Staphylococcus* and *Corynebacterium* varied significantly between summer-fall and age groups (inter-patient variation). Additionally, OTUs varied significantly within patients between time points in 35 of the 40 patients analyzed. Future cross-sectional studies should be mindful of the temporal dynamics of the nasopharyngeal microbiota.

Introduction

The application of next-generation sequencing (NGS) technology to microbial communities residing in the respiratory airways (i.e., airway microbiome) has shown that bacteria may play a significant role in the onset, development, and severity of respiratory diseases such as asthma [1–4]. Both metataxonomics and metagenomics (see [5] for distinction) have demonstrated that airway microbiome composition and structure vary between healthy and diseased individuals and revealed significant associations between representatives of several opportunistic pathogens (*Moraxella*, *Streptococcus*, *Haemophilus*, *Neisseria* and *Staphylococcus*), clinical conditions, and stages of asthma [3, 6–12]. Moreover, microbiome research has shown that all of these pathogens are normal and transient residents of the nasopharynx [10]. As such, they are mainly asymptomatic, but through the nasopharynx they can directly spread to other sections of the respiratory tract and potentially cause asthma, otitis media or pneumonia; or invade the bloodstream to cause sepsis and meningitis [8, 13–15].

Given the general importance of the nasopharynx as a reservoir for microbes associated with respiratory infections [8, 16, 17], several NGS studies have investigated the microbial composition of the nasopharynx during health and disease in both infants and adults, and in relation to clinical factors. Most of these microbiome studies, however, have been designed in a cross-sectional setting (i.e., one time point) [14, 18–24]. Less is known about the temporal dynamics of the nasopharyngeal microbiota (i.e., microbial succession) within patients and, therefore, the influence of changing environmental factors (e.g., seasonality) on the structuring of microbial communities over time. Moreover, temporal variation complicates the prediction of microbial profiles associated with disease progression, the assessment of intra- and inter-patient variation, and the study of microbe-host interactions. Gut microbiome research is increasingly showing that longitudinal (temporal) changes in bacterial community composition and function are associated with disease [25–27]. Similarly, longitudinal studies of the nasopharyngeal microbiome in infants have revealed that certain microbial groups change over the first two years of life in both healthy and asthmatic children [8, 15] and that microbiota profiles vary strongly with the seasons [8, 10]. Additionally, these studies have also shown a high inter-patient microbial diversity, so no nasopharyngeal core microbiome could be defined at the OTU level.

Here, we use NGS technology to explore the nasopharyngeal microbiome. We couple targeted 16S rRNA sequencing (metataxonomics) with a sophisticated analytical pipeline to characterize the nasal microbiome diversity of asthmatic children and adolescents (ages 6 to 18 years) and determine its stability (i.e., microbial succession) over time (5.5 to 6.5 months) and across seasons. Our results complement previous microbiome studies of the nasopharynx focused on younger children (<2 years of age) and adults.

Materials and Methods

Ethics

All participants in this study were part of the AsthMaP2 (Asthma Severity Modifying Polymorphisms) Study. AsthMaP2 is an ongoing study of urban children and adolescents designed to find associations among airway microbes, environmental exposures, allergic sensitivities, genetics, and asthma. AsthMaP2 and the study presented here were approved by the Children's National Medical Center Institutional Review Board (Children's National IRB), which requires that consent is obtained and documented prior to conducting study procedures and collection of samples for research. Written consent was obtained from all independent participants or their legal guardians using the Children's National IRB approved informed consent documents (IRB No PRO00002517).

Samples and molecular analyses

Two nasal washes (W1 and W2) were collected from 40 children and adolescents (ages 6 to 18 years; mean = 11 years) enrolled in the AsthMaP2 study in two consecutive visits (5.5 to 6.5 months apart) at Children's National Medical Center (Washington, DC) [28]. Children were recruited from the metropolitan Washington, DC, area; they had been physician-diagnosed with asthma for at least one year prior to recruitment and have been followed up for another year afterwards. Individuals who reported a medical history of chronic or complex cardiorespiratory disease were ineligible.

Washes were procured by instilling 5 ml of isotonic sterile saline buffer into each nare, holding it for 10 seconds and then blowing into a specimen collection container. Total DNA was extracted using the QIAGEN QIAamp DNA Kit (Catalog # 51304). Before adding the ATL buffer, samples were pre-incubated in 100 μ l of lysozyme-TE buffer pH = 8.0 for 30 minutes at 37°C. All extractions yielding >50 ng of total DNA (as indicated by NanoDrop 2000 UV-Vis Spectrophotometer measuring) were further processed. DNA extractions were prepared for sequencing using the Schloss' MiSeq_WetLab_SOP protocol (09.2015) in Kozich et al. [29]. Each DNA sample was amplified for the V4 region (~250 bp) of the 16S rRNA gene and libraries were sequenced in a single run of the Illumina MiSeq sequencing platform at University of Michigan Medical School. Sequence data have been deposited in GenBank under SRA accession number SRP069020.

Analyses

Raw FASTQ files were processed in mothur v1.35.1 [30]. Default settings were used to minimize sequencing errors as described in Schloss et al. [31]. Clean sequences were aligned to the SILVA123-based bacterial reference alignment at <http://www.mothur.org>. Chimeras were removed using uchime [32] and non-chimeric sequences were classified using the naïve Bayesian classifier of Wang et al. [33]. Sequences were clustered into Operational Taxonomic Units (OTUs) at the 0.03 threshold (species level). OTU sequence representatives and taxonomy were imported (BIOM format) into QIIME [34] for subsequent analyses. The mothur OTU table was filtered to a minimum of 2 observations (sequences) per OTU. Samples were subsampled (rarefaction analysis) to the smallest sample size (2,288 sequences) to remove the effect of sample size bias on community composition.

Trees for phylogenetic diversity calculations were constructed using FastTree [35]. Taxonomic alpha-diversity was estimated as the number of observed OTUs, but also by the Good's coverage, Chao1, and Shannon indices. Phylogenetic alpha-diversity was calculated by the Faith's phylogenetic diversity index [36]. Phylogenetic (unweighted and weighted unfrac)

beta-diversity metrics were calculated between pairs of samples. Dissimilarity between samples was estimated using principal coordinates analysis (PCoA) and both unifrac distances. Procrustes analysis comparing PCoA plots (weighted unifrac distances) of W1 and W2 groups [37] was performed with 10,000 Monte Carlo iterations. Linear mixed effects models (LME) analysis was applied to both alpha-diversity indices and taxa mean relative proportions (response), while accounting for non-independence of intra-patient samples (random effect) and seasons, age (years), gender, body mass index (BMI), National Asthma Education and Prevention Program (NAEPP) severity level [38], and ethnic background (predictors) (S1 Table). Phylogenetic distances were also compared between seasons using the non-parametric adonis test from the vegan R's library [39] while accounting for random effects and all predictors above. Significance was determined through permutation testing after 10,000 permutations. Finally, differences in genus mean relative proportions between visits (W1 versus W2) were assessed for each patient using the Fisher's exact test. The nasopharyngeal core microbiome (OTUs that are present in a certain percentage of samples) in our 80 samples was identified at different levels of stringency ranging from 50% to 100% of the samples. Bonferroni or Benjamini-Hochberg FDR multiple test correction methods were applied. All analyses were performed in mothur, QIIME, STAMP [40], and RStudio [41].

Results and Discussion

The nasopharyngeal microbiome of asthmatic children

A total of 80 nasal microbiomes corresponding to 40 asthmatic children (two samples per patient 5.5 to 6.5 months apart) were analyzed via MiSeq sequencing of 16S rRNA V4 amplicons. A total of 2,096,584 sequences ranging from 2,288 to 60,806 sequences per sample (mean = 26,207.3; median = 22,381.5) were obtained, after quality control analyses and OTU filtering. From these data, we identified 34–394 OTUs (mean = 167) per sample belonging to 397 classified bacterial genera and 23 classified phyla (S2 and S3 Tables). About 86% of the sequences corresponded to the following nine genera: *Moraxella* (35.3%), *Staphylococcus* (13.9%), *Dolosigranulum* (9.3%), *Corynebacterium* (8.8%), *Prevotella* (6.0%), *Streptococcus* (4.9%), *Haemophilus* (3.5%), *Fusobacterium* (3.0%) and Neisseriaceae genus (1.3%) (see Fig 1 and S3 Table).

Our study does not include samples from healthy children, hence we cannot assess to what extent the microbial profiles we observe here are unique to asthmatic children. Moreover, we are not aware of any study that has described the nasopharyngeal microbiome of healthy children (2 to 18 years old). For comparative purposes, we have then compiled results from several published studies describing the nasopharyngeal microbiomes of non-asthmatic infants (defined as children between the ages of 1–18 months) and adults (Table 1). Some genera in our study show different mean relative proportions than those seen both in infants (*Corynebacterium*, *Dolosigranulum*, *Flavobacterium*, *Haemophilus*, *Prevotella* and *Streptococcus*) and adults (*Corynebacterium*, *Moraxella*, *Prevotella* and *Streptococcus*), while others seem to show similar mean relative proportions across all age groups (*Fusobacterium*, *Neisseria* and *Staphylococcus*). This comparison, therefore, suggests that the nasopharyngeal microbiome profiles observed here are different from those seen in healthy infants and adults.

The nasopharynx is considered a reservoir for commensal and potentially pathogenic microbes inhabiting the respiratory airways and associated with acute respiratory infections, including asthma [4, 8–10, 16, 42]. All dominant genera reported here for the nasopharynx [several of them including pathogenic species (e.g., *Moraxella*, *Staphylococcus* and *Streptococcus*)] have been detected in other microbiome studies of lower and upper (excluding the nasopharynx) respiratory airways of asthmatics [1, 16, 43, 44]. Thus, our results confirm in




Fig 1. Taxonomic profiles of 80 nasopharyngeal microbiomes from 40 asthmatic children. Only OTUs with a minimum total observation count of 0.1% are shown. W1 = nasal microbiome from wash 1; W2 = nasal microbiome from wash 2 collected 5.5 to 6.5 months later. Sample pairs are alternatively colored in red and black to facilitate their visualization.

doi:10.1371/journal.pone.0170543.g001

Table 1. Summary of studies of the nasopharyngeal microbiome in non-asthmatic patients. Number of samples (Ns), age group by month (m) and year (yr), sequenced 16S region, patient condition and genus mean relative proportions (with SD estimates for our study) are indicated. Studies (included ours) are ordered by patient age group.

Study	Teo et al. (2015)	Bogaert et al. (2011)	This study	Hilty et al. (2010)
Ns	487	96	80	40
Age group	Infants (1–12 m)	Infants (18 m)	Children (6–17 yr)	Adults (≥ 18 yr)
16S	V4	V5–V6	V4	V3–V6
Patient condition	Healthy	Healthy	Asthmatic	Healthy
<i>Corynebacterium</i>	10–22	2	8.8 (13.1)	32.8
<i>Dolosigranulum</i>	14–26	5	9.3 (13.6)	9.9
<i>Flavobacterium</i>	NR	10	<1	<1
<i>Fusobacterium</i>	NR	1	3 (6.4)	2.6
<i>Haemophilus</i>	NR	20	3.5 (12.6)	4.5
<i>Moraxella</i>	9–41	40	35.5 (37.0)	4.8
<i>Neisseria</i>	NR	2	<1	<1
<i>Prevotella</i>	NR	<1	6 (14.4)	3.5
<i>Staphylococcus</i>	11–41	<1	13.9 (18.3)	9.1
<i>Streptococcus</i>	NR	12	4.9 (12.8)	9.1

NR = Proportion not reported

doi:10.1371/journal.pone.0170543.t001

children the role of the nasopharynx as a source of potential pathogens for other sections of the respiratory tract [4, 16]. Previous studies of the nasopharyngeal microbiome in both infants [8] and adults [16] with respiratory infections showed that their microbiomes were comprised of the same main genera reported here for asthmatic children and adolescents (ages of 6 to 18 years), although their mean relative proportions varied across studies. Nasopharyngeal microbiomes of infants (1 to 14 months) with acute respiratory infections seem to be dominated by *Haemophilus*, *Streptococcus*, and *Moraxella* [8]; while nasopharyngeal microbiomes of adults with asthma seem to be dominated with *Staphylococcus*, *Corynebacterium*, and *Neisseria* [16]. Given that in our study *Moraxella* and *Staphylococcus* are the dominant taxa, this seems to suggest that microbial succession towards more adult-like configurations in the nasopharynx of asthmatics does not end at early childhood (~2 years) [8, 15] and may take longer than in other body compartments such as the gut [45].

The asthma core microbiome

The differentiation of the core microbiome of a host meta-organism or host microhabitat is a means to differentiate the stable and consistent members and associations from the whole community [46, 47]. The identification of core members of the meta-organism subsequently allows for the differentiation of core pathways and metabolic functions that are provided by the host—microbe interaction [48]. A core microbiome composed of a variable number of OTUs was observed in the studied nasopharyngeal samples. The least stringent definition of the core (presence in at least 50% of the samples) identified 43 OTUs, while the most stringent definition (presence in all sampled children) included only one OTU of the genus *Staphylococcus*. At the 95% level, the nasopharyngeal core microbiome of asthmatic children was represented by five OTUs of the genera *Moraxella*, *Staphylococcus*, *Streptococcus*, *Haemophilus*, and *Fusobacterium*. These five OTUs may represent fingerprints or biological markers (pulmotypes; [17]) of the nasopharyngeal (and potentially the airway) microbiome in asthmatic children. Most studies of the nasopharyngeal microbiome in infants with and without respiratory infections [8, 10, 15, 49] were not able to define a core microbiome at the OTU level due to the high inter-individual variability of their early (and potentially more dynamic) microbiomes. Using a less strict definition of core microbiome (i.e., OTUs present in more than 50% of all samples and representing >0.1% of the sequences), Bogaert et al. [10] described a core microbiome in children 18 months of age composed of *Moraxella*, *Haemophilus*, *Enhydrobacter*, *Streptococcus*, *Dolosigranulum*, and *Corynebacterium*. All these genera were observed in our 50% core microbiome except *Enhydrobacter*, and three genera (*Moraxella*, *Haemophilus*, and *Streptococcus*) were observed in our 95% core microbiome. Therefore, if the consistency and specificity (i.e., dominant in asthmatics and absent in healthy controls) of these taxa across all “ages of asthma” are confirmed, they could potential be used to phenotype this complex and heterogeneous disease (i.e., pulmotypes).

Temporal diversity in the nasopharyngeal microbiome

Procrustes analysis comparing PCoA plots of W1 (nasal wash 1) and W2 (nasal wash 2) microbiomes collected 5.5 to 6.5 months apart showed a great dissimilarity between ($M^2 = 0.663$) both datasets (Fig 2). Microbial profiles (Fig 1) and pairwise comparisons of the most abundant genera between W1 and W2 samples (S1 Fig) also varied greatly across patients, with some patients showing increases in bacterial mean relative proportions, some showing the opposite trend, and others showing no visible change. Similarly, some bacterial taxa seemed to fluctuate more than others (e.g., *Moraxella* versus *Streptococcus*) between W1 and W2 samples (S1 Fig). Differences in OTU mean relative proportions between W1 and W2




Fig 2. Procrustes analysis of 40 nasopharyngeal sample pairs collected 5.5 to 6.5 months apart. Sample pairs (red dots) collected from the same patient are connected by a yellow bar. The length of the bar is proportional to the dissimilarity between sample pairs.

doi:10.1371/journal.pone.0170543.g002

microbiomes were significant in 35 (88%) of the 40 sample pairs analyzed. The number of significant ($P < 0.05$) OTUs detected varied from one to 19, as indicated by the Fisher's exact test (effective size $\geq 1\%$ for the difference in mean relative proportions). All these results combined indicate that members of the nasopharyngeal bacteriome of asthmatic children change over time (5.5 to 6.5 months) on an individual basis. Our study complements previous longitudinal studies revealing differences in microbial composition between infants 2, 6




Fig 3. 3D Principal Coordinates Analyses of weighted (A) and unweighted (B) unifracs distances between 80 nasopharyngeal microbiomes colored by season.

doi:10.1371/journal.pone.0170543.g003

Table 2. Linear mixed effects (LME) models analysis. Significance (P-values) of the LME analysis of mean relative proportions of most abundant genera (>1% of reads) and clinical and demographic variables. All four seasons were compared, but only the lowest P-values are reported.

	Season	Age	Gender	BMI	Severity Classification	Ethnic Background
<i>Moraxella</i>	0.20	0.002	0.932	0.139	0.082	0.415
<i>Staphylococcus</i>	0.11	0.045	0.897	0.578	0.482	0.388
<i>Dolosigranulum</i>	0.12	0.610	0.680	0.520	0.270	0.130
<i>Corynebacterium</i>	0.12	0.003	0.106	0.845	0.335	0.052
<i>Prevotella</i>	0.54	0.629	0.769	0.220	0.723	0.055
<i>Streptococcus</i>	0.16	0.065	0.456	0.762	0.691	0.387
<i>Haemophilus</i>	0.014	0.510	0.355	0.594	0.510	0.612
<i>Fusobacterium</i>	0.36	0.250	0.220	0.057	0.499	0.846
Neisseriaceae genus	0.35	0.780	0.890	0.330	0.710	0.470

Haemophilus: 0.014* (only summer-fall was significant)

doi:10.1371/journal.pone.0170543.t002

and 12 months old at risk of developing respiratory infections and asthma [8]. Future research assessing microbial succession and host-microbe interactions during asthma should account for both intra- and inter-patient microbiome variability in the studied cohorts and temporal dynamics at even shorter time spans than those studied here.

To further explore inter-patient diversity in our cohort, we investigated if sampling seasonality impacted alpha- and beta-diversity and microbe (genera) mean relative proportions while accounting for non-independence of samples, age (years), gender, BMI, NAEPP severity level, and ethnic background (S1 Table). We did not detect significant differences ($P > 0.05$) in alpha diversity (observed OTUs, Good's coverage, Chao1, Shannon and Faith—S2 Table) between seasonal groups or any of the confounders in our linear mixed effects models (LME) analyses (data not shown). Additionally, ordination analysis (PCoA—Fig 3) did not reveal clear dissimilarities between nasopharyngeal microbiomes grouped by season for both unifac distances. Similarly, no significant differences ($P > 0.05$) in beta-diversity (unifrac distances) were detected by the adonis tests between seasons while accounting for all other confounders (data not shown). This contrasts with previous nasopharyngeal microbiome studies showing large differences in the microbiota profiles of healthy infants during seasons [10]. No study so far has addressed if those differences are also seen in children and adults.

At the taxon-level, our LME analyses of mean relative proportions of the most abundant genera (Table 2) found no significant differences between seasons for all taxa but *Haemophilus*, which showed a significantly ($P = 0.014$) higher mean relative proportion in summer (9.3% average) compared to fall (0.3% average) (Table 2). Previous nasopharyngeal microbiome studies have also revealed significant seasonal effects on the proportions of *Haemophilus* and *Moraxella* in infants at risk of developing asthma [8]. Our LME analyses also showed that patient age (6 to 18 years) is significantly associated with variation in the mean relative proportions of *Moraxella*, *Staphylococcus*, and *Corynebacterium*. This agrees with our previous results suggesting that the composition of the nasopharyngeal microbiome changes towards adulthood [8, 15].

Our study (and others) shows that nasopharyngeal microbial communities are not stable; rather they experience structural rearrangements (deterministic or stochastic) over time. Given the exposure of the nasopharynx, such longitudinal changes could be related to environmental factors (e.g., seasonality, diet, pets), although clinical factors (e.g., age, body mass index, neutrophil counts, medication), other microbiotas (i.e., gastrointestinal, oral), anatomical diversity (i.e., nasal microenvironments), and technical biases (e.g., DNA extraction

protocol) could also play a role [1–3, 8–10, 42, 50]. Whatever the cause, future cross-sectional studies of the nasopharyngeal microbiome need to be mindful of the short-time dynamics of the microbiotas occupying this human anatomical cavity.

Supporting Information

S1 Table. Clinical and demographic characteristics of the cohort of 40 patients analyzed in this study.

(XLSX)

S2 Table. Alpha-diversity indices (Good's coverage, observed OTUs, Chao1, Shannon and PD) for 80 nasopharyngeal microbiomes from 40 asthmatic children.

(XLSX)

S3 Table. Absolute counts and taxonomic classification of OTUs in 80 nasopharyngeal microbiomes from 40 asthmatic children.

(XLSX)

S1 Fig. Proportions of sequences of the most abundant bacterial genera in 80 nasopharyngeal microbiomes from 40 asthmatic children.

(PDF)

Acknowledgments

We thank the GWU Colonial One High Performance Computing Cluster for computational time.

Author Contributions

Conceptualization: MP-L KAC RJF.

Data curation: MP-L LA.

Formal analysis: MP-L.

Funding acquisition: MP-L KAC RJF.

Investigation: MP-L KAC RJF.

Methodology: MP-L LA.

Project administration: MP-L KAC RJF.

Resources: MP-L KAC RJF.

Software: MP-L.

Supervision: MP-L KAC RJF.

Validation: MP-L KAC RJF.

Visualization: MP-L.

Writing – original draft: MP-L KAC RJF.

Writing – review & editing: MP-L KAC RJF.

References

1. Huang YJ, Boushey HA. The microbiome and asthma. *Annals of the American Thoracic Society*. 2014; 11 Suppl 1:S48–51.
2. Huang YJ, Boushey HA. The microbiome in asthma. *The Journal of allergy and clinical immunology*. 2015; 135(1):25–30. doi: [10.1016/j.jaci.2014.11.011](https://doi.org/10.1016/j.jaci.2014.11.011) PMID: [25567040](https://pubmed.ncbi.nlm.nih.gov/25567040/)
3. Brar T, Nagaraj S, Mohapatra S. Microbes and asthma: the missing cellular and molecular links. *Current opinion in pulmonary medicine*. 2012; 18(1):14–22. doi: [10.1097/MCP.0b013e32834dccc0](https://doi.org/10.1097/MCP.0b013e32834dccc0) PMID: [22113000](https://pubmed.ncbi.nlm.nih.gov/22113000/)
4. Dickson RP, Huffnagle GB. The Lung Microbiome: New Principles for Respiratory Bacteriology in Health and Disease. *PLoS pathogens*. 2015; 11(7):e1004923. doi: [10.1371/journal.ppat.1004923](https://doi.org/10.1371/journal.ppat.1004923) PMID: [26158874](https://pubmed.ncbi.nlm.nih.gov/26158874/)
5. Marchesi JR, Ravel J. The vocabulary of microbiome research: a proposal. *Microbiome*. 2015; 3:31. doi: [10.1186/s40168-015-0094-5](https://doi.org/10.1186/s40168-015-0094-5) PMID: [26229597](https://pubmed.ncbi.nlm.nih.gov/26229597/)
6. Bisgaard H, Hermansen MN, Bonnelykke K, Stokholm J, Baty F, Skytt NL, et al. Association of bacteria and viruses with wheezy episodes in young children: prospective birth cohort study. *Bmj*. 2010; 341: c4978. doi: [10.1136/bmj.c4978](https://doi.org/10.1136/bmj.c4978) PMID: [20921080](https://pubmed.ncbi.nlm.nih.gov/20921080/)
7. Castro-Nallar E, Shen Y, Freishtat RJ, Pérez-Losada M, Manimaran S, Liu G, et al. Integrating metagenomics and host gene expression to characterize asthma-associated microbial communities. *BMC Medical Genomics*. 2015; 8:50.
8. Teo SM, Mok D, Pham K, Kusel M, Serralha M, Troy N, et al. The infant nasopharyngeal microbiome impacts severity of lower respiratory infection and risk of asthma development. *Cell Host & Microbe*. 2015; 17:704–15.
9. Perez-Losada M, Castro-Nallar E, Bendall ML, Freishtat RJ, Crandall KA. Dual Transcriptomic Profiling of Host and Microbiota during Health and Disease in Pediatric Asthma. *PloS one*. 2015; 10(6): e0131819. doi: [10.1371/journal.pone.0131819](https://doi.org/10.1371/journal.pone.0131819) PMID: [26125632](https://pubmed.ncbi.nlm.nih.gov/26125632/)
10. Bogaert D, Keijsers B, Huse S, Rossen J, Veenhoven R, van Gils E, et al. Variability and diversity of nasopharyngeal microbiota in children: a metagenomic analysis. *PloS one*. 2011; 6(2):e17035. doi: [10.1371/journal.pone.0017035](https://doi.org/10.1371/journal.pone.0017035) PMID: [21386965](https://pubmed.ncbi.nlm.nih.gov/21386965/)
11. Jounio U, Juvonen R, Bloigu A, Silvennoinen-Kassinen S, Kaijalainen T, Kauma H, et al. Pneumococcal carriage is more common in asthmatic than in non-asthmatic young men. *The clinical respiratory journal*. 2010; 4(4):222–9. doi: [10.1111/j.1752-699X.2009.00179.x](https://doi.org/10.1111/j.1752-699X.2009.00179.x) PMID: [20887345](https://pubmed.ncbi.nlm.nih.gov/20887345/)
12. Bisgaard H, Hermansen MN, Buchvald F, Loland L, Halkjaer LB, Bonnelykke K, et al. Childhood asthma after bacterial colonization of the airway in neonates. *The New England journal of medicine*. 2007; 357(15):1487–95. doi: [10.1056/NEJMoa052632](https://doi.org/10.1056/NEJMoa052632) PMID: [17928596](https://pubmed.ncbi.nlm.nih.gov/17928596/)
13. Bogaert D, De Groot R, Hermans PW. Streptococcus pneumoniae colonisation: the key to pneumococcal disease. *The Lancet Infectious diseases*. 2004; 4(3):144–54. doi: [10.1016/S1473-3099\(04\)00938-7](https://doi.org/10.1016/S1473-3099(04)00938-7) PMID: [14998500](https://pubmed.ncbi.nlm.nih.gov/14998500/)
14. Garcia-Rodriguez JA, Fresnadillo Martinez MJ. Dynamics of nasopharyngeal colonization by potential respiratory pathogens. *The Journal of antimicrobial chemotherapy*. 2002; 50 Suppl S2:59–73.
15. Biesbroek G, Tsvitshivadze E, Sanders EA, Montijn R, Veenhoven RH, Keijsers BJ, et al. Early respiratory microbiota composition determines bacterial succession patterns and respiratory health in children. *American journal of respiratory and critical care medicine*. 2014; 190(11):1283–92. doi: [10.1164/rccm.201407-1240OC](https://doi.org/10.1164/rccm.201407-1240OC) PMID: [25329446](https://pubmed.ncbi.nlm.nih.gov/25329446/)
16. Hilty M, Burke C, Pedro H, Cardenas P, Bush A, Bossley C, et al. Disordered microbial communities in asthmatic airways. *PloS one*. 2010; 5(1):e8578. doi: [10.1371/journal.pone.0008578](https://doi.org/10.1371/journal.pone.0008578) PMID: [20052417](https://pubmed.ncbi.nlm.nih.gov/20052417/)
17. Dickson RP, Erb-Downward JR, Huffnagle GB. The role of the bacterial microbiome in lung disease. *Expert review of respiratory medicine*. 2013; 7(3):245–57. doi: [10.1586/ers.13.24](https://doi.org/10.1586/ers.13.24) PMID: [23734647](https://pubmed.ncbi.nlm.nih.gov/23734647/)
18. Mackenzie GA, Leach AJ, Carapetis JR, Fisher J, Morris PS. Epidemiology of nasopharyngeal carriage of respiratory bacterial pathogens in children and adults: cross-sectional surveys in a population with high rates of pneumococcal disease. *BMC infectious diseases*. 2010; 10:304. doi: [10.1186/1471-2334-10-304](https://doi.org/10.1186/1471-2334-10-304) PMID: [20969800](https://pubmed.ncbi.nlm.nih.gov/20969800/)
19. Feazel LM, Santorico SA, Robertson CE, Bashraheil M, Scott JA, Frank DN, et al. Effects of Vaccination with 10-Valent Pneumococcal Non-Typeable Haemophilus influenzae Protein D Conjugate Vaccine (PHiD-CV) on the Nasopharyngeal Microbiome of Kenyan Toddlers. *PloS one*. 2015; 10(6):e0128064. doi: [10.1371/journal.pone.0128064](https://doi.org/10.1371/journal.pone.0128064) PMID: [26083474](https://pubmed.ncbi.nlm.nih.gov/26083474/)
20. Prevaes SM, de Winter-de Groot KM, Janssens HM, de Steenhuijsen Piters WA, Tramper-Stranders GA, Wyllie AL, et al. Development of the Nasopharyngeal Microbiota in Infants with Cystic Fibrosis. *American journal of respiratory and critical care medicine*. 2015.

21. Cremers AJ, Zomer AL, Gritzfeld JF, Ferwerda G, van Hijum SA, Ferreira DM, et al. The adult nasopharyngeal microbiome as a determinant of pneumococcal acquisition. *Microbiome*. 2014; 2:44. doi: [10.1186/2049-2618-2-44](https://doi.org/10.1186/2049-2618-2-44) PMID: [25671106](https://pubmed.ncbi.nlm.nih.gov/25671106/)
22. Allen EK, Koepffel AF, Hendley JO, Turner SD, Winther B, Sale MM. Characterization of the nasopharyngeal microbiota in health and during rhinovirus challenge. *Microbiome*. 2014; 2:22. doi: [10.1186/2049-2618-2-22](https://doi.org/10.1186/2049-2618-2-22) PMID: [25028608](https://pubmed.ncbi.nlm.nih.gov/25028608/)
23. Biesbroek G, Bosch AA, Wang X, Keijsers BJ, Veenhoven RH, Sanders EA, et al. The impact of breast-feeding on nasopharyngeal microbial communities in infants. *American journal of respiratory and critical care medicine*. 2014; 190(3):298–308. doi: [10.1164/rccm.201401-0073OC](https://doi.org/10.1164/rccm.201401-0073OC) PMID: [24921688](https://pubmed.ncbi.nlm.nih.gov/24921688/)
24. Sakwinska O, Bastic Schmid V, Berger B, Bruttin A, Keitel K, Lepage M, et al. Nasopharyngeal microbiota in healthy children and pneumonia patients. *Journal of clinical microbiology*. 2014; 52(5):1590–4. doi: [10.1128/JCM.03280-13](https://doi.org/10.1128/JCM.03280-13) PMID: [24599973](https://pubmed.ncbi.nlm.nih.gov/24599973/)
25. Clemente JC, Ursell LK, Parfrey LW, Knight R. The impact of the gut microbiota on human health: an integrative view. *Cell*. 2012; 148(6):1258–70. doi: [10.1016/j.cell.2012.01.035](https://doi.org/10.1016/j.cell.2012.01.035) PMID: [22424233](https://pubmed.ncbi.nlm.nih.gov/22424233/)
26. Cho I, Blaser MJ. The human microbiome: at the interface of health and disease. *Nature reviews Genetics*. 2012; 13(4):260–70. doi: [10.1038/nrg3182](https://doi.org/10.1038/nrg3182) PMID: [22411464](https://pubmed.ncbi.nlm.nih.gov/22411464/)
27. Caporaso JG, Lauber CL, Costello EK, Berg-Lyons D, Gonzalez A, Stombaugh J, et al. Moving pictures of the human microbiome. *Genome Biol*. 2011; 12(5):R50. doi: [10.1186/gb-2011-12-5-r50](https://doi.org/10.1186/gb-2011-12-5-r50) PMID: [21624126](https://pubmed.ncbi.nlm.nih.gov/21624126/)
28. Benton AS, Wang Z, Lerner J, Foerster M, Teach SJ, Freishtat RJ. Overcoming heterogeneity in pediatric asthma: tobacco smoke and asthma characteristics within phenotypic clusters in an African American cohort. *The Journal of asthma: official journal of the Association for the Care of Asthma*. 2010; 47(7):728–34.
29. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol*. 2013; 79(17):5112–20. doi: [10.1128/AEM.01043-13](https://doi.org/10.1128/AEM.01043-13) PMID: [23793624](https://pubmed.ncbi.nlm.nih.gov/23793624/)
30. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al. Introducing mothur: Open-Source, Platform-Independent, Community-Supported Software for Describing and Comparing Microbial Communities. *Appl Environ Microb*. 2009; 75(23):7537–41.
31. Schloss PD, Gevers D, Westcott SL. Reducing the effects of PCR amplification and sequencing artifacts on 16S rRNA-based studies. *PloS one*. 2011; 6(12):e27310. doi: [10.1371/journal.pone.0027310](https://doi.org/10.1371/journal.pone.0027310) PMID: [22194782](https://pubmed.ncbi.nlm.nih.gov/22194782/)
32. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics*. 2011; 27(16):2194–200. doi: [10.1093/bioinformatics/btr381](https://doi.org/10.1093/bioinformatics/btr381) PMID: [21700674](https://pubmed.ncbi.nlm.nih.gov/21700674/)
33. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microb*. 2007; 73(16):5261–7.
34. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods*. 2010; 7(5):335–6. doi: [10.1038/nmeth.f.303](https://doi.org/10.1038/nmeth.f.303) PMID: [20383131](https://pubmed.ncbi.nlm.nih.gov/20383131/)
35. Price MN, Dehal PS, Arkin AP. FastTree 2—Approximately Maximum-Likelihood Trees for Large Alignments. *PloS one*. 2010; 5(3).
36. Faith DP. Conservation evaluation and phylogenetic diversity. *Biol Conserv*. 1992; 61:1–10.
37. Gower JC. Generalized Procrustes Analysis. *Psychometrika*. 1975; 40(1):33–51.
38. Expert Panel Report 3 (EPR-3). Guidelines for the Diagnosis and Management of Asthma—Summary Report 2007. *J Allergy Clin Immun*. 2007; 120(5 Suppl):S94–S138. PMID: [17983880](https://pubmed.ncbi.nlm.nih.gov/17983880/)
39. Dixon P. VEGAN, a package of R functions for community ecology. *J Veg Sci*. 2003; 14(6):927–30.
40. Parks DH, Tyson GW, Hugenholtz P, Beiko RG. STAMP: statistical analysis of taxonomic and functional profiles. *Bioinformatics*. 2014; 30(21):3123–4. doi: [10.1093/bioinformatics/btu494](https://doi.org/10.1093/bioinformatics/btu494) PMID: [25061070](https://pubmed.ncbi.nlm.nih.gov/25061070/)
41. RStudioTeam. RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL <http://www.rstudio.com/>. 2015.
42. Pérez-Losada M, Crandall KA, Freishtat RJ. Comparison of two commercial DNA extraction kits for the analysis of nasopharyngeal bacterial communities. *AIMS Microbiology*. 2016; 2(2):108–19.
43. Huang YJ, Nelson CE, Brodie EL, Desantis TZ, Baek MS, Liu J, et al. Airway microbiota and bronchial hyperresponsiveness in patients with suboptimally controlled asthma. *The Journal of allergy and clinical immunology*. 2011; 127(2):372–81 e1–3. doi: [10.1016/j.jaci.2010.10.048](https://doi.org/10.1016/j.jaci.2010.10.048) PMID: [21194740](https://pubmed.ncbi.nlm.nih.gov/21194740/)

44. Cardenas PA, Cooper PJ, Cox MJ, Chico M, Arias C, Moffatt MF, et al. Upper airways microbiota in anti-biotic-naive wheezing and healthy infants from the tropics of rural Ecuador. *PloS one*. 2012; 7(10): e46803. doi: [10.1371/journal.pone.0046803](https://doi.org/10.1371/journal.pone.0046803) PMID: [23071640](https://pubmed.ncbi.nlm.nih.gov/23071640/)
45. Yatsunenko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, et al. Human gut microbiome viewed across age and geography. *Nature*. 2012; 486(7402):222–+. doi: [10.1038/nature11053](https://doi.org/10.1038/nature11053) PMID: [22699611](https://pubmed.ncbi.nlm.nih.gov/22699611/)
46. Backhed F, Fraser CM, Ringel Y, Sanders ME, Sartor RB, Sherman PM, et al. Defining a Healthy Human Gut Microbiome: Current Concepts, Future Directions, and Clinical Applications. *Cell Host & Microbe*. 2012; 12(5):611–22.
47. Shade A, Handelsman J. Beyond the Venn diagram: the hunt for a core microbiome. *Environ Microbiol*. 2012; 14(1):4–12. doi: [10.1111/j.1462-2920.2011.02585.x](https://doi.org/10.1111/j.1462-2920.2011.02585.x) PMID: [22004523](https://pubmed.ncbi.nlm.nih.gov/22004523/)
48. Shafquat A, Joice R, Simmons SL, Huttenhower C. Functional and phylogenetic assembly of microbial communities in the human microbiome. *Trends Microbiol*. 2014; 22(5):261–6. doi: [10.1016/j.tim.2014.01.011](https://doi.org/10.1016/j.tim.2014.01.011) PMID: [24618403](https://pubmed.ncbi.nlm.nih.gov/24618403/)
49. Biesbroek G, Wang X, Keijsers BJ, Eijkemans RM, Trzcinski K, Rots NY, et al. Seven-valent pneumococcal conjugate vaccine and nasopharyngeal microbiota in healthy children. *Emerging infectious diseases*. 2014; 20(2):201–10. doi: [10.3201/eid2002.131220](https://doi.org/10.3201/eid2002.131220) PMID: [24447437](https://pubmed.ncbi.nlm.nih.gov/24447437/)
50. Pérez-Losada M, Crandall KA, Freishtat RJ. Two sampling methods yield distinct microbial signatures in the nasopharynx of asthmatic children. *Microbiome*. 2016 (in press).