



POSTER PRESENTATION

Open Access

MISIPI study: Melanoma ImmunoScore evaluation in patients treated with Ipilimumab

Carlo Bifulco^{1*}, Marilena Capone², Zip Feng¹, Gabriele Madonna², Ester Simeone², Marcello Curvietto², Nicola Mozzillo², Gennaro Ciliberto², Gerardo Botti², Bernard A Fox¹, Paolo A Ascierto²

From Melanoma Bridge meeting 2013
Naples, Italy. 5-8 December 2013

Background

Increasing evidence has supported the hypothesis that cancer development is influenced by the host immune system. Immune infiltrates of the primary tumors and metastases are independent prognostic biomarkers and may represent predictive factors, suggesting that pre-therapeutic immune response determines the efficacy of different immunotherapies [1]. An immune-classification of tumors based on a scoring system of density and location of immune-cells within the tumor has been recently proposed in human colorectal cancers, where a high density of CD3+, CD8+ memory T lymphocytes in the primary tumor is associated with long disease free and overall survival and low risk of relapse and metastasis [2,3]. The impact of the immune score needs to be evaluated in other tumor types given the universal importance of the immune system in cancers. We have focused on the prognostic and predictive value of the immunoscore in patients with advanced melanoma treated with Ipilimumab in the previous 2-3 years, and correlated it with outcomes and response to the treatment.

Materials and methods

We have collected 190 FFPE metastatic samples from melanoma patients treated with Ipilimumab, using the last lesion excised before the treatment: lymph node involvement or distant metastatic sites. We are characterizing the immunoscore with the expression of CD3, CD8, CD20, Foxp3, by immunohistochemistry and evaluated correlation with clinical outcome. Serial sections (3-4 micron in thickness) have been stained with each marker and counter-stained with H/E. Moreover, one section has been stained

with a multiplex of all markers including tumor marker for melanoma (S100). All slides will be digitized on a Leica SCN-400 system, and the results will be quantified by assessing the density of different cellular immune population at the center and invasive margin of the tumor using a digital images analysis application (Definiens Architect XD). The markers expression will be subsequently matched with the most important clinical information about all the patients.

Results

We have considered the clinical benefit from treatment with Ipilimumab as the parameter for evaluating the effectiveness of immunoscore. The treatment has been considered as effective if a patient met a partial or complete response, or had stable disease for more than 6 months from the start of therapy, or with an overall survival > 1 year. The evaluation of prognostic and predictive power of the immunoscore, in patients with metastatic melanoma, treated with ipilimumab, are nearing completion and definition.

Conclusion

We believe that the immunoscore is key as prognostic and predictive markers for immunotherapies in metastatic melanoma.

Authors' details

¹Earle A. Chiles Research Institute – Providence Cancer Center, Portland, OR, USA. ²Istituto Nazionale Tumori Fondazione “G. Pascale”, Napoli, Italy.

Published: 6 May 2014

References

1. Galon J, Pagès F, Marincola FM, Thurin M, Trinchieri G, Fox BA, Gajewski TF, Ascierto PA: **The immune score as a new possible approach for the classification of cancer.** *J Transl Med* 2012, **10**:1.

* Correspondence: carlo.bifulco@providence.org

¹Earle A. Chiles Research Institute – Providence Cancer Center, Portland, OR, USA

Full list of author information is available at the end of the article

2. Galon J, Costes A, Sanchez-Cabo F, Kirilovsky A, Mlecnik B, Lagorce-Pages C, Tosolini M, Camus M, Berger A, Wind P, *et al*: **Type, density, and location of immune cells within human colorectal tumors predict clinical outcome.** *Science* 2006, **313**:1960-1964.
3. Pages F, Kirilovsky A, Mlecnik B, Asslaber M, Tosolini M, Bindea G, Lagorce C, Wind P, Marliot F, Bruneval P, *et al*: **In situ cytotoxic and memory T cells predict outcome in patients with early-stage colorectal cancer.** *J Clin Oncol* 2009, **27**:5944-5951.

doi:10.1186/1479-5876-12-S1-P11

Cite this article as: Bifulco *et al.*: MISIPI study: Melanoma ImmunoScore evaluation in patients treated with IPilimumab. *Journal of Translational Medicine* 2014 **12**(Suppl 1):P11.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

