



Contents lists available at ScienceDirect

Saudi Journal of Biological Sciences

journal homepage: www.sciencedirect.com



Original article

Polychromatic luminescence and improved antifungal performance of succinic acid in the lattice of L-Lysine monohydrochloride



B. Aneeba ^{a,*}, S.V. Ashvin Santhia ^a, R. Sheela Christy ^a, Fahad A. Al-Misned ^b, Hamed A. El-Serehy ^b, S. Vinu ^{c,*}

^a Department of Physics and Research Centre, Nesamony Memorial Christian College, Marthandam, Affiliated to Manonmaniam Sundaranar University, Abishekappatti, Tirunelveli, Tamilnadu, India

^b Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^c Department of Physics, Govt. Arts and Science College, Nagercoil, Tamilnadu 629004, India

ARTICLE INFO

Article history:

Received 20 August 2020

Revised 14 October 2020

Accepted 15 October 2020

Available online 24 October 2020

Keywords:

Succinic acid

Monoclinic

Transmittance

Candida albicans

LED

ABSTRACT

The incorporation of succinic acid (SA) in the lattice of L-Lysine monohydrochloride (LM) has opened the new avenue in the field of production and application of scintillator materials such as LED and antifungal drug. Crystalline trait and monoclinic structure were scanned by XRD. The existence of carbonyl, carboxylate and protonated amine group were confirmed through FTIR and UV spectra predicted the transmittance of SA: LM crystal. Polychromatic luminescence behaviour had achieved through the incorporation of SA instead of blue luminescence, which is a new result. Also SA: LM exhibited good response towards pathogenic fungi which causes numerous types of infections and diseases in both humans and animals. The high inhibitory zone at 16 mm was formed by the grown SA: LM crystal against the life threatening fungi like *Candida albicans*. Also fungal inhibition against *candida parapsilosis* and *Aspergillus flavus*, respectively, were tuned by the inclusion of succinic acid.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

L-lysine is one of the acentric crystal because the basic chain is made with optically active carbon atom. Also the specular character of this L-Lysine is due to its phonon subsystem with hydrogen bonds (Krishnakumar et al., 2010). Natural characteristic cationic lysine is the electro optic material likewise it has wide potential character like pharmacologic movement, nonlinear optical impact, strong electrostatic interaction and optical parametrical effect as well as it is an indispensable amino acid to the human body especially in the formation of collagen (Rubin et al., 1960; Ozga et al., 2008; Lasser et al., 1960; Dousa et al., 2011; Wang et al., 2018; Cao et al., 2016). Succinic acid is come under the category of metabolite and biologically dynamic carboxylic acids (Kondratenko et al., 2020) formed by microorganisms, animals and plants (Juliet and Subramanian, 2018). Complexes of succinic acid plays a vital role to deliver inhibitory action against microor-

ganism and utilized in pharmaceutical industry (Demir et al., 2010a, 2010b; Lin et al., 2008; Ahn et al., 2016). Furthermore, it can be utilized in the creation of infrared (IR) MALDI analytical methods (Krishnan et al., 2008).

Our environment consists of very diverse microorganisms that include all bacteria, virus and fungus so on. It can be found in rock surfaces, soil, inside rocks, toxic waste, buried mites under the earth and almost in every parts of the atmosphere. The property of toxin production and penetration causes various types of infections, diseases in both humans and animals. Among all microorganism, fungus and bacteria possess greatest threat to all lives because they could cause illness in them (Boehlke et al., 2020; Calzavara-Pinton et al., 2005). The synthesis of antifungal and antibacterial drugs is lower in contrast to the rate of spreading of disease by both fungus and bacteria. So with basic characterization here, concentration given to magnify the optical, luminescence, chemical and biological activities of the grown crystals.

* Corresponding authors.

E-mail addresses: aneebaanu@gmail.com (B. Aneeba), vinusnist@gmail.com (S. Vinu).

Peer review under responsibility of King Saud University.

2. Material and methods

2.1. Materials

Sigma Aldrich Chemicals delivered the analytical grade materials such as L-Lysine monohydrochloride ($C_6H_{15}ClN_2O_2$) and Succinic acid ($C_4H_6O_4$). Deionised water was used as solvent.



Production and hosting by Elsevier

2.2. Growth method of crystals

To grow the crystal under atmospheric condition, slow evaporation is the finest method. This method was used to grow SA doped LM single crystal in the molecular ratio 0.02: 0.98. Enumerate amount of L-Lysine monohydrochloride and succinic acid was used to prepare supersaturated solution. Filtered saturated solution was lidded and kept for slow evaporation. Colourless crystals were obtained nearly after 20 days. Fig. 1. shows the grown SA: LM crystal.

2.3. Characterization of crystals

The structure was found by ENRAF NONIUS CAD4 X-ray diffractometer and the powder XRD of the sample were recorded with the help of XPert Pro- PAalytic powder diffractometer. Using Thermo Nicolet avatar 370 spectrometer, vibrational studies was carried out in the middle infrared region ($400\text{--}4000\text{ cm}^{-1}$). Optical property of SA: LM crystals were assessed in the UV–Visible region by means of Perkin Elmer (Lambda 35) spectrometer. Polychromatic light emission of sample was appraised by make use of Varian Cary Eclipse Photo Luminescence Spectrophotometer. Fungal activity test was undertaken by disk diffusion method.

3. Results

Crystalline quality of the grown samples are identified from the sharp narrow diffractions peaks obtained by powder XRD analysis (Fig. 2) and Phase analysis demonstrate that crystal SA: LM is monoclinic. The various vibrational analysis of the samples are obtained from Fig. 3 and the peculiar vibration of amino acid due to NH_3 stretching is obtained in the range 2924, 1585 and 1506 cm^{-1} . The synthesized crystals exhibit good inhibitory action against the life threatening pathogens which are shown in Fig. 4. SA: LM crystal shows a notable action against *Candida* species such as *Candida albicans*, *candida parapsilosis*, and *Aspergillus flavus* by forming 16 mm, 14 mm, and 16 mm diameter zone. The transmittance behaviour of SA doped LM crystal is depicted in Fig. 5. The SA doped LA crystal shows 67 percentage of transmittance. At 345 nm wavelength the sample is excited and the emission spectrum was recorded, it is given in Fig. 6. Also it revealed the various luminescence property at different wavelength.




Fig. 1. Photograph of SA: LM crystal.




Fig. 2. Powder XRD pattern of SA: LM crystal.

4. Discussion

4.1. X-Ray diffraction analysis

The structure of L- Lysine monohydrochloride was not affected by the small amount of SA doping undoubtedly cleared from the phase analysis which does not shows any secondary phases. Phase analysis demonstrate that crystal SA: LM is monoclinic with unit cell dimensions $a = 5.88\text{ \AA}$, $b = 13.32\text{ \AA}$, $c = 7.49\text{ \AA}$, $\alpha = \gamma = 90^\circ$, $\beta = 97.79^\circ$ and $V = 581.271\text{ \AA}^3$ (Ramesh Babu et al., 2006). This is in agreement with the SP_2 space group. The successful incorporation of SA ion in LM is recognised from the shrinkage in the lattice parameter, subtle shift in intensity of peaks and diffraction angle.

4.2. Vibrational analysis

The C—C—N symmetric stretching obtained at the wave number 862 cm^{-1} (Robert et al., 2010). The absorption peak at 3171 cm^{-1} is assigned to N-H bond stretching (Petrosyan and Ghazaryan, 2009). According to Kalaiselvi et al the COO^- asymmetric stretching and NH_3 rocking were observed in the region 1619 and 1182 cm^{-1} (Kalaiselvi et al., 2007). In SA: LM also same vibration obtained at 1619 and 1182 cm^{-1} . C—N vibration imputed in the wave numbers 1348 and 1320 cm^{-1} (Ozga et al., 2008) and C—C stretching is noticed in the wave number 997 cm^{-1} . 3422 cm^{-1} is assigned to stretching vibration of water molecule. Krishnakumar et al reported that at $3450\text{--}3250\text{ cm}^{-1}$ stretching vibration of H_2O molecule take place. C—O stretching vibration is noticed at 1216 cm^{-1} (Krishnakumar et al., 2010). These wave numbers slightly shifted from the reference values which is due to the addition of SA.

4.3. Fungal activity

The *Candida* species are unicellular fungi which is present in humans at gastrointestinal, mucosal surfaces of the oral cavity, skin, vagina and urogenital tracts (Tsai et al., 2013). But on the other hand it is also a threat to human health by causing oropharyngeal candidiasis (OPC) and vulvovaginal candidiasis (VVC) (Coleman et al., 1993; Vargas and Joly, 2002; Sanchez et al., 1993, Hosseini et al., 2020). Furthermore VVC is a major threat for women infected with HIV. In United States by substantiation, the candida infection is the fourth most nosocomial infection and ranked second in death due to that infection (Pfaller et al., 1998; David Trofa et al., 2008). Aspergillosis is an infection brought about




Fig. 3. Vibrational spectra of SA: LM crystal.




Fig. 4. Zone of inhibition of SA: LM against (a) *candida parapsilosis* (b) *Candida albicans* and (c) *Aspergillus flavus*.




Fig. 5. Transmittance spectra of SA: LM crystal.




Fig. 6. Photoluminescence spectrum of SA: LM crystal.

by sort of fungi in particular *Aspergillus flavus* ordinarily trouble the respiratory framework (Smith and Denning, 2010; Ramirez-Camejo et al., 2012). Due to these infections on humans by fungi the curiosity to find antifungal drug is aroused. 16 mm extent of inhibitory zone was formed in opposition to *Candida albicans* by SA: LM which is more than that of antifungal activity of Nystatin. This Nystatin is used to patients who is affected by fungal infection (Ma et al., 2007; Monteiro et al., 2013; Zahabi et al., 2020). Moreover SA: LM material exhibit inhibitory action against *candida parapsilosis* and *Aspergillus flavus* by forming 14 mm and 16 mm diameter zone, respectively, compared to *Candida albicans* in the diameter 9 mm reported by other authors (Kalaiselvi et al., 2018). Nano silver (Tantyani and Taufikurohmah, 2020) also showed less inhibitory zone than SA: LM crystal. These resisting activity of the sample against fungi leads to the development of effective anti-fungal drugs for wide range of applications.

4.4. Optical analysis

Depending on defects, orientation, optically active functional groups the range of transmittance varied (Anis et al., 2019; Senthil Pandian et al., 2011). The SA doped LM crystal shows about 67 percentage of transmittance and it is less than pure LM which shows 70 percentage transmittance (Aneeba et al., 2020). Because the concentration of doping reduced the transmittance of the crys-

tal due to the creation of extra defects (Zhou et al., 2019). This defects opened the way to SA: LM crystal towards polychromatic luminescence.

optical material:

L-lysine hydrochloride dihydrate

D. Kalaiselvi, R. Mohan Kumar, R. Jayavel

*

Crystal Growth Centre, Anna University, Chennai 600025, India

Received 8 May 2007; received in revised form 22 June 2007; acc

5. anti-microbial activities

5.1. Photoluminescence study

The typical approach for characterizing defects and other imperfection present in the sample is studied by using Photoluminescence Spectroscopic (PL) technique. It enables to examine the electronic transition and optically active intrinsic defect densities occurs due to absorption of light (Alexandar et al., 2017; Rajesh et al., 2017; Shejwal et al., 2016). The sample LM has a sturdy luminescence behaviour, which itself has its potential to act as a scintillator substance (chen et al., 2016; Rani et al., 2013). The PL of SA: LM yields information about multiple energy levels existing between VB and CB which is accountable for radioactive recombination. The high intense PL emission of SA: LM is at 535 and 544 nm which are accountable for the powerful green luminescence. The first two peaks shows indigo and blue emission in the relative wavelength 459 and 494 nm and the rest shows yellow and orange emission at 572, 582 and 595 nm. The blue emission is due to the presence of NH₂ and Cl ions present in the sample. Chen et al depicted the luminescence behaviour of lysine, which is in the wavelength range 490 nm corresponding to blue emission (Chen et al., 2016). Notwithstanding in addition to above emission, SA: LM emitted the following colours like violet, indigo, yellow, orange owing to the inclusion of succinic acid. Because many of the literature showed the effect of SA in the luminescence property (Zhang et al., 2008; Wang et al., 2009; Cui et al., 2005; Demir et al., 2010ab). The enhancement of polychromatic emission of lysine is due to the doping of succinic acid.

6. Conclusion

The antifungal and high photo luminescence crystal SA: LM was synthesised and its crystalline properties were examined by XRD, FTIR, UV and photoluminescence spectroscopy. The monoclinic structure and functional groups are identified from XRD and FTIR. The formation of defects in grown crystal were identified from the transmittance graph. The intense responsive action against these pathogens proved that the succinic acid doped L-Lysine monohydrochloride could act as a potential antifungal drug in the field of medicine. The UV visible spectrum reveals that the crystals transmittance percentage was reduced by the defects present in them. The Photoluminescence characterisation shows the reinforced emission of violet, indigo, blue, green, yellow and orange colours due to polychromatic luminescence. These leading properties in SA: LM crystal opens the way in the field of antifungal drugs and also an effective supernumerary in the fabrication of scintillator materials.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors wish to thank STIC, Cochin, SAIF, IIT, Chennai, and St. Joseph College, Tiruchirappalli for the help in taking various analysis. The authors would like to extend their sincere appreciation to the Researchers Supporting Project Number (RSP-2020/24), King Saud University, Riyadh, Saudi Arabia.

References

- Alexandar, A., Lakshmanan, A., Sakthy Priya, S., Surendran, P., Rameshkumar, P., 2017. Growth, structural, optical, piezoelectric and etching analysis of L-lysine p-nitrophenolate monohydrate single crystals. *Int. J. Mod. Phys. B.* 31 (23), 1750174.
- Aneeba, B., Ashvin Santhia, S.V., Vinu, S., Sheela Christy, R., Dunia A. Al Farraj., Noorah A. Alkubaisi., 2020. Influence of most reactive inorganic cation in the optical and biological activities of L-Lysine monohydrochloride crystal. *Saudi J. Biolog. Sci.* 27 (2020) 2961–2967. <https://doi.org/10.1016/j.sjbs.2020.07.018>.
- Anis, M., Baig, M.I., Muley, G.G., AlFaify, S., Khan, M.A., 2019. Impact of increasing concentration of L-alanine environment on structural, UV-visible, SHG efficiency, luminescence and dielectric traits of zinc thiourea chloride (ZTC) crystal. *Optik* 185, 317–324.
- Boehlke, C., Rupf, S., Tenniswood, M., Chittur, S.V., Hannig, C., Zierau, O., 2020. Caries and periodontitis associated bacteria are more abundant in human saliva compared to other great apes. *Arch. Oral Biol.* 111, 104648.
- Calzavara-Pinton, P.G., Venturini, M., Sala, R., 2005. A comprehensive overview of photodynamic therapy in the treatment of superficial fungal infections of the skin. *J. Photochem. Photobiol. B*. 78 (1), 1–6.
- Cao, J., Ding, L., Zhang, Y., Wang, S., Fang, Y., 2016. A ternary sensor system based on pyrene derivative-SDS assemblies-Cu²⁺ displaying dual responsive signals for fast detection of arginine and lysine in aqueous solution. *J. Photochem. Photobiol. A*. 314, 66–74.
- Chen, Y., Bao, Y., Wang, X., 2016. Green Emission of Tb-doped Mg-Al Layered Double Hydroxide Response to L-lysine. *J. Fluoresc.* 26 (3), 813–820.
- Coleman, D.C., Bennett, D.E., Sullivan, D.J., Gallagher, P.J., Henman, M.C., Shanley, D. B., 1993. Oral Candida in HIV infection and AIDS: new perspectives/new approaches. *Crit. Rev. Microbiol.* 19 (2), 61–82.
- Cui, G.-H., Li, J.-R., Zhang, R.-H., Bu, X.-H., 2005. Hydrothermal synthesis, crystal structures and luminescent properties of two new Ln (III)-succinate (Ln=Eu, Tb) complexes exhibiting three dimensional networks. *J. Mol. Struct.* 740 (1–3), 187–191.
- Trofa, D., Gacsér, A., Nosanchuk, J.D., 2008. Candida parapsilosis, an emerging fungal pathogen. *Clin. Microbiol. Rev.* 21 (4), 606–625.
- Demir, S., Yilmaz, V.T., Sariboga, B., Buyukgungor, O., Mrozinski, J., 2010a. Metal (II) Nicotinamide Complexes Containing Succinate, Succinate and Succinic Acid: Synthesis, Crystal Structures, Magnetic, Thermal, Antimicrobial and Fluorescent Properties. *J. Inorg. Organomet. Polym. Mater.* 20 (2), 220–228.
- Dousa, M., Brichac, J., Gibala, P., Lehnert, P., 2011. Rapid hydrophilic interaction chromatography determination of lysine in pharmaceutical preparations with fluorescence detection after postcolumn derivatization with o-phthalodialdehyde. *J. Pharmaceut. Biomed.* 54 (5), 972–978.
- Hosseini, S.S., Joshaghani, H., Shokohi, T., Ahmadi, A., Mehrbakhsh, Z., 2020. Antifungal Activity of ZnO Nanoparticles and Nystatin and down regulation of SAP1-3 Genes Expression in Fluconazole-Resistant *Candida albicans* Isolates from Vulvovaginal Candidiasis. *Infect. Drug Resist.* 13, 385–394.
- Juliet sheela, K., Subramanian, P., 2018. Influence of VO²⁺ ions on Structural and optical properties of potassium succinate-succinic acid single crystal for nonlinear optical applications. *Physica B*. 534, 156–162.
- Ahn, Jung Ho, Jang, Yu-Sin, Yup, Sang, 2016. Lee Production of succinic acid by metabolically engineered microorganisms. *Curr. Opin. Biotech.* 42, 54–66.
- Kalaiselvi, D., Mohankumar, R., Jayavel, R., 2007. Crystal growth, thermal and optical studies of semiorganic nonlinear optical material: L-lysine hydrochloride dehydrate. *Mater. Res. Bull.* 43, 1829–1835.
- Kalaiselvi, V., Mathammal, R., Vijayakumar, S., Vaseeharan, B., 2018. Microwave assisted green synthesis of Hydroxyapatite nano rods using *Moringa oleifera* flower extract and its antimicrobial applications. *I.J.v.s.m.* 6 (2), 286–295.
- Kondratenko, Yu.A., Nikonorova, A.A., Zolotarev, A.A., Ugolkov, V.L., Kochina, T.A., 2020. Tris (hydroxymethyl) methyl ammonium salts of biologically active carboxylic acids. Synthesis, properties and crystal structure. *J. Mol. Struct.* 1207, 127813.
- Krishnakumar, V., Nagalakshmi, R., Manohar, S., Ozga, K., Claudet, B., Piasecki, M., Pisarek, J., 2010. Elasto-optical spectra of novel L-lysine monohydrochloride dehydrate single crystals. *Int. J. Mod. Phys. B*. 24 (05), 629–645.
- Krishnan, S., Justin Raj, C., Das, S.J., 2008. Growth and characterization of novel ferroelectric urea-succinic acid single crystals. *J. Cryst. Growth*. 310 (14), 3313–3317.
- Lasser, R.P., Schoenfeld, M.R., Friedberg, C.K., 1960. L-Lysine Monohydrochloride: A clinical study of its action as a chloruretic acidifying adjuvant to mercurial diuretics. *N. Engl. J. Med.* 263 (15), 728–733.
- Ma, C.-M., Kully, M., Khan, J.K., Hattori, M., Daneshbalab, M., 2007. Synthesis of chlorogenic acid derivatives with promising antifungal activity. *Bioorg. Med. Chem.* 15 (21), 6830–6833.
- Monteiro, D.R., Silva, S., Negri, M., Gorup, L.F., de Camargo, E.R., Oliveira, R., Henriques, M., 2013. Antifungal activity of silver nanoparticles in combination with nystatin and chlorhexidine digluconate against *Candida albicans* and *Candida glabrata* biofilms. *Mycoses* 56 (6), 672–680.
- Ozga, K., Krishnakumar, V., Kityk, I.V., Jasik-Slezak, J., 2008. L-lysine monohydrochloride dihydrate as novel elasto- and electrooptical materials. *Mater. Lett.* 62, 4597–4600.
- Petrosyan, A.M., Ghazaryan, V.V., 2009. Vibrational spectra of L-lysine monohydrochloride dihydrate and its two anhydrous forms. *J. Mol. Struct.* 917 (1), 56–62.
- Pfaller, M.A., Jones, R.N., Messer, S.A., Edmond, M.B., Wenzel, R.P., 1998. National Surveillance of Nosocomial Blood Stream Infection Due to Species of *Candida* Other than *Candida albicans*: Frequency of Occurrence and Antifungal Susceptibility in the SCOPE Program. *Diagn. Microbiol. Infect. Dis.* 30 (2), 121–129.
- Rajesh, P., Rao, G.B., Ramasamy, P., 2017. Effect of Rochelle salt on growth, optical, photoluminescence, photoconductive and piezoelectric properties of the triglycine sulphate single crystal. *J. Cryst. Growth*. 468, 340–344.
- Ramesh Babu, R., Vijayan, N., Gopalakrishnan, R., Ramasamy, P., 2006. Growth and characterization of L-lysine monohydrochloride dihydrate (L-LMHCl) single crystal. *Cryst. Res. Technol.* 41 (4), 405–410.
- Ramirez-Camejo, L.A., Zuluaga-Montero, A., Lazaro-Escudero, M., Hernandez-Kendall, V., Bayman, P., 2012. Phylogeny of the cosmopolitan fungus *Aspergillus flavus*: is everything everywhere?. *Fungal Biol.* 116 (3), 452–463.
- Rani, N., Vijayan, N., Thukral, K., Maurya, K.K., Haranath, D., Bhagavannarayana, G., Wahab, M.A., 2013. Crystalline perfection, optical and third harmonic generation analyses of non-linear optical single crystal of L-lysine acetate. *Spectrochim Acta A Mol Biomol Spectrosc.* 105, 192–199.
- Robert, R., Raj, C.J., Das, S.J., 2010. Growth and characterization of pure and doped L-Lysine monohydrochloride dihydrate (L-LMHCl) nonlinear optical single crystals. *Curr. Appl. Phys.* 10 (2), 670–675.
- Rubin, A.L., Spritz, N., Mead, A.W., Herrmann, R.A., Braverman, W.S., Luckey, E.H., 1960. The Use of L-Lysine Monohydrochloride in Combination with Mercurial Diuretics in the Treatment of Refractory Fluid Retention. *Circulation* 21 (3), 332–336.
- Sanchez, V., Vazquez, J.A., Barth-Jones, D., Dembry, L., Sobel, J.D., Zervos, M.J., 1993. Nosocomial acquisition of *Candida parapsilosis*: An epidemiologic study. *Am. J. Med. Sci.* 94 (6), 577–582.
- Selcuk Demir, Y., Veysel, T., Yilmaz, B., Bahtiyar Sariboga, Orhan Buyukgungor, Jerzy Mrozniski, 2010. Metal (II) Nicotinamide Complexes Containing Succinate, Succinate and Succinic Acid: Synthesis, Crystal Structures, Magnetic, Thermal, Antimicrobial and Fluorescent Properties. *J. Inorg. Organomet. Polym.* 20(2), 220–228.
- Senthil Pandian, M., Pattanaboonmee, N., Ramasamy, P., Manyum, P., 2011. Studies on conventional and Sankaranarayanan-Ramasamy (SR) method grown ferroelectric glycine phosphite (GPI) single crystals. *J. Cryst. Growth*. 314 (1), 207–212.
- Shejwal, N.N., Anis, M., Hussaini, S.S., Shirsat, M.D., 2016. Ab-initio UV-vis., SHG efficiency, photoluminescence, dielectric, thermal and mechanical studies of pure and L-glutamine doped ZTS crystal for optoelectronics device applications. *Optik* 127 (16), 6525–6531.
- Smith, N.L., Denning, D.W., 2010. Underlying conditions in chronic pulmonary aspergillosis including simple aspergilloma. *Eur. Respir. J.* 37 (4), 865–872.
- Lin, Sze Ki Carol, Chenyu, Du., Koutinas, Apostolis, Wang, Ruohang, Webb, Colin, 2008. Substrate and product inhibition kinetics in succinic acid production by *Actinobacillus succinogenes*. *Biochem. Eng. J.* 41, 128–135.
- Tantyan, T.A., Taufikurohmah, T., 2020. Antibacterial and antifungal activity of silver nanoparticles against *neisseria gonorrhoeae* and *candida albicans*. *Diagn. Microbiol. Infect. Dis.* 8 (6), 179–187.
- Tsai, P.W., Chen, Y.T., Hsu, P.C., Lan, C.Y., 2013. Study of *Candida albicans* and its interactions with the host: A mini review. *Bio Medicine*, 3 (1), 51–64.
- Vargas, K.G., Joly, S., 2002. Carriage frequency, intensity of carriage, and strains of oral yeast species vary in the progression to oral candidiasis in human immunodeficiency virus-positive individuals. *J. Clin. Microbiol.* 40 (2), 341–350.
- Wang, C.-G., Xing, Y.-H., Li, Z.-P., Li, J., Zeng, X.-Q., Ge, M.-F., Niu, S.-Y., 2009. Synthesis, crystal structures and properties of a series of three-dimensional lanthanide coordination polymers with the rigid and flexible mixed dicarboxylate ligands of 1, 4-benzene dicarboxylic acid and succinic acid. *J. Mol. Struct.* 921 (1–3), 126–131.
- Wang, X.R., Huang, Z., Du, J., Wang, X.Z., Gu, N., Tian, X., Ding, B., 2018. Hydrothermal Preparation of Five Rare-Earth (Re = Dy, Gd, Ho, Pr, and Sm) Luminescent Cluster-Based Coordination Materials: The First MOFs-based Ratiometric Fluorescent Sensor for Lysine and Bifunctional Sensing Platform for Insulin and Al3+. *Inorg. Chem.* 57 (20), 12885–12899.
- Zahabi, Z.F., Sharififar, F., Almani, P.G.N., Salari, S., 2020. Antifungal activity of different fractions of *Salvia rhytidia* Benth as a valuable medicinal plant against various species of *Candida* in Kerman Province (southeast Iran). *Gene Rep.* 19, 100624.
- Zhang, X.-J., Xing, Y.-H., Han, J., Zeng, X.-Q., Ge, M.-F., Niu, S.-Y., 2008. A Series of Novel Ln-Succinate-Oxalate Coordination Polymers: Synthesis, Structure, Thermal Stability, and Fluorescent Properties. *J. Cryst. Growth Des.* 8 (10), 3680–3688.
- Zhou, G., Li, G., Lü, Y., Ma, Y., Sun, X., Deng, X., Wang, X., 2019. Growth and characterization of L-phenylalanine doped KDP crystals. *Mater. Res. Bull.* 113, 146–151.