



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Clinical benefits in patients with home-based cardiac rehabilitation in the era of COVID-19 pandemic



ARTICLE INFO

Article history:

Received 9 December 2021
 Received in revised form 15 December 2021
 Accepted 19 December 2021
 Available online 23 December 2021

Keywords:

Home based cardiac rehabilitation
 COVID-19 pandemic
 Clinical outcomes
 Risk assessment

I read the study by Jafri et al.¹ with great interest. The authors evaluated outcomes of patients with or without home-based cardiac rehabilitation (HBCR). Although HBCR was significantly associated with a lower risk of the combined all-cause mortality and all-cause hospitalizations up to 12 months, the adjusted hazard ratios (95% confidence intervals) of HBCR for all-cause mortality, cardiovascular hospitalization, and all-cause hospitalization were 0.43 (0.18–1.0), 0.57 (0.22–1.4), 0.53 (0.28–1.01), respectively. About half of the center-based cardiac rehabilitation (CBCR) programs stopped during the coronavirus disease-19 pandemic.² Jafri et al. pointed out that randomized controlled trials (RCTs) are needed to evaluate the effectiveness of HBCR; therefore, I am presenting some comments about the effectiveness of HBCR based on recent academic papers.

Batalik et al.³ conducted an RCT to evaluate the effectiveness of home-based cardiac telerehabilitation (HBCTR) over a period of 1 year in patients with coronary heart disease (CHD). Although the average of peak oxygen uptake significantly increased, there was no significant change in scores of health-related quality of life (QOL) by intervention. Ramachandran et al.⁴ conducted a meta-analysis of RCTs to evaluate the effectiveness of HBCTR as an alternative to CBCR in patients with CHD. Patients with HBCTR showed a significant improvement in functional capacities such as 6-min walking test distance, daily step count, and exercise habits than patients with usual medical care. In addition, depression scores and QOL scores also improved with HBCTR. Furthermore, HBCTR and CBCR were comparable in terms of their impact on functional capacity, physical activity,

depression, QOL score, medication adherence, cardiac-related hospitalization, and behavioral risk factors, including smoking.

There is not enough data present regarding the effectiveness of HBCR/HBCTR in the literature; therefore, further studies are required. In the era of the coronavirus disease-19 pandemic, valid HBCR/HBCTR may serve as an effective alternative to manage patients with CHD.

Declaration of Competing Interest

There is no conflict of interest in this study.

There is no financial or other relationship with other people or organizations, that may inappropriately influence my work.

References

- Jafri SH, Imran TF, Medbury E, et al. Cardiovascular outcomes of patients referred to home based cardiac rehabilitation. *Heart Lung*. 2022;52:1–7.
- Ghisi GLM, Xu Z, Liu X, et al. Impacts of the COVID-19 pandemic on cardiac rehabilitation delivery around the world. *Glob Heart*. 2021;16(1):43.
- Batalik L, Dosbaba F, Hartman M, Konecny V, Batalikova K, Spinar J. Long-term exercise effects after cardiac telerehabilitation in patients with coronary artery disease: 1-year follow-up results of the randomized study. *Eur J Phys Rehabil Med*. 2021;57(5):807–814.
- Ramachandran HJ, Jiang Y, Tam WWS, Yeo TJ, Wang W. Effectiveness of home-based cardiac telerehabilitation as an alternative to Phase 2 cardiac rehabilitation of coronary heart disease: a systematic review and meta-analysis. *Eur J Prev Cardiol*. 2021 Jul. <https://doi.org/10.1093/eurjpc/zwab106>.

Tomoyuki Kawada, MD,*

Department of Hygiene and Public Health, Nippon Medical School

*Corresponding author at: Department of Hygiene and Public Health, Nippon Medical School, 1-1-5 Sendagi, Bunkyo-Ku, Tokyo 113-8602, Japan.

E-mail address: kawada@nms.ac.jp

Received 9 December 2021

Revised 15 December 2021

Accepted 19 December 2021

Available online 23 December 2021