

miR-26b inhibits isoproterenol-induced cardiac fibrosis via the Keap1/Nrf2 signaling pathway

SHAOHUA XIANG¹, JING LI² and ZHENGFU ZHANG²

¹Department of Cardiothoracic Surgery, Dianjiang County Hospital of Traditional Chinese Medicine, Chongqing 408300;

²Department of Cardiothoracic Surgery, People's Hospital of Changshou, Chongqing 401220, P.R. China

Received March 7, 2019; Accepted November 5, 2019

DOI: 10.3892/etm.2020.8455

Abstract. A critical event in cardiac fibrosis is the transformation of cardiac fibroblasts (CFs) into myofibroblasts. MicroRNAs (miRNAs) have been reported to be critical regulators in the development of cardiac fibrosis. However, the underlying molecular mechanisms of action of miRNA (miR)-26b in cardiac fibrosis have not yet been extensively studied. In the present study, the expression levels of miR-26b were down-regulated in isoproterenol (ISO)-treated cardiac tissues and CFs. Moreover, miR-26b overexpression inhibited the cell viability of ISO-treated CFs and decreased the protein levels of collagen I and α -smooth muscle actin (α -SMA). Furthermore, bioinformatics analysis and dual luciferase reporter assays indicated that Kelch-like ECH-associated protein 1 (Keap1) was the target of miR-26b, and that its expression levels were decreased in miR-26b-treated cells. In addition, Keap1 overexpression reversed the inhibitory effects of miR-26b on ISO-induced cardiac fibrosis, as demonstrated by cell viability, and the upregulation of collagen I and α -SMA expression levels. Furthermore, inhibition of Keap1 expression led to the activation of nuclear factor erythroid 2-related factor 2 (Nrf2), which induced the transcriptional activation of antioxidant/detoxifying proteins in order to protect against cardiac fibrosis. Taken together, the data demonstrated that miR-26b attenuated ISO-induced cardiac fibrosis via the Keap1-mediated activation of Nrf2.

Introduction

Cardiac fibrosis is an important contributor to the development of multiple cardiovascular diseases, such as myocardial infarction and hypertension (1). It is primarily characterized by an excessive accumulation of extracellular matrix, including collagen I

and collagen III (2-5). Since myofibroblasts exhibit potent proliferative and secretory activities, the transformation of cardiac fibroblasts (CFs) to myofibroblasts is regarded as a crucial step in the development and progression of cardiac fibrosis (6-8). Despite significant advancements in the treatment of cardiac fibrosis, the underlying molecular mechanisms of cardiac fibroblast-to-myofibroblast transformation remain undiscovered.

Accumulating evidence has suggested that miRNAs are highly associated with the pathogenesis of various cardiovascular diseases, including cardiac fibrosis (9,10). Several miRNAs, such as miRNA (miR)-130, miR-328, miR-30a and miR-29a, have been reported to function as potential antifibrotic targets in cardiac fibrosis (11-14). Notably, an emerging role has been proposed for miR-26 family members in cardiovascular disease. For example, Wei *et al* (15) reported that miR-26a expression was markedly decreased in angiotensin II-induced neonatal CFs, and overexpression of miR-26a suppressed the expression of the connective tissue growth factor and collagen I. Although miR-26b has been reported to inhibit the development of hypertrophy (16), the exact role of miR-26b in the pathogenesis of cardiac fibrosis is still unclear.

Nuclear factor erythroid 2-related factor 2 (Nrf2) is a key regulator of the antioxidative response (17,18). Under normal conditions, Nrf2 remains in an inactive state and is constitutively suppressed by Kelch-like ECH-associated protein 1 (Keap1). In response to stimulation, Nrf2 dissociates from Keap1, translocates to the nucleus and activates the antioxidant response element. Increasing evidence has demonstrated that oxidative stress induces fibroblast activation and increases collagen production in various organs, leading to pathological conditions such as liver, pulmonary and cardiac fibrosis (19-21). This evidence suggests that the Keap1/Nrf2 signaling pathway is a potential candidate for antifibrotic therapy.

The present study demonstrated that miR-26b could protect heart cells from isoproterenol (ISO)-induced cardiac fibrosis via the Keap1/Nrf2 signaling pathway. The findings may provide a novel understanding of the occurrence of cardiac fibrosis and could be used for the development of a promising therapeutic modality for the treatment of this disease.

Materials and methods

Animals and treatments. The experiments involving animals were approved by the Animal Experimentation

Correspondence to: Dr Zhengfu Zhang, Department of Cardiothoracic Surgery, People's Hospital of Changshou, 16 Fengcheng Street, Chongqing 401220, P.R. China
E-mail: zhengfu_z81@163.com

Key words: cardiac fibrosis, cardiac fibroblasts, microRNA-26b, Kelch-like ECH-associated protein 1, nuclear factor erythroid 2-related factor 2

Ethics Committee of the People's Hospital of Changshou (Chongqing, China). A total of 60 adult male Sprague-Dawley rats (200-220 g) were purchased from the Laboratory of the Animal Center of the Soochow University (Suzhou, China) and randomly divided into the cardiac fibrosis model group (n=30) and the saline group (n=30). The mice were maintained under conditions of 50% relative humidity, a 12-h light/dark cycle and 22°C, and received food and water *ad libitum*. To establish the cardiac fibrosis group, ISO was injected subcutaneously for 10 days (5 mg/kg/day) (22-24). The Sprague-Dawley rats in the saline group were given equal volumes of saline. The rats were sacrificed by cervical dislocation after deep anesthesia with 2% isoflurane (Baxter International, Inc.) and their hearts were isolated for the subsequent experiments.

Cell culture and treatment. CFs were isolated from neonatal Sprague-Dawley rats. A total of eight neonatal Sprague-Dawley rats (age, 1-3 days; weight, 5-7 g) were obtained from the Laboratory of the Animal Center of the Soochow University. The ventricles of the neonatal rats were minced and digested with a mixed enzyme solution (trypsin: Collagenase I; ratio, 2:1). Subsequently, the cells were plated with DMEM (Sigma-Aldrich; Merck KGaA) containing 10% fetal bovine serum (FBS; Thermo Fisher Scientific, Inc.) for 60 min at 37°C in humidified air with 5% CO₂ until the CFs had adhered to the wall of plate. The culture plates were washed twice with PBS to detach the weakly attached and non-adherent cells and then pure CFs were obtained. The extracted cells were cultured in 90% DMEM (HyClone; GE Healthcare Life Sciences) supplemented with 10% FBS (Thermo Fisher Scientific, Inc.) in a humidified incubator at 37°C with 5% CO₂. Second- or third-generation CFs were used in these experiments.

Cell transfection. miR-26b mimics (5'-UUCAAGUAAUUC AGGAUAGGU-3'), miR-26b inhibitor (5'-ACCUAUCCU GAAUUAUCUUGAA-3') and their respective negative controls miR-NC (5'-UUCUCCGAACGUGUCACGUUU-3') and inh-miR-NC (5'-ACGUGACACGUUCGGAGAAUU-3') were synthesized by Shanghai GenePharma Co., Ltd. Keap1 cDNA (GenePharma Co., Ltd.) was cloned into pCDNA3.1 vectors (Invitrogen; Thermo Fisher Scientific, Inc.) by Shanghai GenePharma Co., Ltd. in order to construct corresponding Keap1 expression vector. The sequence of small interfering RNA (siRNA) against Keap1 (siKeap1, 5'-GCGCAAUG UUGACACGGA-3') and the sequence of the scrambled siRNA (siNC, 5'-GUACCUGACUAGUCGCAGAUU-3') were synthesized by Shanghai GenePharma Co., Ltd. CFs (5x10⁴/ml) were transfected with miR-26b mimics (20 nM) or miR-NC (20 nM), the miR-26b inhibitor (20 nM) or inh-miR-NC (20 nM) and 20 nM vectors or 50 nM siRNA. The transfection was conducted using Lipofectamine[®] 2000 transfection reagent (Invitrogen; Thermo Fisher Scientific, Inc.). At 48-h post-transfection, the subsequent experiments were conducted.

MTT assay. The cell viability of CFs was measured by MTT assay. CFs were seeded in a 96-well plate with 5x10⁵ cells/well. After transfection, the cells were treated with ISO for 24 h. Subsequently, MTT solution was added into each well and the cells were incubated for 3 h at 37°C. A total of 200 μ l DMSO

was added to solubilize the MTT crystals, and the optical density was measured at 570 nm.

Reverse transcription-quantitative (RT-q)PCR analysis. Total RNA was extracted from cardiac tissues and cells using TRIzol[®] reagent (Invitrogen; Thermo Fisher Scientific, Inc.). The RNA was reverse transcribed into cDNA using the Takara RT kit (Takara Biotechnology Co., Ltd.) at 37°C for 15 min. The expression levels of miR-26b, collagen I, α -smooth muscle actin (α -SMA), Keap1 and Nrf2 were determined using the SYBR Green PCR kit (Takara Biotechnology Co., Ltd.). GAPDH and U6 were used as controls. The following amplification conditions were used: Pre-denaturation at 95°C for 15 sec, denaturation at 94°C for 30 sec, annealing at 60°C for 20 sec, and extension at 72°C for 40 sec for 40 cycles. Relative gene expression was calculated using the 2^{- $\Delta\Delta$ Cq} method (25). The sequences of the primers were as follows: miR-26b forward, 5'-CCCAGTTCAAGTAATTCAGG-3'; and reverse, 5'-TTTGGCACTAGCACATT-3'; collagen I forward, 5'-CAG AGCACGATGTCCTGAGA-3'; and reverse, 5'-GCAAATGTG AGCTTCTGTGC-3'; α -SMA forward, 5'-GGAGTGATGGTT GGAATGG-3'; and reverse, 5'-ATGATGCCGTGTTCTATC G-3'; Keap1 forward, 5'-GGACGGCAACACTGATTC-3'; and reverse, 5'-TCGTCTCGATCTGGCTCATA-3'; Nrf2 forward, 5'-CACATCCAGACAGACACCAGT-3'; and reverse, 5'-CTA CAAATGGGAATGTCTCTGC-3'; GAPDH forward, 5'-CAA GCTCATTTCCTGGTATGAC-3'; and reverse, 5'-CAGTGA GGGTCTCTCTTCTTCT-3'; U6 forward, 5'-CTCGCTTCG GCAGCACATATACTA-3'; and reverse, 5'-ACGAATTTG CGTGTCATCCTTGCG-3'.

Bioinformatics prediction and luciferase reporter assay. The downstream target genes of miR-26b were predicted using the TargetScan database (<http://www.targetscan.org>). Keap1 was identified as a potential downstream target of miR-26b. 293T cells were obtained from the American Type Culture Collection and cultured in DMEM (Gibco; Thermo Fisher Scientific, Inc.) supplemented with 10% FBS and 1% penicillin-streptomycin (Thermo Fisher Scientific, Inc.) at 37°C in a humidified atmosphere with 5% CO₂. Luciferase reporter assays were used to investigate the regulatory relationship between miR-26b and Keap1. Site-directed mutagenesis was used to create the mutant 3'-untranslated region (UTR) sequence of Keap1. The wild-type and mutant-type sequences of Keap1 were cloned into the pGL-luciferase plasmid (Promega, Corp.), while the NC and miR-26b mimics sequences were co-transfected into 293T cells using Lipofectamine[®] 2000 transfection reagent (Invitrogen; Thermo Fisher Scientific, Inc.). After 48 h incubation at 37°C, firefly and *Renilla* luciferase activities were evaluated using the Dual-Luciferase Reporter Analysis kit (Promega Corporation). Normalized relative light units represent firefly luciferase activity/*Renilla* luciferase activity.

Western blot analysis. Cultured cells were lysed in RIPA buffer (Beyotime Institute of Biotechnology) for total protein extraction. Protein concentration was determined using a BCA assay (Beyotime Institute of Biotechnology). A total of 10 μ g protein/lane were separated by SDS-PAGE (10% gel) and subsequently transferred to nitrocellulose membranes. Following blocking with 5% bovine serum albumin (Beijing




Figure 1. Upregulation of collagen I and α -SMA in ISO-treated cardiac tissues. (A) RT-qPCR analysis indicated that the expression levels of collagen I and α -SMA were increased in ISO-treated rat cardiac tissues compared with those noted in the saline group. (B) Western blot analysis indicated that the protein levels of collagen I and α -SMA were increased in ISO-treated rat cardiac tissues compared with those noted in the saline group. (C) RT-qPCR analysis indicated that the expression levels of collagen I and α -SMA were increased in ISO-treated CFs compared with those noted in the saline group. (D) Western blot analysis indicated that the protein levels of collagen I and α -SMA were increased in ISO-treated CFs compared with those noted in the saline group. ^{**}P<0.01, ^{##}P<0.01 vs. saline. α -SMA, α -smooth muscle actin; ISO, isoproterenol; RT-qPCR, reverse transcription-quantitative PCR; CF, cardiac fibroblast; Col1A1, collagen I.

Solarbio Science and Technology Co., Ltd.) for 1 h at room temperature, the proteins were incubated with primary antibodies, namely anti-collagen I (cat. no. ab34710), anti- α -SMA (cat. no. ab5694) and anti- β -actin (cat. no. ab8227) (all 1:1,000; all from Abcam), at 4°C overnight. Following washing, the membranes were incubated with horseradish peroxidase-conjugated secondary antibodies (goat anti-mouse IgG; cat. no. ab205719; and goat anti-rabbit IgG; cat. no. ab205718) (both 1:1,000; both from Abcam) for 2 h at room temperature. The proteins were detected using an enhanced chemiluminescence detection system (ProteinSimple) and analyzed using Image-Pro[®] Plus software (version 6.0; Media Cybernetics, Inc.).

Statistical analysis. Statistical analyses were performed using SPSS software (version 18.0; SPSS, Inc.). All data are presented as the mean \pm standard deviation of at least three independent experiments. The group differences were determined by either Student's t-test or one-way ANOVA. Multiple comparisons between the groups was performed using the Student-Newman-Keuls method. P<0.05 was considered to indicate a statistically significant difference.

Results

Upregulation of collagen I and α -SMA in ISO-treated cardiac tissues. ISO has been reported to increase the expression of pro-inflammatory cytokines [interleukin (IL)-1, IL-6 and IL-17] (26). The release of cytokines contributes to cardiac fibrosis via upregulation of matrix metalloproteinase expression in CFs (26). In the present study, Sprague-Dawley rats were treated with ISO to induce cardiac fibrosis. Subsequently, RT-qPCR was used to evaluate the mRNA expression levels of collagen I and α -SMA in cardiac tissues. The mRNA expression levels of collagen I and α -SMA were increased in ISO-treated rats compared with those of the untreated rats (Fig. 1A). Furthermore, western blot analysis indicated that the protein expression levels of collagen I and α -SMA were increased in ISO-treated rat cardiac tissues compared with those noted in the saline group (Fig. 1B). In addition, ISO was used to treat CFs to induce cardiac fibrosis. RT-qPCR and western blot analysis demonstrated that the expression levels of collagen I and α -SMA were upregulated in ISO-treated CFs compared with those of the control groups (Fig. 1C and D). In summary, the results indicated that




Figure 2. Expression of miR-26b and Keap1 in ISO-treated cardiac tissues and CFs. (A) RT-qPCR showed the relative expression of miR-26b in cardiac fibrosis tissues and normal tissues. (B) Western blotting showed the relative expression of Keap1 in cardiac fibrosis tissues and normal tissues. (C) RT-qPCR showed the relative expression of miR-26b in ISO-treated neonatal rat CFs and control group. (D) Western blotting showed the relative expression of Keap1 in ISO-treated neonatal rat CFs and control group. * $P < 0.05$, ** $P < 0.01$ vs. respective saline group. ISO, isoproterenol; CF, cardiac fibroblast; RT-qPCR, reverse transcription-quantitative PCR; Keap1, Kelch-like ECH-associated protein 1; miR, microRNA.

collagen I and α -SMA were overexpressed in the ISO-treated cardiac fibrosis model.

Expression of miR-26b and Keap1 in ISO-treated cardiac tissue and CFs. The expression of miR-26b and Keap1 in ISO-treated cardiac tissues and normal tissues was determined by RT-qPCR. As shown in Fig. 2A and B, the expression of miR-26b was downregulated in ISO-treated cardiac tissues compared with saline-treated group, while the expression of Keap1 was upregulated in ISO-treated cardiac tissues. Furthermore, the expression of miR-26b and Keap1 was analyzed *in vitro* (Fig. 2C and D). Consistent with the results *in vivo*, the expression level of miR-26b was also decreased and the expression level of Keap1 was increased in ISO-treated CFs compared with the saline group.

miR-26b suppresses the cell viability of CFs. To investigate the precise role of miR-26b in ISO-induced cardiac fibrosis, ISO-treated CFs were transfected with miR-26b mimics to overexpress miR-26b, and with the miR-26b inhibitor to decrease miR-26b expression. The transfection efficiency was confirmed by RT-qPCR (Fig. 3A). miR-26b overexpression significantly inhibited the cell viability of ISO-treated CFs, while miR-26b inhibition enhanced the growth of ISO-treated CFs (Fig. 3B). Furthermore, western blot analysis indicated that miR-26b overexpression decreased the protein levels of

collagen I and α -SMA compared with those of the miR-NC group. In contrast to these findings, miR-26b inhibition caused an increase in the expression levels of collagen I and α -SMA proteins (Fig. 3C). Taken collectively, the data indicated that miR-26b inhibited the transformation of CFs into myofibroblasts.

Keap1 is a target of miR-26b. The analysis using the bioinformatics prediction database TargetScan demonstrated that Keap1 was a potential target of miR-26b (Fig. 4A). To determine whether miR-26b could directly interact with Keap1, a dual luciferase reporter assay was performed. The results indicated that miR-26b mimics significantly reduced the luciferase activity of wild-type Keap1, whereas no effect was noted in the cells with the mutant 3'-UTR of Keap1 (Fig. 4B). Moreover, the data indicated that miR-26b mimics significantly reduced Keap1 expression, while the miR-26b inhibitor increased the expression levels of Keap1 (Fig. 4C and D). These results demonstrated that miR-26b could directly target Keap1 to inhibit its expression.

miR-26b-mediated inhibitory effects on ISO-induced cardiac fibrosis are abrogated by Keap1. To examine whether the effect of miR-26b on ISO-induced cardiac fibrosis is regulated by Keap1, two cell groups of ISO-treated CFs were prepared, one that was transfected with miR-NC, miR-26b




Figure 3. miR-26b suppresses the cell viability of CFs. (A) Reverse transcription-quantitative PCR analysis indicating the relative expression level of miR-26b in ISO-treated CFs transfected with miR-26b mimics and the miR-26b inhibitor. (B) MTT assay results showing the cell viability of ISO-treated CFs transfected with miR-26b mimics and the miR-26b inhibitor. (C) Western blot analysis showing the protein levels of collagen I and α -SMA in ISO-treated CFs transfected with miR-26b mimics and the miR-26b inhibitor. * $P < 0.05$, ** $P < 0.01$ vs. respective control. miR, microRNA; inh, inhibitor; ISO, isoproterenol; CF, cardiac fibroblast; NC, negative control; Col1A1, collagen I; α -SMA, α -smooth muscle actin.




Figure 4. Keap1 is one of the targets of miR-26b. (A) Bioinformatics prediction of the binding site for miR-26b in Keap1. (B) The luciferase reporter assay confirmed that miR-26b was able to bind to Wt and not Mut Keap1 in 293T cells. (C) RT-qPCR analysis showed the expression levels of miR-26b and Keap1 in ISO-treated CFs (NC and miR-26b mimics). (D) RT-qPCR analysis showed the expression levels of miR-26b and Keap1 in ISO-treated CFs (NC and miR-26b inhibitor). * $P < 0.05$ vs. respective control. Mut, mutant; Wt, wild-type; NC, negative control; miR, microRNA; Keap1, Kelch-like ECH-associated protein 1; inh, inhibitor; RT-qPCR, reverse transcription-quantitative PCR; CF, cardiac fibroblast; ISO, isoproterenol; NC, negative control.




Figure 5. Restoration of Keap1 abolishes the inhibitory effect of miR-26b on ISO-treated CFs. (A) Western blot analysis showing the relative expression levels of Keap1 in ISO-treated CFs transfected with miR-NC, miR-26b, miR-26b+pcDNA and miR-26b+Keap1. (B) Western blot analysis showing the relative expression levels of Keap1 in ISO-treated CFs transfected with inh-miR-NC, inh-miR-26b, inh-miR-26b+siNC and inh-miR-26b+siKeap1. (C) Cell viability of ISO-treated CFs transfected with miR-NC, miR-26b, miR-26b+pcDNA and miR-26b+Keap1, as determined by MTT assay. (D) Cell viability of ISO-treated CFs transfected with inh-miR-NC, inh-miR-26b, inh-miR-26b+siNC and inh-miR-26b+siKeap1 as determined by MTT assay. (E) The protein levels of collagen I and α -SMA were detected in ISO-treated CFs transfected with miR-NC, miR-26b, miR-26b+pcDNA and miR-26b+Keap1 by western blotting. (F) The protein levels of collagen I and α -SMA were detected by western blotting in ISO-treated CFs transfected with inh-miR-NC, inh-miR-26b, inh-miR-26b+siNC and inh-miR-26b+siKeap1. * P <0.05. Keap1, Kelch-like ECH-associated protein 1; NC, negative control; α -SMA, α -smooth muscle actin inh, inhibitor; ISO, isoproterenol; miR, microRNA; CF, cardiac fibroblast; si, small interfering; Col1A1, collagen I.

mimics, miR-26b+pcDNA or miR-26b mimics+Keap1, and another that was transfected with inh-miR-NC, miR-26b inhibitor, inh-miR-26b+siNC or inh-miR-26b+siKeap1 (Fig. 5A and B). The data indicated that Keap1 overexpression significantly reduced the inhibitory effect of miR-26b on cell viability (Fig. 5C). Conversely, Keap1 silencing attenuated the cell growth of CFs transfected with the miR-26b inhibitor (Fig. 5D). Moreover, western blot analysis demonstrated that miR-26b mimics decreased the protein levels of collagen I and α -SMA in ISO-treated CFs, indicating that miR-26b inhibited ISO-induced cardiac fibrosis; these effects were abrogated by Keap1 overexpression (Fig. 5E). The opposite effect was noted in the ISO-treated CFs co-transfected with inh-miR-26b and siKeap1 (Fig. 5F). In summary, the results demonstrated that Keap1 restoration or silencing abrogated the effects caused by miR-26b mimics or the miR-26b inhibitor in ISO-treated CFs.

miR-26b inhibits ISO-induced cardiac fibrosis via the Keap1/Nrf2 signaling pathway. Previous studies have shown that Nrf2 plays an important role in myofibroblast

dedifferentiation and its activity is under the control of Keap1 (27). Therefore, it was assumed that miR-26b and Keap1 may exert their roles in cardiac fibrosis via activation of Nrf2. The data demonstrated that the upregulation of miR-26b increased the expression levels of Nrf2 in ISO-treated CFs, while this effect was abolished by Keap1 overexpression (Fig. 6A). In contrast to these findings, treatment of the cells with the miR-26b inhibitor caused a downregulation in the expression levels of Nrf2, and Keap1 silencing alleviated the inhibitory effect of miR-26b inhibitor on the expression of Nrf2 (Fig. 6B). In summary, these data demonstrated that miR-26b caused an upregulation of Nrf2 expression levels by targeting Keap1 in ISO-treated CFs.

Discussion

Abnormal activation and cell viability of CFs is a pivotal step in the occurrence and development of cardiac fibrosis. Therefore, identifying approaches that inhibit CF activation may be an effective therapeutic strategy to prevent cardiac fibrosis. The present study demonstrated that miR-26b




Figure 6. miR-26b targets Keap1 to activate Nrf2. (A) Western blot analysis showing the relative expression levels of Nrf2 in ISO-treated CFs transfected with miR-NC, miR-26b, miR-26b+pcDNA and miR-26b+Keap1. (B) Western blot analysis showing the relative expression levels of Nrf2 in ISO-treated CFs transfected with inh-miR-NC, inh-miR-26b, inh-miR-26b+siNC and inh-miR-26b+siKeap1. * $P < 0.05$. miR, microRNA; Keap1, Kelch-like ECH-associated protein 1; NC, negative control; Nrf2, nuclear factor erythroid 2-related factor 2; inh, inhibitor; si, small interfering; ISO, isoproterenol.

suppressed the activation and cell viability of ISO-treated CFs by upregulating Nrf2 expression and targeting Keap1.

It has been thoroughly documented that miRNAs are involved in the development and progression of cardiac fibrosis. For example, Zhou *et al* (28) reported that miR-21 could promote cardiac fibroblast-to-myofibroblast transition by downregulating the expression of Jagged1. Moreover, several studies have indicated that the miR-26 family plays a central role in the development of cardiovascular disease by controlling critical signaling pathways or downstream gene targets. For example, miR-26a inhibits pathological and physiological angiogenesis by targeting bone morphogenic protein/SMAD1 signaling in endothelial cells (29). Han *et al* (16) reported that miR-26b inhibited the development of cardiac hypertrophy by inhibiting the expression of GATA binding protein 4. However, the functional role of miR-26b, which is a member of the miR-26 family, has not been extensively studied. The present study demonstrated that miR-26b was significantly downregulated in ISO-treated cardiac tissues and CFs. Furthermore, we investigated the effects of miR-26b on ISO-treated CFs. The results indicated that miR-26b mimics inhibited cell viability and decreased the protein levels of collagen I and α -SMA, while the miR-26b inhibitor increased the cell growth rate and the expression levels of collagen I and the α -SMA proteins. The data suggested that miR-26b exerts a suppressive role on ISO-induced cardiac fibrosis.

The induction of oxidative stress contributes to the occurrence and development of cardiac fibrosis. The Nrf2-antioxidant response element signaling pathway acts as a critical cellular

defense mechanism that antagonizes oxidative stress (30,31). Keap1 is an important regulator and repressor of this signaling pathway (30,31). For example, Civantos *et al* (32) reported that sitagliptin ameliorated oxidative stress in diabetic kidney rat tissues through downregulation of miR-200a, which further regulated the Keap1/Nrf2 signaling pathway. Moreover, Yang *et al* (33) reported that Nrf2 could protect against hepatic stellate fibrosis by functionally activating the transcription of antioxidant response genes. In the present study, bioinformatics analysis and luciferase reporter assays demonstrated that Keap1 could directly interact with miR-26b, and that its overexpression abrogated the inhibitory effect of miR-26b on ISO-treated CFs, as determined by increased cell viability and expression of fibrosis-related indices in ISO-treated CFs. Moreover, it was demonstrated that the targeting of Keap1 by miR-26b caused an upregulation of Nrf2 expression. Therefore, the data suggested that miR-26b could regulate the Keap1/Nrf2 signaling pathway to inhibit ISO-induced cardiac fibrosis.

Overall, the present study demonstrated for the first time, to the best of our knowledge, that miR-26b could attenuate ISO-induced cardiac fibrosis by activating Nrf2 and by interacting with Keap1. These findings may provide additional evidence for the development of a novel therapeutic strategy for cardiac fibrosis and for understanding the pathogenesis of this disease.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The datasets used and/or analyzed during the present study are available from the corresponding author upon reasonable request.

Authors' contributions

SX and ZZ designed the present study. ZZ collected the samples, and ZZ and JL performed all the experiments. SX and JL analyzed the data and prepared the figures. SX drafted the initial manuscript. ZZ reviewed and revised the initial manuscript. All authors approved the final published version of this manuscript.

Ethics approval and consent to participate

The present study was approved by the Ethics Committee of the People's Hospital of Changshou. The experiments involving animals were approved by the Animal Experimentation Ethics Committee of the People's Hospital of Changshou.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Molkentin JD, Bugg D, Ghearing N, Dorn LE, Kim P, Sargent MA, Gunaje J, Otsu K and Davis J: Fibroblast-specific genetic manipulation of p38 mitogen-activated protein kinase in vivo reveals its central regulatory role in fibrosis. *Circulation* 136: 549-561, 2017.
- Segura AM, Frazier OH and Buja LM: Fibrosis and heart failure. *Heart Fail Rev* 19: 173-185, 2014.
- Edgley AJ, Krum H and Kelly DJ: Targeting fibrosis for the treatment of heart failure: A role for transforming growth factor- β . *Cardiovasc Ther* 30: e30-e40, 2012.
- Espira L and Czubyrt MP: Emerging concepts in cardiac matrix biology. *Can J Physiol Pharmacol* 87: 996-1008, 2009.
- Fan D, Takawale A, Lee J and Kassiri Z: Cardiac fibroblasts, fibrosis and extracellular matrix remodeling in heart disease. *Fibrogenesis Tissue Repair* 5: 15, 2012.
- Barnes JL and Gorin Y: Myofibroblast differentiation during fibrosis: Role of NAD(P)H oxidases. *Kidney Int* 79: 944-956, 2011.
- Weber KT, Sun Y, Tyagi SC and Cleutjens JP: Collagen network of the myocardium: Function, structural remodeling and regulatory mechanisms. *J Mol Cell Cardiol* 26: 279-292, 1994.
- Swynghedauw B: Molecular mechanisms of myocardial remodeling. *Physiol Rev* 79: 215-262, 1999.
- Romaine SP, Tomaszewski M, Condorelli G and Samani NJ: MicroRNAs in cardiovascular disease: An introduction for clinicians. *Heart* 101: 921-928, 2015.
- Gurha P: MicroRNAs in cardiovascular disease. *Curr Opin Cardiol* 31: 249-254, 2016.
- Li L, Bounds KR, Chatterjee P and Gupta S: MicroRNA-130a, a potential antifibrotic target in cardiac fibrosis. *J Am Heart Assoc* 6: pii: e006763, 2017.
- Du W, Liang H, Gao X, Li X, Zhang Y, Pan Z, Li C, Wang Y, Liu Y, Yuan W, *et al*: MicroRNA-328, a potential anti-fibrotic target in cardiac interstitial fibrosis. *Cell Physiol Biochem* 39: 827-836, 2016.
- Chen L, Ji Q, Zhu H, Ren Y, Fan Z and Tian N: miR-30a attenuates cardiac fibrosis in rats with myocardial infarction by inhibiting CTGF. *Exp Ther Med* 15: 4318-4324, 2018.
- Qin RH, Tao H, Ni SH, Shi P, Dai C and Shi KH: microRNA-29a inhibits cardiac fibrosis in Sprague-Dawley rats by downregulating the expression of DNMT3A. *Anatol J Cardiol* 20: 198-205, 2018.
- Wei C, Kim IK, Kumar S, Jayasinghe S, Hong N, Castoldi G, Catalucci D, Jones WK and Gupta S: NF- κ B mediated miR-26a regulation in cardiac fibrosis. *J Cell Physiol* 228: 1433-1442, 2013.
- Han M, Yang Z, Sayed D, He M, Gao S, Lin L, Yoon S and Abdellatif M: GATA4 expression is primarily regulated via a miR-26b-dependent post-transcriptional mechanism during cardiac hypertrophy. *Cardiovasc Res* 93: 645-654, 2012.
- Niture SK, Khatri R and Jaiswal AK: Regulation of Nrf2-an update. *Free Radic Biol Med* 66: 36-44, 2014.
- Baird L and Dinkova-Kostova AT: The cytoprotective role of the Keap1-Nrf2 pathway. *Arch Toxicol* 85: 241-272, 2011.
- Shi H, Shi A, Dong L, Lu X, Wang Y, Zhao J, Dai F and Guo X: Chlorogenic acid protects against liver fibrosis in vivo and in vitro through inhibition of oxidative stress. *Clin Nutr* 35: 1366-1373, 2016.
- Cheresh P, Kim SJ, Tulasiram S and Kamp DW: Oxidative stress and pulmonary fibrosis. *Biochim Biophys Acta* 1832: 1028-1040, 2013.
- Wang LP, Fan SJ, Li SM, Wang XJ, Gao JL and Yang XH: Oxidative stress promotes myocardial fibrosis by upregulating KCa3.1 channel expression in AGT-REN double transgenic hypertensive mice. *Pflugers Arch* 469: 1061-1071, 2017.
- Wu Y, Liu Y, Pan Y, Lu C, Xu H, Wang X, Liu T, Feng K and Tang Y: MicroRNA-135a inhibits cardiac fibrosis induced by isoproterenol via TRPM7 channel. *Biomed Pharmacother* 104: 252-260, 2018.
- Tao H, Zhang JG, Qin RH, Dai C, Shi P, Yang JJ, Deng ZY and Shi KH: LncRNA GAS5 controls cardiac fibroblast activation and fibrosis by targeting miR-21 via PTEN/MMP-2 signaling pathway. *Toxicology* 386: 11-18, 2017.
- Lu J, Wang QY, Zhou Y, Lu XC, Liu YH, Wu Y, Guo Q, Ma YT and Tang YQ: Astragaloside IV against cardiac fibrosis by inhibiting TRPM7 channel. *Phytomedicine* 30: 10-17, 2017.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Periasamy S, Chen SY and Liu MY: The study of ISO induced heart failure rat model. *Exp Mol Pathol*. 2010;88:299-304. *Exp Mol Pathol* 90: 84, 2011.
- Artaud-Macari E, Goven D, Brayer S, Hamimi A, Besnard V, Marchal-Somme J, Ali ZE, Crestani B, Kerdine-Römer S, Boutten A and Bonay M: Nuclear factor erythroid 2-related factor 2 nuclear translocation induces myofibroblastic dedifferentiation in idiopathic pulmonary fibrosis. *Antioxid Redox Signal* 18: 66-79, 2013.
- Zhou XL, Xu H, Liu ZB, Wu QC, Zhu RR and Liu JC: miR-21 promotes cardiac fibroblast-to-myofibroblast transformation and myocardial fibrosis by targeting Jagged1. *J Cell Mol Med*: May 28, 2018 (Epub ahead of print).
- Icli B, Wara AK, Moslehi J, Sun X, Plovie E, Cahill M, Marchini JF, Schissler A, Padera RF, Shi J, *et al*: MicroRNA-26a regulates pathological and physiological angiogenesis by targeting BMP/SMAD1 signaling. *Circ Res* 113: 1231-1241, 2013.
- Köhler UA, Kurinna S, Schwitter D, Marti A, Schäfer M, Hellerbrand C, Speicher T and Werner S: Activated Nrf2 impairs liver regeneration in mice by activation of genes involved in cell-cycle control and apoptosis. *Hepatology* 60: 670-678, 2014.
- Gambhir L, Checker R, Thoh M, Patwardhan RS, Sharma D, Kumar M and Sandur SK: 1,4-Naphthoquinone, a pro-oxidant, suppresses immune responses via KEAP-1 glutathionylation. *Biochem Pharmacol* 88: 95-105, 2014.
- Civantos E, Bosch E, Ramirez E, Zhenyukh O, Egido J, Lorenzo O and Mas S: Sitagliptin ameliorates oxidative stress in experimental diabetic nephropathy by diminishing the miR-200a/Keap-1/Nrf2 antioxidant pathway. *Diabetes Metab Syndr Obes* 10: 207-222, 2017.
- Yang JJ, Tao H, Hu W, Liu LP, Shi KH, Deng ZY and Li J: MicroRNA-200a controls Nrf2 activation by target Keap1 in hepatic stellate cell proliferation and fibrosis. *Cell Signal* 26: 2381-2389, 2014.

