

Associations Between Gait Speed and Fat Mass in Older Adults

Peter Sagat

GSD/Health and Physical Education Department, Sport Sciences and Diagnostics Research Group, Prince Sultan University, Riyadh, 11586, Saudi Arabia

Correspondence: Peter Sagat, Email sagat@psu.edu.sa

Purpose: Although both gait speed and fat mass are crucial for healthy aging, evidence suggests that the associations between these components remain unclear. Therefore, the main purpose of the study was to examine the associations between gait speed and fat mass.

Patients and Methods: In this cross-sectional study, we recruited 643 older men and women aged >60 years. Fat mass was assessed using bioelectrical impedance analysis, while gait speed was determined by calculating the time an individual has taken to walk across a 4.6-m distance. Receiver operating characteristic (ROC) curves and odds ratios (OR) were performed to determine cut-off points and mutual associations.

Results: In older men, the optimal threshold of gait speed to detect high level of fat mass was 1.40 m/s with the area under the curve (AUC) being 0.82 (95% CI 0.76–0.89, $p < 0.001$). In older women, the optimal cut-off point was 1.37 m/s (AUC = 0.85, 95% CI 0.81–0.90, $p < 0.001$). Older men and women who walked below the newly developed threshold were approximately 12 times more likely to have high level of fat.

Conclusion: In summary, newly developed cut-off points of gait speed have adequate discriminatory ability to detect older men and women with high level of fat mass. Although gait speed may be considered as a satisfactory screening tool for fat mass, its utility in clinical practice needs to be further investigated.

Keywords: body composition, physical performance, aging, cut-points, associations

Introduction

In the last 50 years, the number of older adults has dramatically increased by 2%, and the projection estimates that the number will increase to 22% by 2050.¹ Evidence suggests that within the next 20 years, the number of older adults will outnumber younger generations worldwide.² Aging process is accompanied by many health-related consequences, including limitations and disabilities that are more prevalent in this population, compared to younger individuals.³ The co-existing effect of aging and unwanted body composition changes may even accelerate the aging process,⁴ leading to an increased risk of all-cause mortality.⁵

Body composition changes, which are described as a shifting process from lean muscle mass to fat mass, are particularly intensified during the aging process.⁶ Thus, it is not surprising that approximately one-third of older adults are classified as being those with obesity, with an increasing trend being observed in the future.⁷ Indeed, higher level of fat mass has been associated with a variety of metabolic measures, including an unproductive weight gain,⁸ loss of strength,⁹ and higher risk of sarcopenic obesity.¹⁰

Along with body composition, previous studies have shown that poor physical performance has been associated with aging, especially a decline in gait speed in the sixth decade.¹¹ From a practical point of view, gait speed is often used for functional assessment as it is a reliable and easy-to-administrate measurement.^{12,13} The applicability of gait speed has been well established in longitudinal studies, as it may predict survival in older adults.¹⁴ Even though a handgrip strength test is more commonly used to assess the functional status in older adults,¹⁵ due to a more prevalent upper limb

deformities occurring in this age group,¹⁶ gait speed is often considered an appropriate and easy-to-administrate measure of physical performance.¹⁵

Previous cross-sectional^{17–19} and longitudinal²⁰ studies have shown that body composition, ie, body-mass index or fat mass, is associated with gait speed in older adults. In general, the findings suggest that higher level of accumulated fat is related to slower gait speed.^{18–20} Although a gait speed of ≤ 0.8 m/s is often used as a “proxy” of mobility impairment and reduced overall survival,²¹ the threshold of gait speed to identify older adults with high level of fat mass is still unknown. In previous studies, very few older adults have exhibited gait speed values below 0.8 m/s, possibly limiting the significance in clinical practice.^{17–20,22} By detecting an optimal cut-off point of gait speed to define high level of fat mass, one would be able to distinguish between healthy and unhealthy individuals. Also, the newly proposed cut-off values of gait speed may serve as a quick screening tool to detect older adults with high fat mass level.

Therefore, the main purpose of the study was to establish optimal cut-off points of gait speed to predict high level of fat mass in older adults. Based on previous findings,^{17–20} we hypothesized that gait speed would give a relatively high diagnostic accuracy to identify those individuals, who might have high levels of fat mass.

Materials and Methods

Study Participants and Design

For the purpose of this observational prospective cross-sectional study, we recruited 843 older men and women using services of a single rehabilitation center in the “Lipik” county near the city of Zagreb from 2020 to 2022. This study was part of an internal project aiming to determine sociodemographic, physical, mental, and social factors of healthy aging through annual examinations in one of their rehabilitation centers. All participants were asked about the number of days during the week they spent in moderate-to-vigorous physical activity and were classified as “active”, “not active enough”, and “inactive”. Participants included in initial analyses were apparently healthy older adults, who were considered physically active, based on a quick physical activity assessment prior the study. Of note, most of the participants included in this study (87%) attended Pilates, yoga, or corrective gymnastics classes organized for older adults approximately 2x per week. Socioeconomic status was part of the questionnaire and the participants needed to fulfill the question regarding their socioeconomic status in comparison to their sex- and age-specific peers with one-item question: “How would you rate your socioeconomic status, compared to your peers?” with the answers arranged along a 5-item Likert scale: i) very above average, ii) above average, iii) average, iv) below average, and v) very below average. Most of the participants were classified into the “average” socioeconomic group. Socioeconomic status was not significantly correlated to the level of physical activity ($r = 0.19$) and training class attendances ($r = 0.11$); therefore, socioeconomic status was excluded as a barrier towards being physically active. Individuals included in the study should be without chronic diseases, which included cardiovascular, locomotor, or neural complaints. After cleaning the data, 643 men ($N=263$, mean age: 67.4 ± 5.5 years) and women ($N=380$, 66.9 ± 5.2 years) met the inclusion criteria and were included in the final analysis. Sex and age were self-reported. Before initial measurements, all participants had been given information regarding general aims and hypotheses of the project. The participants were ensured confidentiality and informed that their participation was voluntary, and that they had the right to withdraw at any time. Data collection was done in the same setting by the same experimental team, minimizing the effect of a measurement error. All participants have read and signed the informed consent forms. The research related to human use has been complied with all the relevant national regulations, institutional policies and in accordance with the tenets of the Helsinki Declaration, and has been approved by the Ethical Committee of the Home of War veterans (Ethical code: 2022/4).

Fat Mass

The level of fat mass was assessed using bioelectrical impedance analysis (Omron BF500 Body Composition Monitor, Omron Medizintechnik, Vernon Hills, IL, USA). In brief, each participant stood barefoot and with light clothes on metal footpads holding a pair of electrodes attached to the device in front of the chest.²³ After a few seconds, the device used a pre-programmed equation based on sex, height, and weight to estimate the level of fat mass. Following the

recommendations of the American College of Sports Medicine,^{24,25} cut-off points for obesity were 28% and 35% for men and women, compared to their counterparts with obesity (<28% for men and <35% for women).

Gait Speed

As a proxy of physical performance, we determined gait speed by calculating the time an individual has taken to walk across a 4.6-m distance.^{26,27} Specifically, the final time of the walk was obtained by dividing the distance of 4.6-m by the time to cover the same distance. For better transparency, a visible tape was placed from the starting to the finishing position. Each participant was instructed to walk at the usual pace for a distance of 4.6-m, while a stopwatch operated by the measurer recorded time to the nearest 0.01 s. To control for acceleration (a sudden movement from standing to walking) and deceleration (from walking at constant pace to slowing down),^{26,27} an additional distance of 2 m before and after the 4.6-m distance was incorporated.

Data Analysis

Basic descriptive statistics are presented as means and standard deviations (SD) for normally distributed variables and as percentages (%) for categorical variables. Sex differences were examined with Student's *t*-test for independent samples and Chi-square test. The effect size of the comparisons between the sexes was calculated using Cohen's *D* effect size (ES). ES was classified as trivial (<0.2), small (0.2–0.6), moderate (0.6–1.2), large (1.2–2.0), very large (>2.0) and extremely large (>4.0).²⁸ To identify diagnostic accuracy of gait speed to predict high level of fat mass, we performed a set of receiver operating characteristics (ROC) curves quantified by the area under the curve (AUC). ROC curves analyses were used to determine a discriminatory power of a specific diagnostic test, where the curve of the test skews closer to the upper left corner.²⁹ To be able to correctly classify the accuracy of the test, Rice and Harris³⁰ proposed a classification of the AUC as follows: 1) 0.55–0.62 (small), 2) 0.63–0.71 (moderate), and 3) >0.71 (large). Sensitivity and specificity characteristics were calculated and presented as percentages (%). When the proposed cut-off values for gait speed were determined, we used odds ratios (ORs) with 95% confidence intervals (95% CI) to examine binary associations with higher level of fat mass. Finally, the correlation between newly-developed gait speed cut-off points in this study (slow vs fast) and fat mass (those without obesity vs those with obesity) proposed by the American College of Sports Medicine^{24,25} was calculated using the Kappa coefficient (κ). Two-sided *p*-values were used, and the significance was set at $\alpha < 0.05$. All the analyses were calculated in Statistical Packages for Social Sciences v.23 (SPSS, Chicago, IL, United States).

Results

Basic descriptive statistics of the study participants are presented in Table 1. Older men were taller, heavier, and had lower heart rate values, compared to women. Older women had higher fat mass, compared to older men. Interestingly, both older men and women exhibited similar values in gait speed.

Table 1 Basic Descriptive Statistics of the Study Participants, Stratified by Sex

Study Variables	Total Sample (N=643)	Men (N=263)	Women (N=380)	Cohen <i>d</i>	p-value
	Mean (SD)	Mean (SD)	Mean (SD)		
Age (years)	67.4 (5.4)	67.8 (5.7)	67.2 (5.2)	0.11	0.295
Height (cm)	165.7 (8.6)	173.5 (6.2)	160.9 (6.0)	2.07	< 0.001
Weight (kg)	75.1 (13.3)	83.3 (10.7)	70.0 (12.2)	1.24	< 0.001
Body-mass index (kg/m ²)	27.1 (3.9)	27.4 (3.4)	26.9 (4.2)	0.15	0.201
Heart rate (beats/min)	67.3 (9.9)	64.9 (10.2)	68.8 (9.4)	0.38	< 0.001
Gait speed (m/s)	1.35 (0.21)	1.33 (0.22)	1.36 (0.20)	0.14	0.134
Fat mass (%)	34.5 (7.6)	30.8 (6.9)	36.7 (7.2)	0.86	< 0.001
Lean mass (%)	65.5 (7.6)	69.2 (6.9)	63.3 (7.2)	0.86	< 0.001

Note: *P* < 0.05.




Figure 1 The ROC of gait speed to identify older men (A) and older women (B) with high level of fat mass.
Notes: Sensitivity: denotes a test's ability to designate an individual with condition as positive; Specificity: denotes a test's ability to designate an individual without condition as negative; Cut-off points of $\geq 28\%$ and $\geq 35\%$ denote individuals with obesity for older men and women.^{24,25}

Figure 1 shows ROC curves of gait speed to identify older men and women with high level of fat mass. A diagnostic assessment for both sexes is presented in Table 2. For both older men and women, the AUCs were >0.80 . Older men with low gait speed (<1.40 m/s) determined by the ROC were 12.66 (OR=12.66, 95% CI 5.75 to 27.86, $p<0.001$) more likely to have high level of fat mass ($\geq 28\%$). In older women, low gait speed (<1.37 m/s) was associated with 12.09 (OR=12.09, 95% CI 6.75 to 21.65, $p<0.001$) higher likelihood of having high level of fat mass ($\geq 35\%$).

True positive (TP), true negative (TN), false positive (FP), false negative (FN) and Kappa statistics are presented in Table 3. In older men, TP and TN values were 72.5% and 82.7%, while FP and FN were 17.3% and 27.5%. In older

Table 2 Receiver Operating Curve Cut-Offs for Gait Speed to Predict High Level of Fat Mass, Stratified by Sex

Study Variable	High Fat Mass (Men $\geq 28\%$; Women $\geq 35\%$)				
	AUC	95% CI	Std. Error	p-value	Cut-Off Point
Old men (N=260)	0.82	0.76–0.89	0.034	< 0.001	1.40 m/s
Old women (N=383)	0.85	0.81–0.90	0.023	< 0.001	1.37 m/s

Note: $P < 0.05$.
Abbreviations: AUC, Area under the curve; 95% CI, 95% confidence interval; Std. Error, Standard Error.

Table 3 Sensitivity, Specificity and Kappa Statistics for Gait Speed Cut-Offs and High Fat Mass, Stratified by Sex

Old Men (N = 263)	High Fat Mass (Men $\geq 28\%$; Women $\geq 35\%$)					
	Low	High	Chi-Square Test	p-value	Kappa	p-value
Gait Speed (m/s)						
≥ 1.40 m/s	72.5%	17.3%	46.934	< 0.001	-0.43	< 0.001
< 1.40 m/s	27.5%	82.7%				
Old women (N = 380)						
Gait speed (m/s)						
≥ 1.37 m/s	77.8%	22.5%	80.638	< 0.001	-0.53	< 0.001
< 1.37 m/s	22.2%	77.5%				

Notes: True positive (TP): denotes "low" fat mass and gait speed equal or above the threshold in men (≥ 1.40 m/s) and women (≥ 1.37 m/s); True negative (TN): denotes "high" fat mass and gait speed below the threshold in men (< 1.40 m/s) and women (< 1.37 m/s); False positive (FP): denotes "high" fat mass and gait speed equal or above the threshold in men (≥ 1.40 m/s) and women (≥ 1.37 m/s); False negative (FN) denotes "low" fat mass and gait speed below the threshold in men (< 1.40 m/s) and women (< 1.37 m/s) $P < 0.05$.

women, TP and TN values were 77.8% and 77.5%, while FP and FN were 22.5% and 22.2%. Kappa statistics showed moderate correlations between gait speed and fat mass for both sexes.

Discussion

The main purpose of the study was to examine whether gait speed had a sufficient discriminatory power for detecting older individuals with high level of fat mass.

The main findings of the study are as follows: i) gait speed accurately predicts high level of fat mass in both older men and women; ii) the optimal cut-off points of gait speed are 1.40 m/s and 1.37 m/s for older men and women; and iii) strong sensitivity and specificity properties and moderate inverse correlations between gait speed and fat mass are observed in both sexes. A well-documented body of evidence has shown that physical performance assessed through gait speed is associated with activity of daily living and mobility disability in non-disabled, community-dwelling older people.^{31,32} The findings of these studies suggest that slow gait speed (arbitrarily considered as <1 m/s) increases the risk of negative health-related outcomes, including persistent lower extremity limitations,^{33,34} incident activity of daily living disabilities and limitations^{35–37} and all-cause mortality.^{33–35} However, a study by Lee et al³² showed that the optimal cut-off points of gait speed to predict activities of daily living were 0.76 m/s and 0.66 m/s for older men and women. A systematic review of Abellan van Kan et al³¹ concluded that older adults who walked >1 m/s had a lower risk of health events and better survival. Individuals categorized as ‘slow walkers’ (<1 m/s) might suffer from functional and cognitive decline, be extremely dependent, and have higher rates of hospitalization. Although cut-points of 0.8 m/s²¹ and 1 m/s³¹ have been used to detect adverse outcomes, the findings of this study showed that the prevalence of older men and women who walked <0.8 m/s and <1 m/s was 1.2% and 5.3% for men and 0.9% and 4.3% for women, respectively. Thus, such cut-off points generated for geriatric assessment may not be applicable and sensitive enough for other populations of older adults. Indeed, we found that the mean gait speed values were 1.33 m/s and 1.36 m/s for older men and women, which surpassed the aforementioned recommended levels.

Although previous findings support the evidence that gait speed is associated with mutual health-related outcomes,³¹ little information has been provided about the association between gait speed and body composition.^{17–20} Excess body fat may impact hip, knee, and ankle joints, leading to shorter step length that may slower gait speed.³⁸ On the other hand, slow gait speed can lead to obesity through a decreased physical activity and more time spent in sedentary behaviors.³⁹ Studies investigating the optimal cut-off points to detect those with high level of fat mass are scarce.¹⁸ A study by Mendes et al¹⁸ showed that to be able to detect those with obesity, the cut-off point of gait speed normalized for height was 0.87 m/s with the AUC of 0.648 ($p < 0.001$) for older women, while for men, the AUC was statistically non-significant and the cut-off value was not provided (0.525, $p = 0.521$). Although evidence suggests adjusting gait speed for height,¹⁸ we observed no correlations between height and gait speed ($r = 0.06$, $p = 0.474$ for older men and $r = 0.07$, $p = 0.259$ for older women). Therefore, previous cut-off points of gait speed to detect obesity seem to be sex-specific and a lot smaller, compared to our findings.¹⁸ The discrepancy between the studies may be explained by using different methods to assess body composition. For example, body-mass index has been extensively used as a “proxy” of body size, especially in terms of undernutrition and overweight/obesity.^{17–19} Although such measure has merit in population-based studies, it cannot adequately discriminate between fat mass and lean muscle mass in clinical practice, leading to a misclassification of nutritional status in older adults.⁴⁰ Also, it has been observed that poorer nutritional status is more related to slower gait speed in older women, compared to older men,^{18,20} which corresponds to findings of the present study. This is not surprising, since the proportion of lean muscle mass to the body weight is much more evident in men, while in women, a higher ratio between fat mass and lean mass is observed.⁴¹ In recent years, a growing public health challenge of obesity has been rapidly increasing in older adults.⁴² With poor nutritional status, older adults experience a decline in physical performance, limiting their ability for independence and engaging in activities of daily living.^{2,3,9,11} Thus, the coexistence between poor nutritional status and slow gait speed is a result of the aging process, where individuals with obesity may embrace slow walking, often accompanied by balance impairments and fatigue.⁴³

This study has a few limitations. First, a cross-sectional design precluded the possibility of establishing causal associations and cut-points based on longitudinal data. Second, the criterion and convergent validity properties and measurement errors for the bioelectrical impedance were considered when compared with dual X-ray absorptiometry,

resulting in fat mass values different from the “true” values. However, previous evidence suggests that bioelectrical impedance may yield good results in comparison to X-ray absorptiometry. Third, we did not include potential residual covariates, such as diet and genetic factors, which might have influenced cut-points and the strength of the associations. Finally, the generalizability of the findings should be interpreted with caution, since older men and women in our study were Caucasians with a median age of 67 years. Thus, future multi-level studies aiming to investigate the associations between gait speed and fat mass should be investigated in different socioeconomic and ethnic classes with wider age range.

Conclusions

In summary, our findings suggest that gait speed is moderately associated with fat mass and newly developed cut-off points of 1.40 m/s and 1.37 m/s may adequately identify older men and women with high level of fat mass, respectively. Gait speed reveals potential utility as a marker for individuals with obesity, but future research is needed to determine its screening and tracking ability of nutritional dysfunctions during the aging process.

Availability of Supporting Data

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics Approval and Consent to Participate

Research involving human subjects complied with all relevant national regulations and institutional policies and is in accordance with the tenets of the Helsinki Declaration (as revised in 2013) and has been approved by the Ethical Committee of the Home of War veterans (Ethical code: 2022/4).

Consent for Publication

The informed consent was voluntarily signed by the participants.

Acknowledgments

The authors would like to thank all the participants for their enthusiastic participation in the study. Also, we would like to thank to Zvonimir Kalčík, PhD, the head of the Home of War Veterans for the approval for conducting the research.

Funding

The author would like to acknowledge the support of Prince Sultan University for paying the article processing charges (APC) of this publication. This study was conducted by the SSDRL research group.

Disclosure

The author reports no conflicts of interest in this work.

References

1. United Nations. *World Population Prospects: The 2004 Revision*. New York: United Nations; 2005.
2. Fulmer T, Reuben DB, Auerbach J, Fick DM, Galambos C, Johnson KS. Actualizing better health and health care for older adults. *Health Aff*. 2021;40(2):219–225. doi:10.1377/hlthaff.2020.01470
3. Salinas-Rodríguez A, Rivera-Almaraz A, Scott A, Manrique-Espinoza B. Severity levels of disability among older adults in low- and middle-income countries: results from the Study on Global Ageing and Adult Health (SAGE). *Front Med*. 2020;7:562963. doi:10.3389/fmed.2020.562963
4. Santos VRD, Christofaro DGD, Gomes IC, Júnior IFF, Gobbo LA. Relationship between obesity, sarcopenia, sarcopenic obesity, and bone mineral density in elderly subjects aged 80 years and over. *Rev Bras Ortop*. 2017;53(3):300–305. doi:10.1016/j.rbo.2017.05.010
5. Sedlmeier AM, Baumeister SE, Weber A, et al. Relation of body fat mass and fat-free mass to total mortality: results from 7 prospective cohort studies. *Am J Clin Nutr*. 2021;113(3):639–646. doi:10.1093/ajcn/nqaa339
6. St-Onge MP, Gallagher D. Body composition changes with aging: the cause or the result of alterations in metabolic rate and macronutrient oxidation? *Nutrition*. 2010;26(2):152–155. doi:10.1016/j.nut.2009.07.004
7. Malenfant JH, Batsis JA. Obesity in the geriatric population - a global health perspective. *J Glob Health Rep*. 2019;3:e2019045. doi:10.29392/joghr.3.e2019045

8. Batsis JA, Villareal DT. Sarcopenic obesity in older adults: aetiology, epidemiology and treatment strategies. *Nat Rev Endocrinol.* 2018;14(9):513–537. doi:10.1038/s41574-018-0062-9
9. Koster A, Ding J, Stenholm S, et al. Does the amount of fat mass predict age-related loss of lean mass, muscle strength, and muscle quality in older adults? *J Gerontol a Biol Sci Med Sci.* 2011;66(8):888–895. doi:10.1093/gerona/glr070
10. Li CW, Yu K, Shyh-Chang N, et al. Pathogenesis of sarcopenia and the relationship with fat mass: descriptive review. *J Cachexia Sarcopenia Muscle.* 2022;13(2):781–794. doi:10.1002/jcsm.12901
11. Hall KS, Cohen HJ, Pieper CF, et al. Physical performance across the adult life span: correlates with age and physical activity. *J Gerontol a Biol Sci Med Sci.* 2017;72(4):572–578. doi:10.1093/gerona/glw120
12. Beaton K, McEvoy C, Grimmer K. Identifying indicators of early functional decline in community-dwelling older people: a review. *Geriatr Gerontol Int.* 2015;15(2):133–140. doi:10.1111/ggi.12379
13. Inzitari M, Calle A, Esteve A, Casas Á, Torrents N, Martínez N. ¿Mides la velocidad de la marcha en tu práctica diaria? Una revisión [Do you measure gait speed in your daily clinical practice? A review]. *Rev Esp Geriatr Gerontol.* 2017;52(1):35–43. doi:10.1016/j.regg.2015.12.010
14. Studenski S, Perera S, Patel K, et al. Gait speed and survival in older adults. *JAMA.* 2011;305(1):50–58. doi:10.1001/jama.2010.1923
15. Norman K, Stobäus N, Gonzalez MC, Schulzke JD, Pirlich M. Hand grip strength: outcome predictor and marker of nutritional status. *Clin Nutr.* 2011;30(2):135–142. doi:10.1016/j.clnu.2010.09.010
16. Toosizadeh N, Wendel C, Hsu CH, Zamrini E, Mohler J. Frailty assessment in older adults using upper-extremity function: index development. *BMC Geriatr.* 2017;17(1):117. doi:10.1186/s12877-017-0509-1
17. Misu S, Asai T, Doi T, et al. Association between gait abnormality and malnutrition in a community-dwelling elderly population. *Geriatr Gerontol Int.* 2017;17(8):1155–1160. doi:10.1111/ggi.12839
18. Mendes J, Borges N, Santos A, et al. Nutritional status and gait speed in a nationwide population-based sample of older adults. *Sci Rep.* 2018;8(1):4227. doi:10.1038/s41598-018-22584-3
19. Tabue-Teguio M, Perès K, Simo N, et al. Gait speed and body mass index: results from the AMI study. *PLoS One.* 2020;15(3):e0229979. doi:10.1371/journal.pone.0229979
20. Beavers KM, Beavers DP, Houston DK, et al. Associations between body composition and gait-speed decline: results from the Health, Aging, and Body Composition study. *Am J Clin Nutr.* 2013;97(3):552–560. doi:10.3945/ajcn.112.047860
21. Verghese J, LeValley A, Hall CB, Katz MJ, Ambrose AF, Lipton RB. Epidemiology of gait disorders in community-residing older adults. *J Am Geriatr Soc.* 2006;54(2):255–261. doi:10.1111/j.1532-5415.2005.00580.x
22. Dommershuijsen LJ, Raganathan J, Ruiter R, et al. Gait speed reference values in community-dwelling older adults - cross-sectional analysis from the Rotterdam Study. *Exp Gerontol.* 2022;158:111646. doi:10.1016/j.exger.2021.111646
23. Pietiläinen KH, Kaye S, Karmi A, Suojanen L, Rissanen A, Virtanen KA. Agreement of bioelectrical impedance with dual-energy X-ray absorptiometry and MRI to estimate changes in body fat, skeletal muscle and visceral fat during a 12-month weight loss intervention. *Br J Nutr.* 2013;109(10):1910–1916. doi:10.1017/S0007114512003698
24. Bouchard DR, Dionne IJ, Brochu M. Sarcopenic/obesity and physical capacity in older men and women: data from the Nutrition as a Determinant of Successful Aging (NuAge)-the Quebec longitudinal Study. *Obesity.* 2009;17(11):2082–2088. doi:10.1038/oby.2009.109
25. Aubertin-Leheudre M, Lord C, Goulet ED, Khalil A, Dionne IJ. Effect of sarcopenia on cardiovascular disease risk factors in obese postmenopausal women. *Obesity.* 2006;14(12):2277–2283. doi:10.1038/oby.2006.267
26. Amaral TF, Santos A, Guerra RS, et al. Nutritional strategies facing an older demographic: the Nutrition UP 65 Study Protocol. *JMIR Res Protoc.* 2016;5(3):e184. doi:10.2196/resprot.6037
27. Wang CY, Chen TR, Lin YH, Liu MH, Chen YC. Gait speed measure: the effect of different measuring distances and the inclusion and exclusion of acceleration and deceleration. *Percept Mot Skills.* 2012;114(2):469–478. doi:10.2466/10.25.26.PMS.114.2.469-478
28. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009;41(1):3–13. doi:10.1249/MSS.0b013e31818cb278
29. Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology.* 1982;143(1):29–36. doi:10.1148/radiology.143.1.7063747
30. Rice ME, Harris GT. Comparing effect sizes in follow-up studies: ROC Area, Cohen's d, and r. *Law Hum Behav.* 2005;29(5):615–620. doi:10.1007/s10979-005-6832-7
31. Abellan van Kan G, Rolland Y, Andrieu S, et al. Gait speed at usual pace as a predictor of adverse outcomes in community-dwelling older people: an International Academy on Nutrition and Aging (IANA) Task Force. *J Nutr Health Aging.* 2009;13(10):881–889. doi:10.1007/s12603-009-0246-z
32. Lee MC, Hsu CC, Tsai YF, Chen CY, Lin CC, Wang CY. Criterion-referenced values of grip strength and usual gait speed using instrumental activities of daily living disability as the criterion. *J Geriatr Phys Ther.* 2018;41(1):14–19. doi:10.1519/JPT.0000000000000106
33. Cesari M, Kritchevsky SB, Penninx BW, et al. Prognostic value of usual gait speed in well-functioning older people--results from the health, aging and body composition study. *J Am Geriatr Soc.* 2005;53(10):1675–1680. doi:10.1111/j.1532-5415.2005.53501.x
34. Cesari M, Kritchevsky SB, Newman AB, et al. Added value of physical performance measures in predicting adverse health-related events: results from the Health, Aging And Body Composition Study. *J Am Geriatr Soc.* 2009;57(2):251–259. doi:10.1111/j.1532-5415.2008.02126.x
35. Rosano C, Newman AB, Katz R, Hirsch CH, Kuller LH. Association between lower digit symbol substitution test score and slower gait and greater risk of mortality and of developing incident disability in well-functioning older adults. *J Am Geriatr Soc.* 2008;56(9):1618–1625. doi:10.1111/j.1532-5415.2008.01856.x
36. Studenski S, Perera S, Wallace D, et al. Physical performance measures in the clinical setting. *J Am Geriatr Soc.* 2003;51(3):314–322. doi:10.1046/j.1532-5415.2003.51104.x
37. Simonsick EM, Newman AB, Visser M, et al. Mobility limitation in self-described well-functioning older adults: importance of endurance walk testing. *J Gerontol a Biol Sci Med Sci.* 2008;63(8):841–847. doi:10.1093/gerona/63.8.841
38. Bindawas SM. Relationship between frequent knee pain, obesity, and gait speed in older adults: data from the osteoarthritis initiative. *Clin Interv Aging.* 2016;11:237–244. doi:10.2147/CIA.S100546
39. McDermott MM, Guralnik JM, Ferrucci L, et al. Community walking speed, sedentary or lying down time, and mortality in peripheral artery disease. *Vasc Med.* 2016;21(2):120–129. doi:10.1177/1358863X15626521

40. Jeong SM, Lee DH, Rezende LFM, Giovannucci EL. Different correlation of body mass index with body fatness and obesity-related biomarker according to age, sex and race-ethnicity. *Sci Rep.* 2023;13(1):3472. doi:10.1038/s41598-023-30527-w
41. López-Ortega M, Arroyo P. Anthropometric characteristics and body composition in Mexican older adults: age and sex differences. *Br J Nutr.* 2016;115(3):490–499. doi:10.1017/S0007114515004626
42. Peralta M, Ramos M, Lipert A, Martins J, Marques A. Prevalence and trends of overweight and obesity in older adults from 10 European countries from 2005 to 2013. *Scand J Public Health.* 2018;46(5):522–529. doi:10.1177/1403494818764810
43. Dufour AB, Hannan MT, Murabito JM, Kiel DP, McLean RR. Sarcopenia definitions considering body size and fat mass are associated with mobility limitations: the Framingham Study. *J Gerontol a Biol Sci Med Sci.* 2013;68(2):168–174. doi:10.1093/gerona/gls109

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>