

● REVIEW

Using antifibrinolytics to tackle neuroinflammation

Stanimir Atsev¹, Nikola Tomov^{2,*}

1 Faculty of Medicine, Trakia University, Stara Zagora, Bulgaria

2 Institute of Anatomy, University of Bern, Bern, Switzerland

Abstract

Plasmin is generally known as a promotor of inflammation. Recent advancement suggests that it has a complex role as immunity modulator. Pharmacological inhibition of plasmin production and activity has been proven to improve neurological outcomes in traumatic brain injury and subarachnoid hemorrhage, most probably by preventing re-bleeding. The immune-modulatory properties of antifibrinolytics, however, suggest that they probably have effects unrelated to fibrinolysis inhibition, which are currently not adequately harnessed. The present work aims to give an account of the existing data regarding antifibrinolytics as agents influencing neuroinflammation. Preclinical and clinical studies on the possible influence of antifibrinolytics on neuroinflammation are scarce. However, the emerging evidence suggests that inhibition of plasmin(ogen) activity can ameliorate neuroinflammation to some extent. This data demonstrate that plasmin(ogen) is not exclusively involved in fibrinolysis, but also has other substrates and can precipitate in inflammatory processes. Investigation on the role of plasmin as the factor for the development of neuroinflammation shows the significant potential of antifibrinolytics as pharmacotherapy of neuroinflammation, which is worthy of further exploration.

*Correspondence to:

Nikola Tomov, MD, PhD,
tomovmd@gmail.com.

orcid:

0000-0002-7072-3832
(Nikola Tomov)

doi: 10.4103/1673-5374.284979

Received: January 16, 2020

Peer review started: January 21, 2020

Accepted: March 4, 2020

Published online: June 19, 2020

Key Words: antifibrinolytics; fibrin; fibrinogen; neuroinflammation; plasmin; plasminogen; tranexamic acid

Introduction

Fibrinogen (coagulation factor I) is a large complex glycoprotein molecule found in the blood of vertebrates. Upon enzymatic cleavage to fibrin by thrombin, it forms the scaffold of the blood clot, functioning primarily to limit bleeding (De Moerloose et al., 2013; Weisel, 2005). Fibrinogen is one of the acute phase proteins, which are upregulated in inflammation, injury, and other pathological settings (Davalos and Akassoglou, 2012). Fibrinogen, and its polymerization product, fibrin, are macromolecules, and are not found in the normal brain tissue due to impenetrability of the blood-brain barrier (BBB) to proteins of this size (Petersen et al., 2018). However, when the BBB is disrupted (in neurodegenerative disease this occurs even before neuronal damage; Zlokovic, 2008), fibrinogen readily extravasates into the brain parenchyma and converted into fibrin (Thomas et al., 1993; Akassoglou and Strickland, 2002).

The physiological process of fibrin clot degradation is known as fibrinolysis. Fibrinolysis is initiated by the conversion of plasminogen to plasmin, mainly through the action of the serine proteases tissue plasminogen activator (tPA) and urokinase plasminogen activator (uPA). Plasminogen, secreted by the liver, binds to fibrin even in its inactivated state. Plasminogen activators act primarily on site of the clot (or, fibrin deposition), converting bound plasminogen to plasmin, which in turn slowly digest fibrin to fibrin-degradation products. Plasmin activity, as well as plasminogen activator activity, are inhibited by several endogenous inhibitors, if not bound to their substrate (Cesarman-Maus and Hajjar, 2005). The fibrinolytic system also regulates the degradation of extracellular matrix in most tissues either directly or by activating matrix metalloproteinases (Mehra et al., 2016).

Antifibrinolytics are a group of medications inhibiting fi-

brinolysis (MeSH ID: D000933); it includes, among others, tranexamic acid (TXA), and aminocaproic acid. The clinical indications usually include bleeding tendencies, such as menorrhagia, as well as hyperfibrinolytic disorders of the hemostatic system (Cai et al., 2019). They are also used in cases in which transfusion of blood products is not an option (Zeybek et al., 2016). The most commonly used antifibrinolytics and their properties are listed in **Table 1**.

Table 1 Some clinically relevant antifibrinolytic agents

Substance	Type of molecule	Main on-label indications
Tranexamic acid	Lysine analogue	Excessive bleeding, hemorrhage prevention in hemophilia
ϵ -Aminocaproic acid	Lysine analogue	Severe postoperative bleeding
4-Aminomethylbenzoic acid	Lysine analogue	Antidote for fibrinolytics

TXA is one of the most widely used and studied antifibrinolytics in the contemporary medical practice, according to the WHO list of essential medicine (2019). TXA (along other antifibrinolytics) is an analogue of the amino acid lysine. Mechanism of action is by occupying lysine binding sites of plasminogen and inhibiting plasmin formation (Reed and Woolley, 2015). TXA is successfully used to treat excessive bleeding in surgery, intensive care, and obstetrics and gynecology (Cai et al., 2019). Recently, the CRASH-3 study has demonstrated that TXA treatment reduces head-injury related death rates, without significantly elevating the thrombotic risk, much concerning in traumatic patients (CRASH-3 trial collaborators, 2019). In another study, it was shown that

TXA treatment reduces mortality of subarachnoid hemorrhage (SAH) patients (Anker-Moller et al., 2017). Despite a second bleed from the same site is often the cause of death following SAH, it is debated, if TXA treatment acts by preventing it, or by other mechanisms (Post et al., 2019). While the most feared complication of antifibrinolysis in general is thrombosis, a recent meta-analysis concluded that the risk of thrombotic complications is not increased by the use of TXA (Chornenki et al., 2019); therefore, TXA treatment is generally safe. TXA is proposed as means to improve clinical condition in patients with hereditary angioedema and to facilitate reduction of skin hyperpigmentation, relying on fibrinolysis-independent mechanisms (Cai et al., 2019). The good clinical results from the application of TXA in traumatic brain injury and SAH, conditions invariably involving neuroinflammation, draw the attention towards the potential of antifibrinolytics as anti-inflammatory drugs.

In January 2020, NCBI PubMed was searched for relevant documents using the following keyword combinations: neuroinflammation and antifibrinolytics, neuroinflammation and 'tranexamic acid', antifibrinolytics and brain. The list of results was manually screened for relevant entries.

Fibrinogen and Fibrin Are Associated with Neuroinflammation

In the affected CNS, fibrinogen stands out among the multiple plasma proteins, which leak through the damaged BBB. Fibrinogen has a distinct molecular structure, which has multiple binding sites for other proteins and receptors, such as albumin, fibronectin, thrombospondin, von Willebrand factor, fibroblast growth factor-2, vascular endothelial growth factor, and interleukin-1 (Weisel, 2005). Fibrinogen can thus interweave multiple signaling pathways, involved in neurological disease (Petersen et al., 2018).

Fibrinogen is a conspicuous finding in lesion areas in neuroinflammation. Its presence in multiple sclerosis (MS) lesions, as well as in amyloid plaques in Alzheimer's disease has been demonstrated on multiple occasions (Adams et al., 2007; Davalos and Akassouglu, 2012), and the amount of fibrinogen deposition and the degree of disruption of BBB usually correlates with disease severity. A neuroinflammatory reaction of the brain is needed to develop the full spectrum of detrimental effects of fibrin(ogen) deposition. Experiments show that in a mouse model of MS, eliminating the affinity of the microglial receptor towards fibrinogen inhibits perivascular clustering of microglia, and reduces axonal damage (Davalos et al., 2012). This demonstrates the role of early BBB leakage and fibrinogen-caused microglial activation in initiating neuroinflammatory disease. For a detailed review of the role of fibrinogen in neurological disease the readers are referred to the work by Petersen et al. (2018), in which they postulate that fibrinogen is a "global mediator of neurodegeneration and activation of innate immunity in the CNS". Data show that this is justified conclusion; however, a conclusion from a mechanistic point of view, that the endogenous fibrinolytic system might counteract fibrin(o-

gen) dependent neuroinflammation by removing fibrin from the CNS has not been substantiated by experiments, as we want to highlight hereafter.

Neuroinflammation Is Contingent on Plasminogen and Plasmin

In the CNS, plasminogen conversion to plasmin is carried out by both tPA and uPA. tPA is constitutively produced by endothelium (Schreiber et al., 1998), but also by neurons (Tsirka et al., 1997; Docagne et al., 1999) and glia (Adhami et al., 2008; Tjärnlund-Wolf et al., 2011). In the same time, uPA has a low baseline expression, but is upregulated in pathological (inflammatory) conditions (Gveric et al., 2001). Plasminogen itself reaches the brain via the systemic circulation, but is also produced by neurons (Basham and Seeds, 2001). Given that, neural tissue, even in its physiological state, possesses a significant plasminogen converting potential. The active plasmin, which could be produced upon plasminogen activation, can impact a number of processes, both physiological and pathological.

Plasmin(ogen) activity in the CNS is physiologically upregulated in axonal growth (Krystosek and Seeds, 1981) and synaptic pruning (Hensch, 2005), and might be crucial in processes of brain development and neural plasticity, although its role in development and regeneration is not completely understood. In excitotoxin-challenged mice, deficiency of tPA indices resistance to neuronal degeneration and toxin-induced seizures (Tsirka et al., 1995), probably by an interaction with the extracellular matrix. An interesting investigation on the role of plasmin(ogen) in neuroinflammation by Shaw et al. (2017) demonstrated that plasminogen deficiency could delay the onset of MS and also protect against demyelination. In the murine model of MS, experimental autoimmune encephalomyelitis, contrary to initial expectations, plasminogen deficient mice had a later onset and attenuated severity of the disease, compared with wild type mice. Pharmacological inhibition of plasmin by TXA also showed comparable effects. The decreased demyelination and microgliosis in plasmin deficiency/inhibition is an evidence that plasmin activity is a modifier of neuroinflammatory demyelination.

Plasminogen deficiency is also linked to attenuated inflammation in another settings, such as LPS induced neuroinflammation. Plasminogen- and tPA-deficient mice showed deficits in the neurovascular integrity, leading to intraparenchymal fibrin deposits in the brain. However, this was not associated with signs of neuroinflammation. Furthermore, upon stimulation with LPS, the neuroinflammatory response was significantly diminished (Hultman et al., 2014). In another mouse study, prolonged activity of the endogenous fibrinolysis system was suspected to be involved in perpetuating posttraumatic neuroinflammation (Hijazi et al., 2015). Additionally, in a mouse model of traumatic brain injury reaching an anti-fibrinolytic state (by application of antifibrinolytic agents or knockout of the plasminogen activator inhibitor) was shown to be neuroprotective (Griemert

et al., 2019). The data from these studies highlight the importance of plasmin(ogen) activity for the propagation of the neuroinflammatory process in various pathological settings. It is worth mentioning (albeit probably not directly plasmin-dependent) that the anti-neuroinflammatory action of antifibrinolytics in mouse models of spinal cord injury, mostly due to reduction of bleeding at lesion (Yoshizaki et al., 2019).

Plasmin-dependent neuroinflammation could be explained by various mechanisms (Figure 1). As a serine protease, plasmin does not have an absolute specificity towards fibrin. Other substrates beyond fibrin could be involved in the reported involvement of plasmin in neuroinflammation. Critical components of the architecture of the BBB are reported to be degraded by plasmin, such as laminin, fibronectin, collagen, and this is critical, since disruption of BBB is prerequisite for initiation and progression of the neuroinflammatory process (Liotta et al., 1981; MacKay et al., 1990; Floris et al., 2004). Furthermore, plasmin is a direct activator of inflammatory cells (Syrovets et al., 2001; Li et al., 2007), and actively binds macrophage receptors that are crucial for their migration (Das et al., 2007; O'Connell et al., 2010; Lighvani et al., 2011). Moreover, plasmin degrades extracellular matrix proteins, thus facilitating inflammatory cell migration across tissue (Rifkin, 1992). Inflammatory cytokines produced by macrophages are also positively correlated with plasmin enzymatic activity *in vitro* (Zalfa et al., 2019).

The fact that plasmin deficiency dampens inflammation even in the presence of significant stimuli (Hultman et al., 2014), shows that plasmin itself is integrally involved in neuroinflammation. Targeting plasmin might be a promising immunomodulatory strategy (Draxler et al., 2019), not only for the nervous system diseases but also for other diseases and injuries.

Perspectives and Outlooks

Considering the growing amount of data elucidating plasmin(ogen) involvement in neuroinflammation, its pharmacological inhibition is an unexplored approach for neuroinflammation suppression. Plasmin activity is unrelated to fibrinolysis, so we believe that the fibrinolytic cascade with its pleiotropic functions is a promising therapeutic target.




Figure 1 A simplified scheme of the involvement of the fibrinolytic cascade in neuroinflammation.

Blue arrows indicate enzymatic conversion, green arrows indicate activation. Red symbols show effect sites of TXA. BBB: Blood-brain barrier; ECM: extracellular matrix; MMPs: metalloproteinases; MΦs: macrophages; PA: plasminogen activators; TXA: tranexamic acid.

When aiming at an antifibrinolytic state, maintaining the fine balance between coagulation and fibrinolysis is of great importance. Antifibrinolysis might exacerbate BBB leakage and therefore aggravate neuroinflammation in some settings (Paul et al., 2007).

It is clear that the enzymatic activities of the individual components of the fibrinolytic cascade are finely interweaved in a network of signaling pathways. For instance, tPA and uPA deficiency on their own do not produce the same results as plasmin(ogen) deficiency/inhibition (Shaw et al., 2017), and inactive tPA has immunomodulatory activity independent on its enzymatic properties (Zalfa et al., 2019). A fine mechanistic dissection of their involvement in neuroinflammation is yet to be done by molecular means.

Inhibition of plasmin(ogen) activity is a justified strategy to suppress neuroinflammation, which has yet to find its translation from animal models to clinical studies. Antifibrinolytics are approved to be safe and well studied, and are powerful tools in the design of future studies. The CRASH-3 study is a fine example of such a study, and its yet to be discussed if anti-inflammatory mechanisms are involved in its results. We believe that exploring their potential off-label use can open the door for novel therapies of neuroinflammatory disease.

Author contributions: Both authors contributed equally to this work and approved the final manuscript.

Conflicts of interest: Both authors declare no conflicts of interest.

Financial support: None.

Copyright license agreement: The Copyright License Agreement has been signed by both authors before publication.

Plagiarism check: Checked twice by iThenticate.

Peer review: Externally peer reviewed.

Open access statement: This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non-Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

References

- Adams RA, Bauer J, Flick MJ, Sikorski SL, Nuriel T, Lassmann H, Degen JL, Akassoglou K (2007) The fibrin-derived γ 377-395 peptide inhibits microglia activation and suppresses relapsing paralysis in central nervous system autoimmune disease. *J Exp Med* 204:571-582.
- Adhami F, Yu D, Yin W, Schloemer A, Burns KA, Liao G, Degen JL, Chen J, Kuan CY (2008) Deleterious effects of plasminogen activators in neonatal cerebral hypoxia-ischemia. *Am J Pathol* 172:1704-1716.
- Akassoglou K, Strickland S (2002) Nervous system pathology: The fibrin perspective. *Biol Chem* 383:37-45.
- Anker-Moller T, Troldborg A, Sunde N, Hvas AM (2017) Evidence for the use of tranexamic acid in subarachnoid and subdural hemorrhage: a systematic review. *Semin Thromb Hemost* 43:750-758.
- Basham ME, Seeds NW (2001) Plasminogen expression in the neonatal and adult mouse brain. *J Neurochem* 77:318-325.
- Cai J, Ribkoff J, Olson S, Raghunathan V, Al-Samkari H, DeLoughery TG, Shatzel JJ (2019) The many roles of tranexamic acid: An overview of the clinical indications for TXA in medical and surgical patients. *Eur J Haematol* 104:79-87.
- Cesarman-Maus G, Hajjar KA (2005) Molecular mechanisms of fibrinolysis. *Br J Haematol* 129:307-321.
- Chornenki NLJ, Um KJ, Mendoza PA, Samienezhad A, Swarup V, Chai-Adisaksopha C, Siegal DM (2019) Risk of venous and arterial thrombosis in non-surgical patients receiving systemic tranexamic acid: A systematic review and meta-analysis. *Thromb Res* 179:81-86.

- CRASH-3 trial collaborators (2019) Effects of tranexamic acid on death, disability, vascular occlusive events and other morbidities in patients with acute traumatic brain injury (CRASH-3): a randomised, placebo-controlled trial. *Lancet* 394:1713-1723.
- Das R, Burke T, Plow EF (2007) Histone H2B as a functionally important plasminogen receptor on macrophages. *Blood* 110:3763-3772.
- Davalos D, Akassoglou K (2012) Fibrinogen as a key regulator of inflammation in disease. *Semin Immunopathol* 34:43-62.
- Davalos D, Kyu Ryu J, Merlini M, Baeten KM, Le Moan N, Petersen MA, Deerinck TJ, Smirnov DS, Bedard C, Hakozaki H, Gonias Murray S, Ling JB, Lassmann H, Degen JL, Ellisman MH, Akassoglou K (2012) Fibrinogen-induced perivascular microglial clustering is required for the development of axonal damage in neuroinflammation. *Nat Commun* 3:1227.
- De Moerloose P, Casini A, Neerman-Arbez M (2013) Congenital fibrinogen disorders: An update. *Semin Thromb Hemost* 39:585-595.
- Docagne F, Nicole O, Marti HH, MacKenzie ET, Buisson A, Vivien D (1999) Transforming growth factor- β 1 as a regulator of the serpins/t-PA axis in cerebral ischemia. *FASEB J* 3:1315-1324.
- Draxler DF, Awad MM, Hanafi G, Daglas M, Ho H, Keragala C, Galle A, Roquilly A, Lyras D, Sashindranath M, Medcalf RL (2019) Tranexamic acid influences the immune response, but not bacterial clearance in a model of post-traumatic brain injury pneumonia. *J Neurotrauma* 36:3297-3308.
- Floris S, Blezer ELA, Schreibelt G, Döpp E, Van Der Pol SMA, Schadee-Eestermans IL, Nicolay K, Dijkstra CD, De Vries HE (2004) Blood-brain barrier permeability and monocyte infiltration in experimental allergic encephalomyelitis: A quantitative MRI study. *Brain* 127:616-627.
- Griemert EV, Schwarzmaier SM, Hummel R, Gözl C, Yang D, Neuhaus W, Burek M, Förster CY, Petkovic I, Trabold R, Plesnila N, Engelhard K, Schäfer MK, Thal SC (2019) Plasminogen activator inhibitor-1 augments damage by impairing fibrinolysis after traumatic brain injury. *Ann Neurol* 85:667-680.
- Gveric D, Hanemaaijer R, Newcombe J, van Lent NA, Sier CF, Cuzner ML (2001) Plasminogen activators in multiple sclerosis lesions: Implications for the inflammatory response and axonal damage. *Brain* 124:1978-1988.
- Hensch TK (2005) Critical period mechanisms in developing visual cortex. *Curr Top Dev Biol* 69:215-237.
- Hijazi N, Fanne RA, Abramovitch R, Yarovoi S, Higazi M, Abdeen S, Basheer M, Maraga E, Cines DB, Higazi AAR (2015) Endogenous plasminogen activators mediate progressive intracerebral hemorrhage after traumatic brain injury in mice. *Blood* 125:2558-2567.
- Hultman K, Cortes-Canteli M, Bounoutas A, Richards AT, Strickland S, Norris EH (2014) Plasmin deficiency leads to fibrin accumulation and a compromised inflammatory response in the mouse brain. *J Thromb Haemost* 12:701-712.
- Krystosek A, Seeds N (1981) Plasminogen activator release at the neuronal growth cone. *Science* 213:1532-1534.
- Li Q, Laumonier Y, Syrovets T, Simmet T (2007) Plasmin triggers cytokine induction in human monocyte-derived macrophages. *Arterioscler Thromb Vasc Biol* 27:1383-1389.
- Lighvani S, Baik N, Diggs JE, Khaldoyanidi S, Parmer RJ, Miles LA (2011) Regulation of macrophage migration by a novel plasminogen receptor P1g-R KT. *Blood* 118:5622-5630.
- Liotta LA, Brundage R, Siegal GP, Goldfarb RH, Terranova V, Garbisa S (1981) Effect of plasminogen activator (urokinase), plasmin, and thrombin on glycoprotein and collagenous components of basement membrane. *Cancer Res* 41:4629-4636.
- MacKay AR, Corbitt RH, Hartzler JL, Thorgeirsson UP (1990) Basement membrane type IV collagen degradation: evidence for the involvement of a proteolytic cascade independent of metalloproteinases. *Cancer Res* 50:5997-6001.
- Mehra A, Ali C, Parcq J, Vivien D, Docagne F (2016) The plasminogen activation system in neuroinflammation. *Biochim Biophys Acta* 1862:395-402.
- O'Connell PA, Surette AP, Liwski RS, Svenningsson P, Waisman DM (2010) S100A10 regulates plasminogen-dependent macrophage invasion. *Blood* 116:1136-1146.
- Paul J, Strickland S, Melchor JP (2007) Fibrin deposition accelerates neurovascular damage and neuroinflammation in mouse models of Alzheimer's disease. *J Exp Med* 204:1999-2008.
- Petersen MA, Ryu JK, Akassoglou K (2018) Fibrinogen in neurological diseases: Mechanisms, imaging and therapeutics. *Nat Rev Neurosci* 19:283-301.
- Post R, Germans MR, Boogaarts HD, Ferreira Dias Xavier B, Van den Berg R, Coert BA, Vandertop WP, Verbaan D (2019) Short-term tranexamic acid treatment reduces in-hospital mortality in aneurysmal sub-arachnoid hemorrhage: A multicenter comparison study. *PLoS One* 14:e0211868
- Reed MR, Woolley LCT (2015) Uses of tranexamic acid. *BJA Educ* 15:32-37.
- Rifkin DB (1992) Plasminogen activator expression and matrix degradation. *Matrix Suppl* 1:20-22.
- Schreiber SS, Tan Z, Sun N, Wang L, Zlokovic BV (1998) Immunohistochemical localization of tissue plasminogen activator in vascular endothelium of stroke-prone regions of the rat brain. *Neurosurgery* 43:909-913.
- Shaw MA, Gao Z, McElhinney KE, Thornton S, Flick MJ, Lane A, Degen JL, Ryu JK, Akassoglou K, Mullins ES (2017) Plasminogen deficiency delays the onset and protects from demyelination and paralysis in autoimmune neuroinflammatory disease. *J Neurosci* 37:3776-3788.
- Syrovets T, Jendrach M, Rohwedder A, Schüle A, Simmet T (2001) Plasmin-induced expression of cytokines and tissue factor in human monocytes involves AP-1 and IKK β -mediated NF- κ B activation. *Blood* 97:3941-3950.
- Thomas WS, Mori E, Copeland BR, Yu JQ, Morrissey JH, del Zoppo GJ (1993) Tissue factor contributes to microvascular defects after focal cerebral ischemia. *Stroke* 24:847-853.
- Tjärnlund-Wolf A, Hultman K, Curtis MA, Faull RLM, Medcalf RL, Jern C (2011) Allelic imbalance of tissue-type plasminogen activator (t-PA) gene expression in human brain tissue. *Thromb Haemost* 105:945-953.
- Tsirka SE, Gualandris A, Amaral DG, Strickland S (1995) Excitotoxin-induced neuronal degeneration and seizure are mediated by tissue plasminogen activator. *Nature* 377:340-344.
- Tsirka SE, Rogove AD, Bugge TH, Degen JL, Strickland S (1997) An extracellular proteolytic cascade promotes neuronal degeneration in the mouse hippocampus. *J Neurosci* 17:543-552.
- Weisel JW (2005) Fibrinogen and fibrin. *Adv Protein Chem* 70:247-299.
- World Health Organization (2019) Model List of Essential Medicines, 21st List. Geneva: World Health Organization.
- Yoshizaki S, Kijima K, Hara M, Saito T, Tamaru T, Tanaka M, Konno DJ, Nakashima Y, Okada S (2019) Tranexamic acid reduces heme cytotoxicity via the TLR4/TNF axis and ameliorates functional recovery after spinal cord injury. *J Neuroinflammation* 16:160.
- Zalfa C, Azmoon P, Mantuano E, Gonias SL (2019) Tissue-type plasminogen activator neutralizes LPS but not protease-activated receptor-mediated inflammatory responses to plasmin. *J Leukoc Biol* 105:729-740.
- Zeybek B, Childress AM, Kilic GS, Phelps JY, Pacheco LD, Carter MA, Borahay MA (2016) Management of the Jehovah's witness in obstetrics and gynecology: A comprehensive medical, ethical, and legal approach. *Obstet Gynecol Surv* 71:488-500.
- Zlokovic BV (2008) The blood-brain barrier in health and chronic neurodegenerative disorders. *Neuron* 57:178-201.