

Improvement of Comorbid Psoriasis in Patients With MS Treated With Natalizumab

Regina Berkovich, MD, PhD, Aida Yakupova, MD, PhD, Jonathan Eskenazi, MD, Noel G. Carlson, PhD, and Lawrence Steinman, MD

Correspondence

Dr. Berkovich
reginaberkovichmd@gmail.com

Neurol Neuroimmunol Neuroinflamm 2021;8:e961. doi:10.1212/NXI.0000000000000961

Abstract

Objective

To present observations on administration of natalizumab to 18 patients with the comorbid MS and psoriasis, who represented a full subset of patients with such comorbidity within the patient records available.

Methods

A retrospective analysis of patient records was performed. Patient histories were gathered and included date of diagnosis of MS and psoriasis, MS disease-modifying therapies (DMTs), Expanded Disability Status Scale (EDSS), reason for DMT switch, and effects on MS and psoriasis status.

Results

On initiation of natalizumab, all 18 patients had a complete cessation of MS disease activity (within 2–8 months) with significant patient-reported improvement of psoriasis (within 1–5 months). This improvement was independent of previous MS therapy and led to 15 of 18 patients needing no additional treatment for MS and psoriasis (remaining 3 patients continued to use topical treatments for psoriasis).

Conclusions

In this cohort of 18 patients with comorbid MS and psoriasis, beneficial results on both diseases were observed after initiation of therapy with natalizumab.

From the LA County and University of Southern California General Hospital and Clinic (R.B.), Los Angeles; Kazan State Medical University (A.Y.), Russia; UCLA (J.E.), Los Angeles, CA; GRECC, VA Salt Lake, University of Utah (N.C.), Salt Lake City; and Stanford University (L.S.), Palo Alto, CA.

Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

DMF = dimethyl fumarate; DMT = disease-modifying therapy; EDSS = Expanded Disability Status Scale; IFN β = interferon-beta.

Natalizumab is a humanized monoclonal antibody that is approved for the treatment of relapsing forms of MS.¹ Its mechanism of action is associated with modulation of lymphocytes trafficking through biological barriers. Current literature on MS and comorbid psoriasis discuss both detrimental and beneficial associations with natalizumab.^{2–5}

In 2010, we reported at the American Academy of Neurology that our first 4 cases showed improvement of psoriasis with natalizumab treatment.⁶ In the present report, we have extended these observations to a total of 18 patients with MS—who represent the entire cohort of patients with such comorbidities from 3 different neurology practices—and were followed with regular intervals over a period from 2004 through 2020.

Methods

The Institutional Review Board found that this research meets requirements for a waiver of consent form under 45 CFR 46.116(f) [2018 Requirements] 45 CFR 46.116(d) [pre-2018 Requirements]. Retrospective data were gathered including date of diagnosis of MS and psoriasis, MS disease-modifying therapies (DMTs), MS activity, Expanded Disability Status Scale (EDSS), reason for DMT switch, concomitant medications, and ongoing data on MS and psoriasis status. To minimize potential bias, clinical observations were obtained from 3 independent neurology clinics.

Every case of comorbid MS and psoriasis was collected, provided that the patient was on natalizumab treatment for MS; this amounted to 18 cases. In all 18 cases, the MS was diagnosed based on concurrent McDonalds criteria. Sixteen of 18 patients had CSF analysis positive for several unique oligoclonal bands (2 patients declined the lumbar puncture procedure). Psoriasis diagnosis was confirmed by treating dermatologists. Psoriasis status was evaluated using the Subject Global Impression scale (grades: much worse, worse, somewhat worse, no change, somewhat better, better, and much better); only “better” and “much better” counted as improvement. No tumor necrosis factor inhibitors were used in the observed cases. All psoriasis cases observed were plaque skin psoriasis. Two cases also had upper extremity nail involvement with cuticle atrophy.

Statistical analysis was performed with Kruskal-Wallis non-parametric analysis of variance using Graph pad InStat software.

Data Availability

Anonymized data will be shared by request from any qualified investigator.

Results

As demonstrated by the data presented in table, interferon-beta (IFN β) treatment was used by 11 patients; in 10 of 11 cases, it coincided either with worsening of psoriasis or with psoriasis developed de novo while on the treatment. IFN β treatment was associated with new MS activity in 3 of 11 patients. According to our data, glatiramer acetate treatment was not associated with psoriasis worsening. Three patients were started on dimethyl fumarate (DMF) but discontinued because of tolerability issues or lack of efficacy for MS. Two patients who discontinued DMF for tolerability reasons were not on DMF for a sufficient amount of time to determine any effects on psoriasis. One DMF patient was on therapy for approximately 2 years and had incomplete control on psoriasis.

All 18 patients at different points were started on natalizumab. One of the patients actually had natalizumab treatment initiated twice—as a first-line DMT and as his 3rd DMT.

In all 18 cases, natalizumab use was associated with patient-reported improvement of the comorbid psoriasis; 16 of 18 patients at the time of last evaluation (2017–2020) were not needed to be on systemic/immunosuppressive therapy for psoriasis. Most cases, positive clinical effects attributed to natalizumab were reported first in the clinical presentation of skin condition before any effects on their MS.

We examined whether the treatment regimens resulted in any significant difference in response time for benefit for either MS or Ps (see last 3 columns in table). There was no significant difference between any of the 3 treatment paradigms (direct transition from IFN to Nat, indirect transfer from IFN to Nat, or no IFN before Nat) for the time for benefit for MS or Ps (data not shown). Furthermore, there was no significant difference between any of these treatment paradigms and the lag in time between MS benefit and beginning of Ps benefit.

Discussion

Given still limited published data, no definite conclusion on direct relationship between natalizumab treatment and psoriasis condition can be drawn. Moreover, no clinical recommendations can be made. However, these observations provide additional data for better understanding of treatment responses of patients having comorbid MS and psoriasis.

Comorbidity of psoriasis in a fraction of patients with MS was recognized several decades ago,^{7,8} and a more recent report suggested that psoriasis comorbidity may influence progression of MS.⁹ There is also similar immunologic connections of

Table Summary of Patient Information

MS Dx	CSF	Base-line EDSS	Psoriasis diagnosis	Psoriasis treatment (t = top, s = steroid)	1st DMT	Year start	Reason for d/c	EDSS	2nd DMT	Year start
Patients treated with IFN										
Direct transfer of IFN to Nat										
2007	Pos	2	2007	t	GA	2007	LOE	2.5	IFNb	2009
2001	Pos	4	2005	t, s	GA	2001	LOE	6	IFNb	2003
2005	Pos	4	2005	t	GA	2005	LOE	4.5	IFNb	2009
2007	Pos	2	2002	t	GA	2007	LOE	4	IFNb	2011
1987	Pos	2.5	1999	t	GA	2000	tlrb	4	IFNb	2002
1999	Pos	4	1998	t	IFNb	1999	Ps	4.5	Nat	2006
1996	Pos	4.5	1999	t	IFNb	1997	LOE, Ps	6	Nat	2006
Indirect transfer of IFN to Nat										
1989	Pos	5	2001	t, s	IFNb	1996	Ps	6.5	GA	2003
2008	Pos	1	2011	t, s	IFNb	2008	Ps	1	GA	2011
2001	Pos	2.5	2000	t, s	IFNb	2001	Ps	2.5	GA	2004
1999	Pos	3	2002	t	IFNb	1999	Ps	6	GA	2004
Patients not treated with IFNb										
2012	Pos	2.5	2010	t, s	GA	2012	LOE	3	Nat	2014
2002	0	3.5	2002	t, s	GA	2004	LOE	4	DMF	2013
2009	0	3	2009	t	GA	2010	LOE	4	DMF	2013
2011	Pos	2.5	1996	t	Nat	2012	JCV+*	2	Fing	2016
2008	Pos	4	2001	t	Nat	2008		3.5		
2006	Pos	3	2004	t, s	GA	2007	Pregn	3.5	GA	2015
2003	Pos	3	2001	t	GA	2006	tlrb	3	DMF	2013

Abbreviations: DMF = dimethyl fumarate; DMT = disease-modifying therapy; Dx = diagnosis; GA = glatiramer acetate; IFNb = interferon-beta; JCV = JC virus; LOE = loss of effectiveness; NA = not applicable; Nat = Natalizumab; pregn = pregnancy; Ps = Psoriasis; tlrb = not tolerated; top = topical. JCV+* occurred in 2016.

psoriasis and MS, with involvement of Th17 cells in both diseases.^{10,11} Although both diseases share common genetic risk factors to influence susceptibility to autoimmune disease, the association of MS with other immune-mediated diseases such as psoriasis has recently been recognized to be due predominantly to surveillance bias or similar influence by environmental factors in both diseases.^{12,13}

Psoriasis is a dermatological chronic immune-mediated disease. The cellular component of the psoriatic lesion tissue includes epidermal infiltration with lymphocytes, including Th1 and Th17, activated T cells, innate lymphocytes, and dendritic cells, causing keratinocyte proliferation.^{14,15} Natalizumab blocks immune cell entry to the CNS including the entry of T cells and B cells.¹⁶ Natalizumab also has been studied for the treatment of the Crohn's disease because it was

shown to modulate adhesion between the leukocytes and endothelial cells.¹⁷ Natalizumab may exert its putative benefit in psoriasis by blocking immune cell infiltration into the skin in a similar manner.

In spite of this relatively close association between MS and psoriasis and higher incidence of psoriasis in MS population compared with the general population, it is well known that at least some MS DMT not only show no efficacy in psoriasis but also may exacerbate the condition. Two hypotheses could be proposed to explain how treatment with natalizumab was associated with a potential improvement of the psoriasis: (1) transition of patients to natalizumab may alleviate exacerbation of psoriasis resulting from an iatrogenic previous DMT or (2) there may be beneficial effects from the natalizumab toward psoriasis.

Table (continued)

Reason for d/c	EDSS	3rd DMT	Year start	EDSS	1st obs benefit MS (mo)	1st obs benefit Ps (mo)	Current Ps treatment
Patients treated with IFN							
Direct transfer of IFN to Nat							
LOE	3	Nat	2011	2.5	4	2	0
Ps	6.5	Nat	2006	6.5	6	3	0
LOE, Ps	4.5	Nat	2012	4	6	4	0
Ps	3	Nat	2012	2.5	2	2	Topical
Ps	5	Nat	2010	5	8	3	0
NA	4.5			4.5	3	2	0
NA	5.5			5.5	4	3	0
Indirect transfer of IFN to Nat							
LOE	6.5	Nat	2007	6	4	1	0
LOE	2	Nat	2012	1	2	1	0
LOE	2.5	Nat	2008	2.5	3	1	0
LOE	6	Nat	2008	6	5	3	Topical
Patients not treated with IFNb							
	3			3	6	3	0
LOE	5	Nat	2015	4	3	2	0
tlrb	4	Nat	2013	3.5	4	1	0
LOE	2.5	Nat	2018	2	3	1	0
					3	2	0
LOE	3.5	Nat	2016	3	5	5	Topical
tlrb	3.5	Nat	2013	3	4	3	0

It is well known, for example, that IFNb treatment has been suggested to potentially trigger onset of psoriasis in patients with MS. In our population, 5 of the 11 patients who received IFNb were subsequently diagnosed with psoriasis after initiation of IFNb therapy. It has been postulated that IFNb may induce the IL-17-driven conditions such as psoriasis because IL-17 enhances keratinocyte proliferation while inhibiting keratinocyte differentiation; in the meantime, keratinocytes induce further Th17 cell recruitment and IL-17 production.¹⁵

Regarding potential improvement of psoriasis after discontinuation of IFNb, 4 of the 11 IFNb patients received another DMT before initiation of natalizumab. Notably, of the remaining 7 IFNb patients who transitioned directly from IFNb to natalizumab, 2 of those patients had lag times between therapies of 2–3 years. These results indicate that mere transition from IFNb may not totally account for the improvement of psoriasis observed with natalizumab treatment and suggest that natalizumab may actually exert some

beneficial effect toward psoriasis. The possible interaction of other MS therapies with psoriasis (and vice versa) has not yet been investigated. Of note, an effective psoriasis treatment with a monoclonal antibody efalizumab that blocks trafficking through inhibition of LFA-1/ICAM-1 interaction was withdrawn because of risk for progressive multifocal leukoencephalopathy.¹⁸ In light of this example where an effective drug for psoriasis blocks trafficking, it would seem quite plausible that another drug for MS, such as natalizumab that also blocks trafficking, might also be effective in psoriasis. This also raises the question as to whether other MS drugs that interfere with trafficking might also have efficacy in psoriasis.

Preclinical studies showed that Th1 cells preferentially infiltrate the spinal cord via an $\alpha 4$ integrin-mediated mechanism, whereas the entry of Th17 cells into the brain parenchyma occurs in the absence of $\alpha 4$ integrins but is dependent on the expression of $\alpha L\beta 2$.¹⁹ If these preclinical findings are applicable to psoriasis, it emphasizes that $\alpha 4$ integrin blockade does provide benefit in psoriasis when accompanied by MS.

There has been one case reported of a patient who developed disseminated psoriasis over the course of six infusions with natalizumab.³ The direct link of natalizumab was not established in this case, but those authors noted that psoriatic (and not rheumatic) arthritis patients can harbor anti-integrin antibodies and that natalizumab might similarly alter laminin function to promote dermal changes seen in psoriasis. However, there are not any subsequent reports of natalizumab-associated exacerbation of psoriasis, suggesting that if this was the cause, then the incidence would be more wide-spread. In our case, we have seen just the opposite in that natalizumab was associated with clinical improvement of psoriasis. It would be potentially more plausible that natalizumab may potentially block the deleterious effects of patient associated anti integrin antibodies. More studies would be needed to test this possibility.

This observational study was limited because only a small number of patients were observed and were not compared against any control group. Further randomized studies will enable neurologists to obtain definitive answers to the optimal therapy when this comorbidity arises.

Acknowledgment

Evan Riddle, a Biogen employee, contributed to the development of this publication. Sergey Grando, Professor of Dermatology, University of California Irvine, and Daniel Arkfeld, Professor of Rheumatology, University of Southern California, contributed to the methodology and results discussion.

Study Funding

There was no funding provided for this study. Partial effort for Noel Carlson was provided by the Department of Veterans Affairs. The authors report no targeted funding.

Disclosure

R. Berkovich received past research support from Biogen, Novartis, Sanofi, Mallinckrodt and Teva and she consulted for Alexion, Bayer, Biogen, Celgene, Genetec, Mallinckrodt, Novartis, Sanofi. A. Yakupova and J. Eskenazi have no financial relationships to disclose. N.G. Carlson received past research support from Biogen outside the scope of this work. L. Steinman has consulted with Roche, Novartis, Tolerion, Atreca, TG Therapeutics, and Atara Biopharma and all these consultations are outside the scope of this work. Go to Neurology.org/NN for full disclosures.

Publication History

Received by *Neurology: Neuroimmunology & Neuroinflammation* August 14, 2020. Accepted in final form December 8, 2020.

Appendix Authors

Name	Location	Contribution
Regina Berkovich, MD, PhD	LA County and University of Southern California General Hospital and Clinic, Los Angeles	Conceptualized the study, major role in the acquisition of data, analyzed the data, and drafted the manuscript for intellectual content
Aida Yakupova, MD	Kazan State Medical University, Russia	Important role in the acquisition of data
Jonathan Eskenazi, MD	UCLA, Los Angeles, CA	Important role in the acquisition of data
Noel G. Carlson, PhD	University of Utah, Salt Lake City	Analyzed the data and drafted the manuscript for intellectual content
Lawrence Steinman, MD	Stanford University, Palo Alto, CA	Conceptualized the study, analyzed the data, and drafted the manuscript for intellectual content

References

- Polman CH, O'Connor PW, Havrdova E, et al; AFFIRM Investigators. A randomized, placebo-controlled trial of natalizumab for relapsing multiple sclerosis. *N Engl J Med* 2006;354:899–910.
- Lambrianides S, Kinnis E, Leonidou E, Pantzaris M. Does natalizumab induce or aggravate psoriasis? A case study and review of literature. *Case Rep Neurol* 2018. 10:286–291.
- Millán-Pascual J, Turpin-Fenoll L, Del Saz-Saucedo P, Rueda-Medina I, Navarro-Muñoz S. Psoriasis during natalizumab treatment for multiple sclerosis. *J Neurol* 2012;259:2758–2760.
- Clark SJ, Wang Q, Mao-Draayer Y. Switching from natalizumab to fingolimod: case report and review of literature. *J Immunol Clin Res* 2016;3:1030.
- Vacchiano V, Foschi M, Sabattini L, Scandellari C, Lugesani A. Arthritic psoriasis during natalizumab treatment: a case report and review of the literature. *Neurol Sci* 2018;39:181–183.
- Berkovich RR. Four cases of comorbid multiple sclerosis and psoriasis: sustained remission of both conditions while on natalizumab. Presented at American Academy of Neurology 62nd Meeting; Toronto, Canada; April 10–17, 2010; Abstract: P06.163.
- Cendrowski W. Multiple sclerosis and psoriasis. *Wiad Lek* 1989;42:575–578.
- Midgard R, Grønning M, Riise T, Kvåle G, Nyland H. Multiple sclerosis and chronic inflammatory diseases. A case-control study. *Acta Neurol Scand* 1996;93:322–328.
- Miron G, Gurevich M, Baum S, Achiron A, Barzilai A. Psoriasis comorbidity affects multiple sclerosis neurological progression: a retrospective case-control analysis. *J Eur Acad Dermatol Venereol* 2017;12:2055–2061.
- Kwok T, Jing Loo W, Guenther L. Psoriasis and multiple sclerosis: is there a link? *J Cutan Med Surg* 2010;14:151–155.
- Axtell RC, Raman C, Steinman L. Interferon- β exacerbates Th17-mediated inflammatory disease. *Trends Immunol* 2011;32:272–277.
- Roshanifefat H, Bahmanyar S, Hillert J, Olsson T, Montgomery S. Shared genetic factors may not explain the raised risk of comorbid inflammatory diseases in multiple sclerosis. *Mult Scler* 2012;18:1430–1436.
- Olafsson S, Stridh P, Bos SD et al. Fourteen sequence variants that associate with multiple sclerosis discovered by meta-analysis informed by genetic correlations. *NPJ Genom Med* 2017;2:24.
- Weller RB, Hunter HJ, Mann MW. *Clinical Dermatology*. Chichester, UK: John Wiley & Sons, Ltd; 2014.
- Deng Y, Chang C, Lu Q. The inflammatory response in psoriasis: a comprehensive review. *Clin Rev Allergy Immunol* 2016;50:377–389.
- Stuve O, Marra CM, Jerome KR, et al. Immune surveillance in multiple sclerosis patients treated with natalizumab. *Ann Neurol* 2006; 59 743–747.
- Guagnozzi D, Caprilli R. Natalizumab in the treatment of Chron's disease. *Biologics* 2008;2:275–284.
- Pugashetti R, Koo JJ. Efalizumab discontinuation: a practical strategy. *Dermatol Treat* 2009;20:132–136.
- Rothhammer V, Heink S, Petermann F, et al. Th17 lymphocytes traffic to the central nervous system independently of $\alpha 4$ integrin expression during EAE. *J Exp Med* 2011;208:2465–2476.