



Draft Genome Sequences of Two *Fusarium oxysporum* Isolates Cultured from Infected *Zinnia hybrida* Plants Grown on the International Space Station

Camilla Urbaniak,^a Gioia Massa,^b Mary Hummerick,^c Christina Khodadad,^c Andrew Schuerger,^d  Kasthuri Venkateswaran^a

^aJet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA

^bExploration Research and Technology Programs, NASA, Kennedy Space Center, Florida, USA

^cAECOM, Kennedy Space Center, Florida, USA

^dSpace Life Sciences Lab, University of Florida, Gainesville, Florida, USA

ABSTRACT Here, we present the whole-genome sequences of two *Fusarium oxysporum* isolates cultured from infected *Zinnia hybrida* plants that were grown on-board the International Space Station (ISS).

The *Fusarium oxysporum* species complex represents one of the most important plant pathogens worldwide, causing disease in many economically important plants and crops (1), but can also cause opportunistic infections in humans (2, 3). However, *F. oxysporum*, like other fungi, produces many bioactive compounds that are beneficial to humans (4, 5) and could be exploited for use in the pharmaceutical/medical industries. During the course of the “Veggie” project (a vegetable production system flown onboard the International Space Station [ISS] to study the effects of the space environment on plant growth and function), *Zinnia hybrida* plants became afflicted with a foliar, stem, and root rot disease caused by the fungus *Fusarium oxysporum*. Isolates were cultured from the leaf (VEG-01C1) and the root (VEG-01C2) of these infected plants, and the draft whole-genome sequences are described herein.

The whole-genome sequences were paired-end sequenced (2×100 bp) on the Illumina HiSeq platform with a 350-bp insert size. A total of 33 million reads (GC content, 47.5%) and 46 million reads (GC content, 46.8%) were obtained from VEG-01C1 and VEG-01C2, respectively. Trimmomatic, on the Galaxy server (<https://usegalaxy.org>), was used to remove the sequencing adaptors (settings, max mismatch, 2; how accurate the match between the two adaptor ligated reads, 30; how accurate the match between any adaptor, 10) and to trim the leading and trailing ends (settings, minimum quality required to keep a base, 3). Postprocessed reads were *de novo* assembled with ABySS version 2.0.2 (6) using *k*-mer sizes of 80 (VEG-01C1) and 88 (VEG-01C2). The VEG-01C1 assembly resulted in a genome size of 49.3 Mb, with an N_{50} value of 376,797 bp. The number of scaffolds generated was 6,455, with a max scaffold length of 1,817,733 bp. The number of scaffolds over 1 kb was 588. The VEG-01C2 assembly resulted in a genome size of 48.9 Mb, with an N_{50} of 334,342 bp. The number of scaffolds generated was 6,398, with a max scaffold length of 2,326,957 bp. The number of scaffolds over 1 kb was 637.

The assembled genomes were compared to those of 66 *F. oxysporum* isolates downloaded from NCBI (ftp://ftp.ncbi.nlm.nih.gov/genomes/genbank/fungi/Fusarium_oxysporum/), as well as those of two isolates cultured from ISS environmental surfaces, using (i) 10 phylogenetically informative loci (7) and (ii) the presence/absence of effector proteins (8). Both methods showed that VEG-01C1 and VEG-01C2 were most closely related to *F. oxysporum* IMV-00293, an isolate cultured in the aftermath of the Chernobyl disaster (GenBank assembly accession number GCA_001931975). To note,

Received 2 April 2018 Accepted 2 April 2018 Published 17 May 2018

Citation Urbaniak C, Massa G, Hummerick M, Khodadad C, Schuerger A, Venkateswaran K. 2018. Draft genome sequences of two *Fusarium oxysporum* isolates cultured from infected *Zinnia hybrida* plants grown on the International Space Station. *Genome Announcements* 6:e00326-18. <https://doi.org/10.1128/genomeA.00326-18>.

Copyright © 2018 Urbaniak et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kasthuri Venkateswaran, kjvenkat@jpl.nasa.gov.

this isolate was incorrectly identified and deposited as *Fusarium solani* (9). Interestingly, *F. oxysporum* strains VEG-01C1, VEG-01C2, and IMV-00293 (i.e., the Chernobyl strain) were very similar to the clinical *F. oxysporum* strain FOSC 3-a (GenBank assembly accession number GCA_000271745), an isolate cultured from blood from a patient in the United States suffering from fusariosis. Comparative genomics of these *Z. hybrida* leaf and root fungal strains with the Chernobyl (IMV-00293) strain could provide insight into which genes could allow for growth in extreme environments, such as those involved in radiation resistance.

Accession number(s). The assembled whole-genome sequences have been deposited in DDBL/EMBL/GenBank under the accession numbers [PXUO00000000](#) (VEG-01C1) and [PXUN00000000](#) (VEG-01C2). The strains have also been deposited in NASA's GeneLab; <https://genelab-data.ndc.nasa.gov/genelab/accession/GLDS-177/>. These are the first versions.

ACKNOWLEDGMENTS

We thank the astronauts for collecting environmental samples aboard the ISS and growing plants in Veggie. We also thank the Microbial Tracking-1 Implementation Team at NASA Ames Research Center for coordinating this effort.

Part of the research described in this publication was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA.

This research was funded by a 2012 Space Biology NNH12ZTT001N grant number 19-12829-26 under Task Order NNN13D111T award to K.V. Funding for C.U. was supported by the NASA Postdoctoral Program. Funding for the Veggie hardware technology validation tests was from NASA's Space Life and Physical Sciences Research and Applications (SLPSRA) Space Biology program. Government sponsorship is acknowledged.

REFERENCES

- Gordon TR. 2017. *Fusarium oxysporum* and the *Fusarium* wilt syndrome. *Annu Rev Phytopathol* 55:23–39. <https://doi.org/10.1146/annurev-phyto-080615-095919>.
- Guarro J, Gené J. 1995. Opportunistic fusarial infections in humans. *Eur J Clin Microbiol Infect Dis* 14:741–754. <https://doi.org/10.1007/BF01690988>.
- O'Donnell K, Sutton DA, Rinaldi MG, Magnon KC, Cox PA, Revankar SG, Sanche S, Geiser DM, Juba JH, van Burik JA, Padhye A, Anaissie EJ, Francesconi A, Walsh TJ, Robinson JS. 2004. Genetic diversity of human pathogenic members of the *Fusarium oxysporum* complex inferred from multilocus DNA sequence data and amplified fragment length polymorphism analyses: evidence for the recent dispersion of a geographically widespread clonal lineage and nosocomial origin. *J Clin Microbiol* 42: 5109–5120. <https://doi.org/10.1128/JCM.42.11.5109-5120.2004>.
- Kour A, Shawl AS, Rehman S, Sultan P, Qazi PH, Suden P, Khajuria RK, Verma V. 2008. Isolation and identification of an endophytic strain of *Fusarium oxysporum* producing podophyllotoxin from *Juniperus recurva*. *World J Microbiol Biotechnol* 24:1115–1121. <https://doi.org/10.1007/s11274-007-9582-5>.
- Nascimento AM, Conti R, Turatti ICC, Cavalcanti BC, Costa-Lotufo LV, Pessoa C, Moraes MO, Manfrim V, Toledo JS, Cruz AK, Pupo MT. 2012. Bioactive extracts and chemical constituents of two endophytic strains of *Fusarium oxysporum*. *Rev Bras Farmacogn* 22:1276–1281. <https://doi.org/10.1590/S0102-695X2012005000106>.
- Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJM, Birol I. 2009. ABySS: a parallel assembler for short read sequence data. *Genome Res* 19:1117–1123. <https://doi.org/10.1101/gr.089532.108>.
- Epstein L, Kaur S, Chang PL, Carrasquilla-Garcia N, Lyu G, Cook DR, Subbarao KV, O'Donnell K. 2017. Races of the celery pathogen *Fusarium oxysporum* f. sp. *apii* are polyphyletic. *Phytopathology* 107:463–473. <https://doi.org/10.1094/PHYTO-04-16-0174-R>.
- van Dam P, Fokkens L, Schmidt SM, Linmans JH, Kistler HC, Ma LJ, Rep M. 2016. Effector profiles distinguish *formae speciales* of *Fusarium oxysporum*. *Environ Microbiol* 18:4087–4102. <https://doi.org/10.1111/1462-2920.13445>.
- Singh NK, Blachowicz A, Romsdahl J, Wang C, Torok T, Venkateswaran K. 2017. Draft genome sequences of several fungal strains selected for exposure to microgravity at the International Space Station. *Genome Announc* 5:e01602-16. <https://doi.org/10.1128/genomeA.01602-16>.