


Article

Internal selective attention is delayed by competition between endogenous and exogenous factors



Edward F. Ester,
Asal Nouri

eester@unr.edu

Highlights

Top-down and bottom-up factors compete to control the locus of external attention

A similar competition also controls the selection of items stored in working memory

Competition produces capture during external but not during internal attention

Ester & Nouri, iScience 26, 107259
 July 21, 2023 © 2023 The Author(s).
<https://doi.org/10.1016/j.isci.2023.107259>



Article

Internal selective attention is delayed by competition between endogenous and exogenous factors

Edward F. Ester^{1,3,*} and Asal Nouri²

SUMMARY

External attention is mediated by competition between endogenous (goal-driven) and exogenous (stimulus-driven) factors, with the balance of competition determining which stimuli are selected. Occasionally, exogenous factors “win” this competition and drive the selection of task-irrelevant stimuli. Endogenous and exogenous selection mechanisms may also compete to control the selection of internal representations (e.g., those stored in working memory), but whether this competition is resolved in the same way as external attention is unknown. Here, we leveraged the high temporal resolution of human EEG to determine how competition between endogenous and exogenous factors influences the selection of internal representations. Unlike external attention, competition did not prompt the selection of task-irrelevant working memory content. Instead, it delayed the endogenous selection of task-relevant working memory content by several hundred milliseconds. Thus, competition between endogenous and exogenous factors influences internal selective attention, but in a different way than external selective attention.

INTRODUCTION

Efficient behavior requires rapid comparison of sensory inputs with internal representations of goal states and motor affordances. Many of these comparisons take place in working memory (WM), a capacity- and duration-limited system that forms a temporal bridge between fleeting sensory phenomena and possible actions.^{1,2} Capacity limits in WM necessitate the existence of external selection mechanisms that gate access to this system (i.e., input gating), while rapidly changing environmental circumstances necessitate the existence of internal selection mechanisms that prioritize behaviorally relevant subsets of information stored in WM for action (i.e., output gating). Whether similar mechanisms mediate the selection of internal and external information is hotly debated.^{3–5}

External sensory inputs can be selected based on behavioral goals (i.e., endogenous selection) or stimulus properties (i.e., exogenous selection), with selection ultimately determined by the balance of competition between these factors. For example, stimulus factors can trigger the selection of task-irrelevant information,^{6–8} disrupting top-down searches for task-relevant stimuli.⁹ These disruptions are frequently accompanied by concurrent shifts in cortical and subcortical spatial priority maps that mediate eye movements and endogenous shifts of covert spatial attention.^{10–12} Endogenous and exogenous factors may also compete to control the selection of internal representations, for example, those stored in WM.¹³ However, little is known about how this competition influences memory performance and is resolved. One obvious possibility is that competition results in the exogenous selection of task-irrelevant information like that seen in external attention. For example, an external stimulus might trigger the exogenous selection of stimulus-matching WM content (i.e., the converse of WM-guided selection, where external attention is oriented to task-irrelevant stimuli that incidentally match attributes of stimuli stored in WM^{13,14}). Alternately, competition between endogenous and exogenous factors could produce a general slowing or delay in the selection of task-relevant memory content without prompting the exogenous selection of task-irrelevant memory content. Although this possibility has been tested and rejected in the external attention literature,¹⁵ it may help explain a recent finding documenting delays in oculomotor biases to the locations of items stored in WM when experimental factors place endogenous and exogenous selection mechanisms in conflict.¹³

¹Department of Psychology and Integrative Neuroscience Program, University of Nevada, Reno, NV, USA

²Center for Complex Systems & Brain Sciences, Florida Atlantic University, 777 Glades Road, Boca Raton, FL, USA

³Lead contact

*Correspondence: eeester@unr.edu

<https://doi.org/10.1016/j.isci.2023.107259>



To test these possibilities, we recorded EEG while participants performed a retrospectively cued WM task typically used to study internal attention.^{16,17} In different experimental blocks, a cue presented during WM storage indicated which of the two memorized positions would be probed for recall (pro-cue trials) or which position would not be probed for recall (anti-cue trials). We reasoned that the anti-cue manipulation would create a state of conflict between endogenous and exogenous selection mechanisms, a point supported by studies documenting visual search costs when participants are cued to the identity of an upcoming distractor.^{13,18} We then examined how informative pro-cues and anti-cues influenced EEG signals that track covert shifts of spatial attention with high temporal precision. Across multiple analyses, we found no evidence for shifts of attention toward cue-matching but task-irrelevant memory representations during the anti-cue task. Instead, we observed a significant delay in the selection of task-relevant WM content during the anti-cue relative to the pro-cue task. Control analyses demonstrated that this result could not be explained by weaknesses in our experimental design or idiosyncrasies in our analytic approach. Thus, we argue that unlike external attention — where competition between endogenous and exogenous selection mechanisms can produce the selection of task-irrelevant information — competition between endogenous and endogenous selection mechanisms during the selection of internal content does not produce an exogenous selection of task-irrelevant information.

RESULTS

We recorded EEG while 40 human volunteers performed a retrospectively cued spatial recall task (Figure 1A). Participants remembered the positions of two discs over a brief delay, and a retrospective color cue presented 1.25 s later instructed participants to continue remembering the positions of both discs (i.e., uninformative trials) or to prioritize one of the discs for subsequent recall (i.e., informative trials). The locations of the two discs were fully randomized across experimental blocks (subject to the constraint that two discs could not appear at the same location on a given trial). At the end of the trial, participants recalled the position of the task-relevant disc via mouse click. Behavioral performance was quantified via average response times and average absolute recall error (i.e., the average absolute difference between the correct and reported disc position). In separate experimental blocks, participants performed a pro-cue task or an anti-cue task. During the pro-cue task informative cues were assigned 100% validity; during the anti-cue task informative cues were assigned 0% validity (i.e., the cue color indicated which disc was task-irrelevant). This allowed us to disentangle the effects of endogenous and exogenous factors on the selection of WM content: during the pro-cue task the color cue indicates which of the two remembered stimuli are task relevant, and endogenous and exogenous selection mechanisms are aligned. During the anti-cue task, however, the color cue indicates which of the two stimuli are task-irrelevant, placing endogenous and exogenous selection mechanisms in competition.¹³ Task order (i.e., pro-followed by anti-cue or vice versa) was counterbalanced across participants, and participants were explicitly reminded about cue validity at the beginning of every block.

Endogenous and exogenous factors influence the selection of task-relevant WM content, but in different ways

A two-factor repeated measures analysis of variance (ANOVA) applied to participants' average absolute recall errors (Figure 1B) revealed a main effect of cue type (i.e., informative versus uninformative; $[F(1,39) = 15.854, p = 0.0003, \eta^2 = 0.289]$), with lower errors during informative versus uninformative cue trials. Likewise, this analysis revealed a main effect of task (i.e., pro-versus anti-cues; $[F(1,39) = 8.168, p = 0.0068, \eta^2 = 0.1732]$), with lower errors during the pro-versus anti-cue task, and a significant interaction between these factors $[F(1,39) = 5.35, p = 0.0261]$. A complementary analysis of response times (Figure 1D) revealed a main effect of cue type $[F(1,39) = 483.046, p < 0.0001, \eta^2 = 0.925]$, with response times faster during informative versus uninformative cue trials, no main effect of task $[F(1,39) = 0.022, p = 0.884, \eta^2 = 0.060]$, and a significant interaction between these factors $[F(1,39) = 30.362, p < 0.0001]$.

Importantly, average absolute recall errors can be influenced by the precision of participants' memory as well as random guessing or accidental reports of a non-probed object ("swap errors"). To quantify the frequencies of random guessing and swap errors, we pooled participants' recall data across all cue conditions (e.g., pro versus anti; informative versus uninformative) and used a hierarchical Bayesian approach to fit participants' data with a parametric model which assumes that on a given trial (1) participants report the position of the probed disk with precision k , (2) participants report the position of the non-probed disk with precision k , or (3) participants randomly guess (Bays et al., 2009). Maximum a posteriori estimates obtained from model fitting indicated that swap errors and random guesses accounted for 1e-04% and




Figure 1. Retro-cue task and memory performance

(A) Participants remembered the locations of two discs over a blank delay. Each disc could appear at one of eight positions along the perimeter of an imaginary circle centered at fixation (upper right panel). (B) Effects of cue type (informative, uninformative) and task type (pro-cue, anti-cue) on average absolute recall errors. (C) We estimated the effects of exogenous factors on recall performance by computing the difference between informative pro-cue trials (i.e., where endogenous and exogenous factors are aligned) and informative anti-cue trials (i.e., where endogenous and exogenous factors are opposed). We estimated the effects of endogenous factors on recall performance by computing the difference between informative pro-cue trials and uninformative pro-cue trials minus the estimated effect of exogenous factors (see text for specifics). Identical analyses were also applied to participants response times (D and E). Error bars depict the 95% confidence interval of the mean.

2.20% of responses at the population level, respectively. This outcome suggests that cue effects on average absolute recall error reflect changes in the precision of participants' memory rather than changes in the frequency of guessing or incorrect responses.

In planned comparisons, we sought further clarity on how endogenous and exogenous factors influenced participants' memory performance. Our approach is based on two assumptions. The first assumption is that during the pro-cue task endogenous and exogenous selection mechanisms are aligned. That is, participants can select the task-relevant position via top-down or bottom-up factors. The second assumption is that during the anti-cue task endogenous and exogenous selection mechanisms conflict: participants can select the task-relevant position only via a top-down interpretation of the cue, whereas bottom-up interpretation of the cue would result in selecting the wrong position. Thus, to isolate the effects of exogenous factors on memory performance we compared participants' recall errors and response times across informative cue trials during the pro- and anti-cue tasks (i.e., the simple effect of task for informative cues).

Next, we calculated the effect of endogenous factors in memory performance via a two-step process. In the first step, we calculated differences in participants' recall errors and response times across informative and uninformative cue trials in the pro-cue task (i.e., the simple effect of cue type for the pro-cue task). In the second step of the analysis, we subtracted the effects of exogenous factors (estimated using the procedure in the preceding paragraph) from these differences, i.e., while accounting for the fact that during the pro-cue task endogenous and exogenous factors are aligned whereas during the anti-cue task they are opposed.

The results of these analyses are summarized in Figures 1C and 1E. Endogenous factors had a facilitatory effect on task performance, lowering recall errors ($M = 1.78^\circ$; 95% CI = 0.645° – 3.112° ; Figure 1C) and speeding response times ($M = 0.165$ s; 95% CI = 0.014 – 0.305 s; Figure 1E). In contrast, exogenous factors significantly worsened participants' recall errors ($M = 0.961^\circ$; 95% CI = 0.191° – 1.863° ; Figure 1C) but had no effect on response times ($M = -0.055$, 95% CI = -0.073 – 0.189 ; Figure 1E). Thus, endogenous and exogenous factors had facilitatory and deleterious effects on participants' memory performance, respectively.

Manipulation check: The anti-cue task requires a greater degree of cognitive control than the pro-cue task

A key assumption of our experimental approach holds that the anti-cue task produces conflict between endogenous and exogenous selection mechanisms. We reasoned that cognitive control is needed to




Figure 2. Frontal theta power is greater during the anti-versus pro-cue task, reflecting a greater need for cognitive control

(A) Time-resolved differences in pro- and anti-cue theta power computed from frontal electrode sites. Theta power estimates were larger during the anti-versus pro-cue task beginning approximately 600 ms after cue onset. Shaded regions depict the 95% confidence interval of the mean. Vertical solid lines at times 0.00 and 3.00 depict the onset of the sample and recall displays, respectively; the vertical dashed line at time 1.75 depicts the onset of an informative retro-cue. The horizontal black bar at the top of the plot marks periods where the difference between anti- and pro-cue theta power was significantly greater than zero (cluster-based permutation tests; see STAR Methods).

(B) Difference in theta-power (4–7 Hz) scalp topography during the pro- and anti-cue tasks. Pro- and anti-cue theta power estimates were averaged over a period spanning 2.5–3.0 s after trial start (i.e., 750–1250 ms after cue onset). Electrode-wise power estimates during the pro-cue task were subtracted from corresponding estimates during the anti-cue task, i.e., larger values indicate higher theta power during the anti-versus pro-cue task.

(C) Task-level differences in power were absent from frequency bands adjacent to theta, including delta (1–3 Hz) and alpha (8–13 Hz).

resolve this competition and drive the selection of task-relevant WM content, and that therefore a greater degree of cognitive control would be required during the anti-cue task compared to the pro-cue task (i.e., when endogenous and exogenous selection mechanisms are aligned). We tested this prediction by estimating and comparing theta power (4–7 Hz) over frontal electrode sites during the pro- and anti-cue tasks. Frontal theta power has robustly linked with the need for cognitive control,¹⁹ scales with WM load,²⁰ and predicts successful WM updating.²¹ Thus, we expected larger frontal theta power estimates during the anti-cue versus the pro-cue task. Indeed, we observed significantly greater frontal theta power during the anti-cue versus pro-cue task that was maximal over frontal midline electrode sites (Figures 2A and 2B). Note that this effect emerged only after presentation of the retro-cue, consistent with a need for “on-line” cognitive control rather than a general increase in difficulty during the anti-cue versus pro-cue task. We also verified that cue-locked frontal power differences were limited to theta-band activity but not neighboring frequency bands (e.g., 1–3 Hz delta-band activity or 8–13 Hz alpha-band activity; Figure 2C). These data support our contention that the anti-cue task produces significant conflict between endogenous and exogenous selection mechanisms.

Competition between endogenous and exogenous selection delays the selection of task-relevant WM content

To understand how competition between endogenous and exogenous factors influence the selection of WM content, we examined how pro- and anti-cues influenced our ability to decode stimulus positions from scalp EEG. Our approach builds on studies demonstrating that stimulus- and location-specific information can be decoded from alpha-band EEG signals²² and that attending to an item or location stored in WM selectively boosts decoding for the attended information.^{23–26} We implemented a multivariate distance-based decoding analysis²⁷ that was customized for our (parametric, circular) location space. This approach is similar to image reconstruction techniques (i.e., “inverted encoding models”) but does not require the experimenter to specify a specific coding model or basis set. To facilitate comparisons across

cue conditions and tasks, participant-level decoding time series were sorted by task relevance: During the pro-cue task decoding performance for the cue-matching disc was designated task-relevant and decoding performance for the cue-nonmatching disc was designated task-irrelevant; during the anti-cue task this mapping was reversed.

We tested two models describing how competition between endogenous and exogenous selection mechanisms influences the prioritization of task-relevant and task-irrelevant WM content. The first model — which we term “retro-capture” — was motivated by studies reporting exogenous shifts of attention to task-irrelevant stimuli in the external attention literature.^{6,7} This model predicts a transient increase in position decoding performance for the cue-matching but task-irrelevant stimulus during the anti-cue task, followed by a later increase in position decoding performance for the cue-nonmatching but task-relevant position (i.e., after the effects of selecting the task-irrelevant stimulus have been resolved). The second model — which we term “delayed selection” — predicts that competition between endogenous and exogenous selection mechanisms merely delays the selection of task-relevant WM content until this competition is resolved. Thus, this model predicts a significant delay in the onset of above-chance position decoding for the cue-nonmatching but task-irrelevant stimulus anti- versus pro-cue task, but no evidence for above-chance decoding of the cue-matching but task-irrelevant stimulus during the anti-cue task.

Our experimental task (Figure 1A) was deliberately constructed so that the effects of endogenous and exogenous factors on the selection of WM contents could be measured during informative and uninformative trials. For example, during uninformative trials participants received an uninformative retrospective cue instructing them to remember the positions of both discs. On presentation of the probe display, this uninformative cue was replaced by a 100% valid (pro-) or 0% valid (anti-) cue instructing participants which disc to report. Conversely, during informative trials pro-cues and anti-cues were presented midway through the storage period. Because informative and uninformative trials had different response demands (i.e., pro- and anti-cues presented at the end of uninformative trials required an immediate response whereas pro- and anti-cues presented during the memory delay during informative trials did not), we analyzed data from these conditions separately.

We first considered data from uninformative cue trials (Figure 3). Task-relevant and task-irrelevant location decoding performance in the pro-cue (Figure 3A) and anti-cue (Figure 3B) tasks increased rapidly during the sample display but returned to chance levels by the time the (uninformative) retro-cue was presented 1.75 s later. Task-irrelevant decoding performance remained at chance levels through the retro-cue and probe displays whereas task-relevant decoding performance increased from chance-to above-chance levels during the probe period. Visual comparisons of probe-locked task-relevant decoding performance suggested that above-chance decoding performance was reached earlier during the pro- relative to the anti-cue task (Figure 3C). To quantify this effect, we extracted and compared probe-locked task-relevant decoding time courses during the pro- and anti-cue tasks via cross-correlation. Specifically, we computed correlations between the timeseries of task-relevant decoding performance during the pro- and anti-cue tasks while temporally shifting the former by -1.0 to $+1.0$ s in 4 msec intervals relative to the latter, yielding a correlation-by-lag function (see STAR Methods). Observed cross-correlation coefficients (Figure 3D) exceeded those expected by chance over lags spanning -0.33 to -0.22 s and fell below those expected by chance over a period spanning $+0.25$ to $+0.35$ s, confirming that task-relevant decoding performance reached above chance levels earlier during the pro- versus anti-cue task.

A complementary analysis of cue-locked decoding performance during informative trials yielded a nearly identical pattern of findings (Figure 4). Specifically, we once again found no evidence for above-chance decoding of the cue-matching but task-irrelevant stimulus position during the anti-cue task (Figure 4B). We did, however, observe a significant delay in the onset of task-relevant decoding performance during the anti- versus pro-cue tasks (Figures 4C and 4D). Thus, the results of probe- and cue-display-locked position decoding performance reveal (1) no evidence for above-chance decoding of the cue- or probe-matching but task-irrelevant stimulus position (Figures 3B and 4B) and (2) a significant delay in the onset of above-chance decoding of the task-relevant stimulus position compared to the pro-cue task (Figures 3C, 3D, 4C, and 4D). These findings are incompatible with a model of internal selective attention where competition between exogenous and endogenous selection mechanisms produces shifts of attention to cue-matching but task-irrelevant WM content.




Figure 3. Location decoding performance during uninformative trials

(A and B) Decoding performance for task-relevant and task-irrelevant locations during pro-cue and anti-cue blocks, respectively.

(C) Overlay of task-relevant location decoding performance for pro-cue and anti-cue blocks (i.e., the blue lines in panels (A and B)). Solid vertical lines at time 0.00 and 3.00 depict the onset of the sample and probe displays, respectively. The dashed vertical line at time 1.75 depicts the onset of the (uninformative) retro-cue. Gray shaded region spanning 0.00–0.50 marks the duration of the sample display. Horizontal bars at the top of each plot mark intervals where decoding performance was significantly greater than zero (nonparametric cluster-based randomization test; see STAR Methods) or intervals where decoding performance for one location was significantly greater than decoding performance for the other location. Shaded regions around each line depict bootstrapped confidence intervals of the mean.

(D) Cross-correlation analysis showing a significant delay in the onset of above-chance probe-locked task-relevant decoding performance during the anti-versus pro-cue task. The null distribution was obtained by repeating the cross-correlation analysis 10,000 times while randomizing participant-level condition labels (i.e., randomly switching the pro- and anti-cue labels). Horizontal bars at the top of the plot depict intervals where the observed cross-correlation coefficient was significantly greater than that expected by chance.

A motivated critic could dismiss our conclusions as based on a null result. For example, perhaps our anti-cue task was insufficient to produce selection of cue-matching yet task-irrelevant stimuli. This argument is difficult to reconcile with behavioral findings showing clear memory impairments during the anti- versus pro-cue task (Figure 1C) and higher cue-locked frontal theta power during the anti- versus pro-cue task (Figure 2) uninformative and informative cue trials in the anti- versus pro-cue tasks (Figure 2). A second possibility is that the parametric similarity-based decoding approach we used is somehow insensitive to resolve the selection of cue-matching but task-irrelevant WM content during the anti-cue task. We tested this possibility by re-analyzing data from informative cue trials using a support vector machine (SVM) based decoding approach (Figure 5) and an inverted encoding model (Figure 6). SVM-based decoding failed to reveal above-chance decoding of the cue-matching but task-irrelevant position during the anti-cue task (Figure 5B). Likewise, the results of the inverted encoding model analysis are a perfect qualitative replication of the pattern reported in Figure 4: we observed no evidence for robust above-chance representations of the cue-matching but task-irrelevant stimulus during the anti-cue task (Figure 6B) and a significant delay in above-chance reconstructions for the task-relevant position during the anti- versus the pro-cue task (Figure 6C). Thus, the findings summarized in Figure 4 generalize across multiple analytic approaches.

Next, we considered the possibility that our decoding approach (Figures 3 and 4) lacked the temporal sensitivity to detect the selection of task-irrelevant WM content. For example, perhaps the temporal dynamics of changes in alpha power are insufficient to measure weak or intermittent (i.e., occurring on only a subset of trials) shifts of attention to the cue-matching but task-irrelevant position during the anti-cue task. We investigated this possibility by tracking the N2pc, an event-related potential (ERP) component known to track covert shifts of attention across visual hemifields. The N2pc is a difference wave defined by




Figure 4. Location decoding performance during informative trials

(A and B) Decoding performance for task-relevant and task-irrelevant locations during pro-cue and anti-cue blocks, respectively.

(C) Overlay of task-relevant location decoding performance for pro-cue and anti-cue blocks (i.e., the blue lines in panels (A and B)).

(D) Cross-correlation analysis showing a significant delay in the onset of above-chance probe-locked task-relevant decoding performance during the anti-versus pro-cue task. All conventions are identical to Figure 3.

greater negative voltages over occipitoparietal electrode sites contralateral (versus ipsilateral) to a visual target beginning ~ 200 ms after stimulus onset²⁸ and can be used to track endogenously and exogenously driven shifts of covert attention with exceptionally high temporal precision.^{29,30} We reasoned that if competition between endogenous and exogenous selection mechanisms produces a selection of cue-matching but task-irrelevant information, then we should observe a significant N2pc over electrode sites contralateral to the visual hemifield containing the cue-matching but task-irrelevant disc during the anti-cue task. Conversely, if competition between endogenous and exogenous selection mechanisms delays the selection of task-relevant WM content, then we should (a) observe a robust N2pc over electrode sites contralateral to the visual hemifield containing the task-relevant disc during the pro-cue task, and (b) observe a significant delay in the onset of the N2pc over electrode sites contralateral to the visual hemifield containing the task-relevant disc during the anti-cue versus pro-cue task.

To test these predictions, we computed voltage differences over occipitoparietal electrode sites contralateral to the visual hemifield containing the task-relevant disc during the pro- and anti-cue tasks. To control for possible sensory confounds we restricted our analyses to trials where the task-relevant and task-irrelevant discs appeared in opposite visual hemifields (approximately 70 trials/task). The N2pc was defined as the average voltage difference over contralateral and ipsilateral electrodes spanning 200–300 ms after cue onset. Because we defined the N2pc with respect to the visual hemifield containing the task-relevant disc, and because we restricted our analysis to trials where the task-relevant and task-irrelevant discs appeared in opposite visual hemifields, shifts of attention toward the cue-matching but task-irrelevant disc during the anti-cue task should manifest as a positive-going waveform 200–300 ms after cue onset.

We observed a statistically robust N2pc from 200 to 300 ms following the appearance of an informative pro-cue (Figure 7), indicating that participants executed a shift of covert visual attention to the visual hemifield containing the cue-matching and task-relevant disc. Conversely, we observed no evidence for a positive-going waveform during the same interval following the presentation of an informative anti-cue. That is, we found no evidence suggesting that participants executed a shift of covert spatial attention toward the visual hemifield containing the cue-matching but task-irrelevant disc during anti-cue trials. Instead, we observed a robust negative-going difference wave beginning ~ 350 ms after the appearance of an anti-cue. We speculate that this negative-going difference wave is identical to the N2pc elicited during the pro-cue task whose onset has been delayed by competition between endogenous and exogenous




Figure 5. Support vector machine-based decoding of stimulus position

To ensure the generality of our findings (e.g., Figure 4), we decoded the positions of the task-relevant and -irrelevant discs during the pro-cue task (A and C) and the anti-cue task (B and C). Plotting conventions are identical to Figure 4. We did not perform a cross-correlation analysis due to the absence of above-chance decoding of the cue-matching but task-relevant stimulus during the anti-cue task.

selection mechanisms. Nevertheless, the results of this analysis provide converging evidence against the hypothesis that competition between endogenous and exogenous selection mechanisms drives the inadvertent selection of cue-matching but task-irrelevant information.

Other alternative explanations

Next, we considered the possibility that evidence for the selection of the task-irrelevant disc during the anti-cue task was obscured by trial averaging. For example, perhaps the selection effect is small, short lived, or intermittent (i.e., occurring on only a subset of trials). We tested this possibility by recomputing alpha-band-based decoding performance for the task-irrelevant disc after sorting participants' anti-cue task performance by median recall error (i.e., "high" versus "low"). Here, we reasoned that because exogenous factors have a deleterious effect on participants' recall errors during the anti-cue task (Figure 1C), exogenous selection of the task-irrelevant disc — as indexed by higher task-irrelevant decoding performance — should be more evident during high recall error trials. However, this was not the case: We observed no evidence for above-chance task-irrelevant decoding performance during low- or high-error informative (Figure 8A) or uninformative trials (Figure 8B). Thus, it is unlikely that the pattern of exogenous-then-endogenous selection predicted by the retro-capture model was obscured by trial-averaging.

We also considered the hypothesis that selection of the cue-matching but task-irrelevant disc during the anti-cue task was obscured by successful cognitive control. Specifically, we reasoned that shifts of attention toward the location of the task-irrelevant disc might be more likely during trials contaminated by lapses of attention. To test this hypothesis, we re-computed cue-matching but task-irrelevant location decoding performance after sorting participants' alpha-band EEG data by frontal theta power (Figure 2), reasoning that inadvertent selection of cue-matching but task-irrelevant stimuli would be more likely during trials where frontal theta power (indexing cognitive control) was low versus high. However, we observed no evidence for above-chance decoding of the cue-matching but task-irrelevant position during high- or low-theta power trials (Figure 9). This analysis provides converging evidence suggesting that exogenous factors do not lead to a selection or re-activation of cue-matching but task-irrelevant WM content, but instead delay the endogenous selection of task-relevant WM content.

Finally, although our analyses reveal no evidence for a selection of cue-matching but task-irrelevant information during the anti-cue task, they do reveal a significant delay in the selection of task-relevant




Figure 6. Inverted encoding model analysis

(A–C) We modeled patterns of alpha-band activity at each electrode site as a weighted combination of eight position filters, each with an idealized tuning curve. Filter weights from each electrode were used to reconstruct a representation of remembered position(s) in an independent test dataset. Conventions are identical to Figure 4.

information during the anti-versus pro-cue tasks (Figures 3C, 3D, 4C, and 4D). This effect could reflect a delay in the selection of task-relevant information caused by competition between endogenous and exogenous selection mechanisms during the anti-cue task or some other task-specific factor. For example, one trivial possibility is that it simply takes participants longer to interpret anti-cues versus pro-cues. However, this explanation is difficult to reconcile with the fact that neither the main effect of task (i.e., pro-cue versus anti-cue; Figure 1D) nor the simple effect of task (Figure 1E) on response times during informative cue trials reached significance. A second possibility is that delayed above-chance decoding performance during the anti-cue task was caused by carryover effects. For example, although task order was counterbalanced across observers, perhaps participants who completed the pro-cue task followed by the anti-cue task had extra difficulty interpreting anti-cues compared to participants who performed the anti-cue task followed by the pro-cue task. To test this possibility, we compared the time-courses of task-relevant decoding performance during informative anti-cue trials in participants who performed the pro-cue task followed by the anti-cue task ($N = 17$) or vice versa ($N = 23$). For both groups, task-relevant decoding performance reached above chance levels shortly before or immediately after the onset of the probe display (Figure 10). If anything, the onset of above-chance decoding performance occurred earlier for participants who performed the anti-cue task second versus those who performed the anti-cue task first, though this difference was not significant ($p = 0.146$; randomization test, see STAR Methods). Thus, order effects cannot account for delays in task-relevant decoding performance during the anti-cue versus pro-cue blocks.

Eye movement control analysis

Finally, we investigated the possibility that our key findings (e.g., Figures 3, 4, 5, 6, and 7) were influenced by oculomotor artifacts. Although we used independent components analysis to identify and remove large oculomotor artifacts from the EEG data, several recent reports have documented the existence of small ($\leq 0.5^\circ$ visual angle) but consistent gaze position biases toward the position of a behaviorally relevant item stored in WM, especially following the appearance of a retrospective cue.¹³ Moreover, there is some evidence suggesting that these gaze position biases can contribute to EEG decoding performance.^{31,32} Although we did not collect precise eye position data during this experiment, we reasoned that because of volume conduction gaze biases would have the largest effects on EEG signals at extreme frontal electrode sites. Therefore, if gaze biases contribute to decoding performance, it should be possible to decode stimulus positions from alpha-band filtered data at these same electrode sites.³³ To investigate this possibility, we attempted to decode the positions of the cue-matching and cue-nonmatching stimuli from 10 to 20 electrode sites Fp1, Fp2, AF7, AF3, AFz, AF4, and AF8 during informative pro- and anti-cue trials (using the same parametric




Figure 7. Event-related potentials reveal delayed selection of task-relevant WM content during the anti-cue task

(A) Average contralateral and ipsilateral ERP waveforms during the pro-cue task, time-locked to trial start (0.00 s). The vertical lines at times 1.75 and 3.00 s depict the onset of the retro-cue and probe displays, respectively. The shaded region depicts the duration of the sample display.

(B) Identical to (A), but for the anti-cue task.

(C) Difference waves (i.e., contralateral-ipsilateral) time locked to retro-cue onset (time 0 ms). The shaded region spanning 200–300 ms depicts the canonical N2pc window. Horizontal bars at the top of the plot mark epochs where difference wave voltage was significantly greater than chance (red bar) or when anti-cue difference wave voltage was significantly greater than pro-cue difference wave voltage (maroon bar). Shaded regions depict the 95% confidence interval of the mean.

(D) N2pc amplitudes, defined as the average difference wave voltage over a period spanning 200–300 ms after cue onset. Error bars depict the 95% confidence interval of the mean; *, $p < 0.05$, bootstrap test.

decoding analysis used to produce the data shown in Figure 4). Apart from a brief epoch of above-chance decoding performance for the cue-matching disc during the pro-cue task that was limited to the probe epoch, this analysis failed to reveal robust above-chance decoding of stimulus position resembling that seen in our primary analyses (Figure 11; compare with Figure 4). Thus, it is unlikely that our key findings can be attributed to subtle differences in gaze bias across experimental conditions.

DISCUSSION

Selective attention can be allocated to sensory inputs and internal representations based on voluntary, endogenous factors or involuntary, exogenous factors. An enormous literature suggests that external selection is mediated by competition between endogenous and exogenous factors, with the focus of selection determined by the balance of competition between these factors.⁹ Endogenous and exogenous factors may also compete to control the selection of internal representations, for example, those stored in WM.¹³ Here, we show that - unlike external attention - this competition does not result in a selection of task-irrelevant stimuli. This, in turn, supports the hypothesis that internal and external selective attention are mediated by at least partially non-overlapping mechanisms.

A motivated critic could dismiss our conclusion as based on a null result. For example, perhaps our experimental approach was insufficient at creating conditions conducive to the exogenous selection of cue-matching but task-irrelevant stimuli during the anti-cue task. While we cannot fully exclude this possibility, we note the following: First, we point critics toward a recent paper by van Ede et al.¹³ who used a behavioral task and cue manipulation like the one reported here to document evidence for oculomotor capture by cue-matching but task-irrelevant stimuli during an anti-cue task. In that study, participants memorized the orientations of two colored bars (one per visual hemifield), and a color cue presented during storage indicated which bar should be probed for report. Using this approach, van Ede et al.¹³ reported that during anti-cue trials gaze position was subtly biased toward the location of the cue-matching but task-irrelevant stimulus before “flipping” to the cue-nonmatching but task-relevant stimulus (see their Figures 2C and 3A). Thus, their experimental setup - which was highly similar to ours - was sufficient to produce oculomotor capture by task-irrelevant stimuli (we return to this point below). Second, in the current study participants’ memory performance was significantly worse during the anti- versus pro-cue tasks (Figure 1B) and that the




Figure 8. Split-half analysis of task-irrelevant decoding performance during the anti-cue task

To examine whether exogenous selection of the task-irrelevant disc during anti-cue blocks was obscured by trial averaging, we sorted task-irrelevant decoding performance during uninformative (top) and informative (bottom) trials by participants' recall errors. We reasoned that because exogenous factors have a deleterious effect on participants' recall errors (Figure 1C), exogenous selection of the task-irrelevant disc – as indexed by higher task-irrelevant decoding performance – should be more evident during high recall error trials (black lines) than low recall error trials (green lines). However, we observed no evidence for above-chance task-irrelevant decoding performance in any of the conditions we examined. Plotting conventions are identical to those in Figure 4.

appearance of an anti-cue led to a significant increase in frontal theta power compared to the appearance of a pro-cue (Figure 2), consistent with a need for greater cognitive control during the anti- versus pro-cue task. Third, perhaps our specific decoding approach lacked the sensitivity to identify the selection of task-irrelevant information during the anti-cue task. Again, it is difficult to fully exclude this possibility, but we note that we observed qualitatively different patterns of findings across two different decoding methods (similarity-based versus support vector machine-based; Figures 4 and 5, respectively) and the results of an inverted encoding model analysis where we reconstructed remembered positions from EEG activity (Figure 6). Fourth, perhaps the posterior alpha-band signal (8–13 Hz) lacks the temporal resolution necessary to resolve fleeting or intermittent selection of task-irrelevant information during the anti-cue task. However, analyses of the N2pc ERP component responses (with a temporal resolution of ~4 ms) revealed clear evidence for the selection of the task-relevant disc during the pro- and anti-cue tasks but no evidence for the selection of the task-irrelevant disc during the anti-cue task (Figure 7). Fifth, a variety of additional control analyses demonstrate that the selection of task-irrelevant information during the anti-cue task was not obscured by high behavioral performance (Figure 8), successful cognitive control (Figure 9), or task order effects (Figure 10). Importantly, in many of these analyses we *did* find evidence for a temporal delay in the selection of task-relevant WM content during the anti-cue versus the pro-cue task. Thus, we argue that unlike external attention — where competition between endogenous and exogenous selection mechanisms produces clear evidence for the selection of irrelevant stimuli — competition between endogenous and endogenous internal selection mechanisms does not produce a selection of task-irrelevant memory content and is resolved in a fundamentally different way. More generally, this result points toward important differences in how voluntary and involuntary selection mechanisms compete to control the processing of external sensory inputs versus internal memory representations.




Figure 9. Task-irrelevant decoding performance during the anti-cue task sorted by frontal theta power
Conventions are identical to Figure 4B.

As noted above, a recent study by van Ede et al.¹³ reported evidence for oculomotor capture by cue-matching but task-irrelevant stimuli using a task design and cue manipulation nearly identical to that used in the previous study. Conversely, we found no evidence for this kind of “retro-capture” effect in our EEG data. How can these results be reconciled? We believe that the key lies within recent studies demonstrating dissociations between shifts of covert spatial attention indexed by oculomotor biases and EEG signals. For example, although attention-related modulations of cortical and subcortical processing are larger during trials containing microsaccades toward the location of a (covertly) attended stimulus, clear attention-related modulations are also observed in the absence of microsaccades.^{33,34} The apparent contradiction between our EEG findings and earlier oculomotor findings¹³ provides additional impetus to further explore relationships between oculomotor and EEG signatures of covert spatial attention.

Our study complements earlier efforts that examined the role of active forgetting in WM.^{35,36} For example, Williams et al.³⁶ showed participants successive displays of to-be-remembered stimuli that were followed by a cue instructing participants which display to remember (i.e., “directed remembering”) or which display to ignore (i.e., “directed forgetting”). These authors found that directed forgetting cues improved WM performance compared to an uninformative cue condition, but less so than directed remembering cues. Importantly, these studies utilized spatial or conceptual cues, including pointed arrow symbols³⁶ or written words.³⁵ Conversely, in the current study, our retrospective cues always matched one feature of an item stored in WM. Like prior work¹³ independently manipulating the feature match and the meaning of the cue (i.e., pro-versus anti-) allowed us to track the consequences of placing endogenous and exogenous selection mechanisms in conflict.

The current findings may inform neurocomputational models of WM. For example, conjunctive coding models predict that WM representations are maintained by spiking activity in feature- and/or location-specific neural populations.^{37,38} Although the exact mechanisms vary by implementation, these models generally predict that a feature probe in one dimension (e.g., orientation) activates spiking patterns in neural populations that code this feature and those that code other features of the same object (e.g., color) and/or its location. This, in turn, enables robust readout of the probed and non-probed stimulus dimensions by downstream neural populations. Although these models were not developed to describe the anti-cue task contemplated here, one could reasonably predict an increase in task-irrelevant decoding performance after presentation of an anti-cue based on their general architecture. We observed no evidence for such an effect, and it remains to be seen whether these models can be modified to predict behavioral and neural data during pro- and anti-cue tasks. Alternately, pattern completion models predict that the contents of WM reside in different neural states — an “active” state mediated by sustained spiking activity and a “latent” state mediated by short-term synaptic plasticity.^{39,40} Presentation of a feature probe that matches a stimulus stored in a latent format reinstates activity patterns evoked when that stimulus was encoding, prompting and/or “refreshing” of the neural representation of the probe-matching item through pattern completion. This prediction enjoys some support: a representation stored in WM item can be “re-activated” (as indexed by above-chance EEG decoding performance) by a task-irrelevant sensory input²⁷ or




Figure 10. Delayed improvements in task-relevant decoding performance during the anti-cue task cannot be explained by order effects

We tested whether delayed improvements in task-relevant decoding performance during the anti-cue (versus pro-cue) task were caused by order effects by splitting decoding performance across participants who performed the anti-cue task followed by the pro-cue task (green) or vice versa (maroon). If anything, above-chance decoding performance was reached earlier for participants who completed the pro-cue followed by the anti-cue tasks, though this effect was not significant ($p = 0.141$; randomization test).

a transcranial magnetic stimulation (TMS) pulse applied over sensory cortex.⁴¹ Conversely, in the current study we found no evidence for a reactivation of cue-matching but task-irrelevant WM content following presentation of an anti-cue. However, one salient difference between the current study and prior work is that in the latter, an informative retrospective cue instructed presented before the “impulse” stimulus instructed participants which of two remembered stimuli should be prioritized for report. Thus, re-activation




Figure 11. Stimulus position cannot be decoded from frontal alpha-band activity

Conventions are identical to Figures 4A and 4B.

of information stored in synaptic traces may be contingent on the network responsible for storing information to be selected or otherwise primed for decision making and action.

To summarize, the current findings support recent suggestions that endogenous and exogenous selection mechanisms compete to control access to internal WM representations. However, this competition is resolved in a fundamentally different way than that seen during external attention. Specifically, endogenous and exogenous competition does not produce an errant selection or refreshing of salient but task-irrelevant WM content. Instead, this competition delays the selection of task-relevant memory content by endogenous mechanisms.

Limitations of the study

Our findings reveal an apparent contradiction between oculomotor and electrophysiological signatures of internal attention during the selection of task-relevant WM content. On the one hand, an earlier eye tracking study that used an experimental design like the one reported here found evidence for oculomotor capture by cue-matching but task-irrelevant stimuli during an anti-cue task.¹³ Conversely, we found no evidence for such an effect in EEG. Although caution is always required in interpreting a null result, several control analyses (summarized in the first paragraph of the discussion) suggest that the absence of capture by cue-matching but task-irrelevant stimuli during the anti-cue task are not because of limitations in our experimental design or analytic sensitivity. When compared to earlier results,¹³ our findings complement recent demonstrations suggesting that oculomotor and electrophysiological measurements may index different selection mechanisms³⁴ and provide further motivation for studies directly comparing oculomotor and electrophysiological indices of attentional selection.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- [KEY RESOURCES TABLE](#)
- [RESOURCE AVAILABILITY](#)
 - Lead contact
 - Materials availability
 - Data and code availability
- [EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS](#)
- [METHOD DETAILS](#)
 - Testing environment
 - Spatial retrocue task
 - Quantifying memory performance
 - EEG acquisition and preprocessing
 - Data cleanup
 - Decoding spatial positions from posterior alpha-band EEG signals
 - Decoding and filter cutoffs
 - Cross-correlation analyses
 - Quantifying frontal theta power
 - N2pc analyses
 - Inverted encoding model
- [QUANTIFICATION AND STATISTICAL ANALYSIS](#)
 - General
 - Statistical power
 - Statistical comparisons – Behavioral data
 - Statistical comparisons – N2pc, decoding performance, and inverted encoding model

ACKNOWLEDGMENTS

Funding: National Science Foundation Grant 2050833 (EFE).

AUTHOR CONTRIBUTIONS

Conceptualization: E.F.E. and A.N.

Methodology: E.F.E. and A.N.

Investigation: A.N.

Visualization: E.F.E. and A.N.

Supervision: E.F.E.

Writing – original draft: E.F.E.

Writing – review and editing: E.F.E. and A.N.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: March 28, 2023

Revised: May 18, 2023

Accepted: June 27, 2023

Published: July 3, 2023

REFERENCES

- D'Esposito, M., and Postle, B.R. (2015). The cognitive neuroscience of working memory. *Annu. Rev. Psychol.* *66*, 115–142.
- van Ede, F., and Nobre, A.C. (2023). Turning attention inside-out: How working memory serves behavior. *Annu. Rev. Psychol.* *74*, 137–165.
- Chun, M.M., Golomb, J.D., and Turk-Browne, N.B. (2011). A taxonomy of external and internal attention. *Annu. Rev. Psychol.* *62*, 73–101.
- Chatham, C.H., and Badre, D. (2015). Multiple gates on working memory. *Curr. Opin. Behav. Sci.* *1*, 23–31.
- Rac-Lubashevsky, R., and Frank, M.J. (2021). Analogous computations in working memory input, output, and motor gating: Electrophysiological and computational modeling evidence. *PLoS Comput. Biol.* *17*, e1008971.
- Theeuwes, J. (1992). Perceptual selectivity for color and form. *Percept. Psychophys.* *51*, 599–606.
- Folk, C.L., Remington, R.W., and Johnston, J.C. (1992). Involuntary covert orienting is contingent on attentional control settings. *J. Exp. Psychol. Hum. Percept. Perform.* *18*, 1030–1044.
- Anderson, B.A., Laurent, P.A., and Yantis, S. (2011). Value-driven attentional capture. *Proc. Natl. Acad. Sci. USA* *108*, 10367–10371.
- Wolfe, J.M. (2020). Visual search: How do we find what we are looking for? *Atten. Percept. Psychophys.* *82*, 1–6.
- Bisley, J.W., and Goldberg, M.E. (2010). Attention, intention, and priority in the parietal lobe. *Annu. Rev. Neurosci.* *33*, 1–21.
- Sprague, T.C., and Serences, J.T. (2013). Attention modulates spatial priority maps in the human occipital, parietal, and frontal cortices. *Nat. Neurosci.* *16*, 1879–1887.
- Luck, S.J., Gaspelin, N., Folk, C.L., Remington, R.W., and Theeuwes, J. (2021). Progress towards resolving the attentional capture debate. *Vis. cogn.* *29*, 1–21.
- van Ede, F., Board, A.G., and Nobre, A.C. (2020). Goal-directed and stimulus-driven selection of internal representations. *Proc. Natl. Acad. Sci. USA* *117*, 24590–24598.
- Olivers, C.N.L., Peters, J., Houtkamp, R., and Roelfsema, P.R. (2011). Different states in visual working memory: When it guides attention and when it does not. *Trends Cogn. Sci.* *15*, 327–334.
- Remington, R.W., Folk, C.L., and Mclean, J.P. (2001). Contingent attentional capture or delayed allocation of attention? *Percept. Psychophys.* *63*, 298–307.
- Griffin, I.C., and Nobre, A.C. (2003). Orienting attention to locations in internal representations. *J. Cogn. Neurosci.* *15*, 1176–1194.
- Landman, R., Spekreijse, H., and Lamme, V.A.F. (2003). Large capacity storage of integrated objects before change blindness. *Vision Res.* *43*, 149–164.
- Moher, J., and Egeth, H.E. (2012). The ignoring paradox: Cueing distractor features leads first to selection, then to inhibition of to-be-ignored items. *Atten. Percept. Psychophys.* *74*, 1590–1605.
- Cavanagh, J.F., and Frank, M.J. (2014). Frontal theta as a mechanism for cognitive control. *Trends Cogn. Sci.* *18*, 414–421.
- Jensen, O., and Tesche, C.D. (2002). Frontal theta activity in humans increases with memory load in a working memory task. *Eur. J. Neurosci.* *15*, 1395–1399.
- Itthipuripat, S., Wessel, J.R., and Aron, A.R. (2013). Frontal theta is a signature of successful working memory manipulation. *Exp. Brain Res.* *224*, 255–262.
- Foster, J.J., Sutterer, D.W., Serences, J.T., Vogel, E.K., and Awh, E. (2016). The topography of alpha-band activity tracks the content of spatial working memory. *J. Neurophysiol.* *115*, 168–177.
- Ester, E.F., Nouri, A., and Rodriguez, L. (2018). Retrospective cues mitigate information loss in human cortex during working memory storage. *J. Neurosci.* *38*, 8538–8548.
- LaRocque, J.J., Lewis-Peacock, J.A., Drysdale, A.T., Oberauer, K., and Postle, B.R. (2013). Decoding attended information in short-term memory: an EEG study. *J. Cogn. Neurosci.* *25*, 127–142.
- Sprague, T.C., Ester, E.F., and Serences, J.T. (2016). Restoring latent visual working memory representations in human cortex. *Neuron* *91*, 694–707.
- Samaha, J., Sprague, T.C., and Postle, B.R. (2016). Decoding and reconstructing the focus of spatial attention from the topography of alpha-band oscillations. *J. Cogn. Neurosci.* *28*, 1090–1097.
- Wolff, M.J., Jochim, J., Akyürek, E.G., and Stokes, M.G. (2017). Dynamic hidden states underlying working-memory-guided behavior. *Nat. Neurosci.* *20*, 864–871.
- Luck, S.J., and Hillyard, S.A. (1994). Spatial filtering during visual search: Evidence from human electrophysiology. *J. Exp. Psychol. Hum. Percept. Perform.* *20*, 1000–1014.
- Hickey, C., McDonald, J.J., and Theeuwes, J. (2006). Electrophysiological evidence of the

- capture of visual attention. *J. Cogn. Neurosci.* 18, 604–613.
30. Burra, N., and Kerzel, D. (2013). Attentional capture during visual search is attenuated by target predictability: Evidence from the N2pc, Pd, and topographic segmentation. *Psychophysiology* 50, 422–430.
 31. Mostert, P., Albers, A.M., Brinkman, L., Todorova, L., Kok, P., and de Lange, F.P. (2018). Eye movement-related confounds in neural decoding of visual working memory representations. *Eneuro* 5. ENEURO.0401, 17.2018.
 32. Quax, S.C., Dijkstra, N., van Staveren, M.J., Bosch, S.E., and van Gerven, M.A.J. (2019). Eye movements explain decodability during perception and cued attention in MEG. *Neuroimage* 195, 444–453.
 33. Ester, E.F., and Pytel, P. (2023). Changes in behavioral priority influence the accessibility of working memory content. *Neuroimage* 272, 120055.
 34. Liu, B., Nobre, A.C., and van Ede, F. (2022). Functional but not obligatory link between microsaccades and neural modulation by covert spatial attention. *Nat. Commun.* 13, 3503.
 35. Williams, M., and Woodman, G.F. (2012). Directed forgetting and directed remembering in visual working memory. *J. Exp. Psychol. Learn. Mem. Cogn.* 38, 1206–1220.
 36. Williams, M., Hong, S.W., Kang, M.-S., Carlisle, N.B., and Woodman, G.F. (2013). The benefit of forgetting. *Psychon. Bull. Rev.* 20, 348–355.
 37. Schneegans, S., and Bays, P.M. (2017). Neural architecture for feature binding in visual working memory. *J. Neurosci.* 37, 3913–3925.
 38. Schneegans, S., and Bays, P.M. (2019). Bays PM New perspectives on binding in visual working memory. *Br. J. Psychol.* 110, 207–244.
 39. Mongillo, G., Barak, O., and Tsodyks, M. (2008). Synaptic theory of working memory. *Science* 319, 1543–1546.
 40. Manohar, S.G., Zokaei, N., Fallon, S.J., Vogels, T.P., and Husain, M. (2019). Neural mechanisms of attention to items in working memory. *Neurosci. Biobehav. Rev.* 101, 1–12.
 41. Rose, N.S., LaRocque, J.J., Riggall, A.C., Gosseries, O., Starrett, M.J., Meyering, E.E., and Postle, B.R. (2016). Reactivation of latent working memories with transcranial magnetic stimulation. *Science* 354, 1136–1139.
 42. Bays, P.M., Catalao, R.F.G., and Husain, M. (2009). The precision of visual working memory is set by allocation of a shared resource. *J. Vis.* 9.
 43. Suchow, J.W., Brady, T.F., Fougny, D., and Alvarez, G.A. (2013). Modeling visual working memory with the MemToolbox. *J. Vis.* 13, 9.
 44. Delorme, A., and Makeig, S. (2004). EEGLAB: An open-source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J. Neurosci. Methods* 134, 9–21.
 45. van Driel, J., Olivers, C.N.L., and Fahrenfort, J.J. (2021). High-pass filtering artifacts in multivariate classification of neural time series data. *J. Neurosci. Methods* 352, 109080.
 46. Nouri, A., and Ester, E.F. (2020). Recovery of information from latent memory stores decreases over time. *Cogn. Neurosci.* 11, 101–110.
 47. Maris, E., and Oostenveld, R. (2007). Nonparametric statistical testing of EEG and MEG data. *J. Neurosci. Methods* 164, 177–190.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Deposited data		
Preprocessed Behavioral and EEG Data	This paper	https://doi.org/10.18112/openneuro.ds004519.v1.0.1
Software and algorithms		
Stimulus presentation and analytic software	This paper	https://osf.io/wat6d/

RESOURCE AVAILABILITY

Lead contact

All questions and matters arising from this paper should be directed to and will be addressed by the lead contact, Dr. Edward F. Ester (ester@unr.edu).

Materials availability

This study did not generate any new reagents.

Data and code availability

- De-identified behavioral and preprocessed EEG Data (BIDS format) have been deposited on OpenNeuro and are publicly available as of the date of publication. Accession information is available in the [key resources table](#). Raw EEG data can be obtained by contacting the [lead contact](#) of this study.
- Original code sufficient to reproduce all figures and reported statistical values are publicly available on the Open Sciences Framework. Accession information is available in the [key resources table](#).
- Any additional information required to re-analyze the data reported in this paper is available from the [lead contact](#) upon request.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

In total, 42 human adult volunteers (ages 18–40) participated in this study, with each participant completing a single 2.5-h testing session. Two participants voluntarily withdrew from the study prior to completing both tasks (i.e., pro-cue vs. anti-cue); data from these participants were excluded from final analyses. Thus, the data reported here reflect the remaining 40 participants. Participants were recruited from the Florida Atlantic University (FAU) community via campus advertisements and remunerated at \$15/h in [Amazon.com](#) gift cards. All participants gave both written and oral informed consent prior to enrolling in the study, and all study procedures were approved by the FAU institutional review board (IRB). All participants self-reported normal or corrected-to-normal visual acuity. We had no *a priori* reason to suspect that task performance or study outcomes would vary as a function of sex, gender identity, race, ethnicity, or any other immutable characteristic; thus, we did not collect this information from participants.

METHOD DETAILS

Testing environment

Participants were seated in a dimly-lit and acoustically shielded recording chamber for the duration of the experiment. Stimuli were generated in MATLAB and rendered on a 17" Dell CRT monitor cycling at 85 Hz (1024 x 768 pixel resolution) via PsychToolbox3 software extensions. Participants were seated approximately 65 cm from the display (head position was unconstrained). To combat fatigue and/or boredom, participants were offered short breaks at the end of each experimental block.

Spatial retrocue task

A task schematic is shown in [Figure 1A](#). Each trial began with the presentation of an encoding display lasting 500 ms. The encoding display contained two colored circles (blue and red; subtending 1.75° visual angle from a viewing distance of 65 cm) rendered at two of eight locations (22.5° to 337.5° in 45°

increments) along the perimeter of an imaginary circle (radius 7.5° visual angle) centered on a circular fixation point (subtending 0.25°) rendered in the middle of the display. The locations of the two discs were counterbalanced across each task (i.e., pro-cue vs. anti-cue), though not necessarily within an experimental block. Participants were instructed to maintain fixation and refrain from blinking for the duration of each trial.

The sample display was followed by a 1.25 sec blank display and a 1.25 sec retrocue display. Retrocues were defined by a change in the color of the fixation point. During informative cue trials the fixation point changed colors from black to either blue or red (i.e., matching the color of a remembered disc) and remained that color for the duration of the trial. During uninformative cue trials the fixation point initially changed colors from black to purple (the “average” of blue and red), before again changing colors from purple to blue or red at the onset of the response display. At the end of the trial, a response display containing a fixation cue (i.e., a blue or red fixation point), a mouse cursor, a “?” symbol, and a response circle appeared. During the pro-cue task, participants were instructed to report the location of the disc matching the color of the fixation cue, while during the anti-cue task participants were instructed to report the location of the disc that did not match the color of the fixation cue. Participants responded by clicking along the perimeter of the response circle. Participants were instructed to prioritize accuracy over speed, and no response deadline was imposed. The trial terminated as soon as the participant clicked on a location. Sequential trials were followed by a 1.5-2.5 sec blank period (randomly and independently selected from a uniform distribution after each trial).

Each experimental block contained 28 informative cue and 28 uninformative cue trials, for a total number of 56 trials per block. Informative cue and uninformative cue trials were randomly intermixed within blocks. During the first half of the experiment (e.g., experimental blocks 1-8), each participant was assigned to the pro-cue or anti-cue task. Participants completed eight blocks in each of the pro- and anti-cue tasks. Task order (i.e., eight blocks of the pro-cue task followed by eight blocks of the anti-cue task or vice versa) was counterbalanced across participants.

Quantifying memory performance

We quantified participants’ memory performance as average absolute recall error (i.e., the difference in polar angle reported by the participant and the polar angle of the probed disk) and average response times. Comparisons of memory performance across task conditions were conducted via repeated-measures analyses of variance (ANOVA) and repeated-measures t-tests.

Importantly, average absolute recall errors can be influenced by the precision of participants’ memory as well as random guessing or accidental reports of a non-probed object (“swap errors”). To quantify the frequencies of random guessing and swap errors, we pooled participants’ recall data across all cue conditions (e.g., valid vs. invalid; 100% vs. 75%) and used a hierarchical Bayesian approach to fit participants’ data with a parametric model which assumes that on a given trial (a) participants report the position of the probed disk with precision k , (b) participants report the position of the non-probed disk with precision k (i.e., a “swap error”), or (c) participants randomly guess.⁴² We used hierarchical Bayesian modeling (implemented via the MemFit MATLAB toolbox⁴³) to obtain maximum a posteriori estimates of memory precision, guessing frequency, and swap error frequency at the single-participant and population levels.

EEG acquisition and preprocessing

Continuous EEG was recorded from 63 uniformly distributed scalp electrodes using a BrainProducts “actiCHamp” system. The horizontal and vertical electrooculogram (EOG) were recorded from bipolar electrode montages placed over the left and right canthi and above and below the right eye, respectively. Live EEG and EOG recordings were referenced to a 64th electrode placed over the right mastoid and digitized at 1 kHz. All data were later re-referenced to the algebraic mean of the left- and right mastoids, with 10-20 site TP9 serving as the left mastoid reference.

Data preprocessing was carried out via EEGLAB software extensions⁴⁴ and custom software. Data preprocessing steps included the following, in order: (1) resampling (from 1 kHz to 250 Hz), (2) bandpass filtering (1 to 50 Hz; zero-phase forward- and reverse finite impulse response filters as implemented by EEGLAB), (3) epoching from -1.0 to +5.0 sec relative to the start of each trial, (4) identification, removal, and interpolation of noisy electrodes via EEGLAB software extensions, and (5) identification and removal of oculomotor

artifacts via independent components analysis as implemented by EEGLAB. After preprocessing, location decoding analyses focused exclusively on the following 10-20 occipitoparietal electrodes: P7, P5, P3, Pz, P2, P4, P6, P8, PO7, PO3, POz, PO2, PO4, PO8, O1, O2, Oz.

Data cleanup

Prior to analyzing participants' behavioral or EEG data, we excluded all trials where the participant responded with a latency of < 0.4 s (we attributed these trials to accidental mouse clicks following the onset of the probe display rather than a deliberative recall of a specific stimulus position) and more than 3 standard deviations above the average response time across all experimental conditions. This resulted in an average (± 1 S.E.M.) loss of 14.43 ± 0.93 trials (or $1.67\% \pm 0.11\%$ of trials) across participants but had no qualitative effect on any of the findings reported here.

Decoding spatial positions from posterior alpha-band EEG signals

Location decoding was based on the multivariate distance between EEG activity patterns associated with memory for specific positions. This approach is an extension of earlier parametric decoding methods designed for use in circular feature spaces.²⁷ We extracted spatiotemporal patterns of alpha-band activity (8–13 Hz) from 17 occipitoparietal electrode sites (see [EEG acquisition and preprocessing](#) above). The raw timeseries at each electrode was bandpass filtered from 8-13 Hz (zero-phase forward-and-reverse filters as implemented by EEGLAB software), yielding a real-valued signal $f(t)$. The analytic representation of $f(t)$ was obtained via Hilbert transformation:

$$z(t) = f(t) + if(t)$$

where i is the imaginary operator and $if(t) = A(t)e^{i\phi(t)}$. Alpha power was computed by extracting and squaring the instantaneous amplitude $A(t)$ of the analytic signal $z(t)$.

Location decoding performance was computed separately for each disc (i.e., blue vs. red), trial type (i.e., informative vs. uninformative) and each task (i.e., pro-cue vs. anti-cue) on a timepoint-by-timepoint basis. In the first phase of the analysis, we sorted data from each condition into 5 unique training and test data sets using stratified sampling while ensuring that each training set was balanced across remembered positions (i.e., we ensured that each training data set contained an equal number of observations where the location of the remembered stimulus was at 22.5° , 67.5° , etc.). We circularly shifted the data in each training and test data set to a common center (0° , by convention) and computed trial-averaged patterns of responses associated with memory for each disc position in each training data set. Next, we computed the Mahalanobis distance between trial-wise activation patterns in each test data set with position-specific activation patterns in the corresponding test data set, yielding a location-wise set of distance estimates. If scalp activation patterns contain information about remembered positions then distance estimates should be smallest when comparing patterns associated with memory for identical positions in the training and test data set and largest when comparing opposite positions (i.e., those $\pm 180^\circ$ apart), yielding an inverted gaussian-shaped function. Trial-wise distance functions were averaged and sign-reversed for interpretability. Decoding performance was estimated by convolving timepoint-wise distance functions with a cosine function, yielding a metric where chance decoding performance is equal to 0. Decoding results from each training- and test-data set pair were averaged (thus ensuring the internal reliability of our approach), yielding a single decoding estimate per participant, timepoint, and task condition. To verify that our findings are not contingent on the specific type of decoding analysis used, we repeated the aforementioned analysis via eight-way support-vector-machine (SVM) based classification using a "one-versus-all" motif (see [results](#)).

Decoding and filter cutoffs

One recent study reported that high-pass filter cutoffs exceeding approximately 0.1 Hz can introduce temporal distortions in decoding timeseries in broadband EEG data.⁴⁵ As noted above, we applied a 1 Hz high-pass filter to the raw EEG data to optimize it for independent components analysis. Importantly, the high-pass filter was applied to the continuous EEG timeseries, that is, prior to sorting the data according to cue type (i.e., pro-vs. anti) or cue informativeness (i.e., informative vs. uninformative). An analogous procedure was used to isolate alpha-band activity within the broadband EEG signal prior to decoding. While we cannot exclude the possibility that our filtering approach introduced some amount of temporal smearing into decoding timeseries, our approach ensures that any filtering effects are agnostic to specific

experimental conditions. Thus, any differences in decoding timeseries across experimental conditions cannot be explained by mere filtering artifacts.

Cross-correlation analyses

Temporal differences in task-relevant location decoding performance were estimated via cross-correlation analyses. For uninformative trials, we extracted task-relevant decoding performance during pro- and anti-cue blocks over a period spanning 0.0 to 1.0 s following the onset of the probe display (i.e., when an informative cue instructed participants which disc to recall). We computed correlation coefficients between pro- and anti-cue decoding time series while systematically shifting the pro-cue time series from -1.0 to $+1.0$ s relative to the anti-cue decoding time series (e.g., Figure 3D, blue line). We compared these correlation coefficients to a distribution of correlation coefficients computed under the null hypothesis (i.e., no systematic difference in pro- and anti-cue decoding time series) by repeating the same analysis 10,000 times while randomizing the decoding condition labels (i.e., pro-vs. anti-cue) for each participant. An identical analysis was performed on task-relevant pro- and anti-cue decoding task performance from 0.0 to 1.75 s during informative trials. We deliberately selected a longer temporal interval for analysis during informative trials as we expected increases in pro- and anti-cue decoding performance to begin during the retrocue period and persist into the ensuing response period.

Quantifying frontal theta power

Analyses of frontal theta power focused on informative trials from the pro- and anti-cue tasks. The raw timeseries at each scalp electrode was bandpass filtered from 4-7 Hz (zero-phase forward-and-reverse filters as implemented by EEGLAB software), yielding a real-valued signal $f(t)$. The analytic representation of this signal was obtained via Hilbert transformation, and theta power was computed by extracting and squaring the instantaneous amplitude $A(t)$ of the analytic signal $z(t)$. Topographic maps of theta power during the pro- and anti-cue tasks were obtained by averaging power estimates over trials and a temporal window spanning 2.5 to 3.0 s following the start of each informative trial (i.e., 750 to 1000 ms after cue onset). Based on these maps, we limited further analyses to power estimates measured at four frontal electrode sites: AFz, Fz, F1, and F2. Data from these electrodes were used to compute time-resolved estimates of theta power during the pro- and anti-cue tasks and task differences in theta power. In a final analysis, we extracted and computed trial-wise estimates of theta power during the anti-cue task (using the same electrodes and temporal window described in the previous paragraph). We sorted participants' anti-cue EEG data into low- and high-theta power groups after applying a media split to theta power estimates, then decoded the location of the cue-matching but task-irrelevant item within each group. This allowed us to test whether evidence for exogenous selection of and/or "refreshing" of cue-matching memory traces was more likely to occur on trials with low- vs. high theta power.

N2pc analyses

The N2pc is a negative-going waveform defined by a greater negativity over occipitoparietal electrode sites contralateral vs. ipsilateral to the hemifield containing a visual target.²⁸ We used the N2pc to track covert spatial selection of the cue-matching and task-relevant position during pro-cue blocks and the cue-matching but task-irrelevant stimulus during anti-cue blocks. To control for sensory imbalances across visual hemifields, we restricted our analysis to trials where the task-relevant and task-irrelevant discs appeared in opposite visual hemifields. We estimated voltages over occipitoparietal electrode site pairs O1/2, PO3/4, and PO7/8 during trials when the task-relevant stimulus was in the left vs. right visual field, then sorted trial-wise voltage estimates by the hemifield containing the task-relevant target, i.e., contralateral vs. ipsilateral. We defined the N2pc as the average difference in voltage across contralateral and ipsilateral electrode sites over a period spanning 200–300 ms.

Inverted encoding model

To verify the generality of our findings across analytic approaches, we reconstructed position-specific WM representations from spatiotemporal patterns of alpha-band activity using an inverted encoding model. Our approach was conceptually and quantitatively identical to that used in earlier studies.^{25,26,46} We modeled alpha power at each scalp electrode as a weighted sum of eight location-selective channels, each with an idealized tuning curve (a half-wave rectified cosine function raised to the 8th power, with the maximum response of each function normalized to 1). The predicted responses of each channel during

each trial were arranged in a k channels by n trials design matrix C . The relationship between the EEG data and the predicted responses in C is given by a general linear model of the form:

$$B = WC + N$$

where B is an m electrode by n trial training data matrix, W is an m electrode by k channel weight matrix, and N is a matrix of residuals (i.e., noise).

To estimate W , we constructed a training data set containing an equal number of trials for each stimulus position (i.e., 22.5-337.5° in 45° increments). We identified the location φ with the fewest r repetitions and constructed a training dataset B_{trn} (m electrodes by n trials) and weight matrix C_{trn} (n trials by k channels) by randomly selecting (without replacement) 1 to r trials for each of the eight possible stimulus positions. The training dataset was used to compute a weight for each channel C_i using ordinary least-squares estimation:

$$V_i = \frac{\sum_i^{-1} W_i}{W_i^T \sum_i^{-1} W_i}$$

where T and -1 are the matrix transpose and inversion operations, respectively. Σ_i is the regularized noise covariance matrix for each channel i , estimated as:

$$\sum_i \frac{1}{n-1} \varepsilon_i \varepsilon_i^T$$

where n is the number of training trials and ε_i is a matrix of residuals:

$$\varepsilon_i = B_{trn} - W_i C_{trn,i}$$

Estimates of ε_i were obtained by regularization-based shrinkage using an analytically determined shrinkage observation. In this way, an optimal spatial filter v_i was estimated for each channel C_i , yielding an m electrode by k filter matrix V .

Next, we constructed a test dataset B_{tst} (m electrodes by n trials) containing data from all trials not included in the training data set and estimated trial-by-trial channel responses C_{tst} (k channels by n trials):

$$C_{tst} = V^T B_{tst}$$

Trial-wise channel response estimates were interpolated to 360°, circularly shifted to a common center (0°, by convention), and averaged, yielding a single reconstruction per participant, time point, cue condition (i.e., informative vs. uninformative) and task (i.e., pro vs. anti-cue). Condition-wise channel response functions were averaged, converted to polar form, and projected onto a vector with angle 0°:

$$r = |z| \cos(\arg(z)), z = c e^{2i\rho}$$

Where c is a vector of estimate channel responses and ρ is a vector of angles at which the channels peak. To ensure internal reliability, this entire analysis was repeated 100 times, and unique (randomly chosen) subsets of trials were used to define the training and test data sets during each iteration. The results were then averaged across permutations to obtain a single reconstruction strength estimate for each participant, task condition, and timepoint.

QUANTIFICATION AND STATISTICAL ANALYSIS

General

All statistical analyses were performed in MATLAB using custom software. Behavioral data were analyzed using repeated-measures parametric statistics (t-tests, ANOVA) with an *a priori* threshold of $p = 0.05$. EEG data were analyzed using nonparametric randomization tests with cluster-based corrections for autocorrelation in the EEG signal with an *a priori* threshold of $p = 0.05$. Where appropriate, test statistics (i.e., a t-score or F-ratio), p-values, and effect sizes are reported in the main text. Additional information on statistical quantification for behavioral and EEG data can be found in specific subsections below. Unless otherwise stated, n (and degrees of freedom) correspond to the number of participants in the analysis, i.e., sample size.

Statistical power

Before commencing data collection we determined that a sample of 40 volunteers would be sufficient to examine effects of interest. This estimate was based on sample sizes and effect sizes reported in prior studies that used similar experimental and analytic approaches.^{22,26,33}

Statistical comparisons – Behavioral data

Participants' behavioral data (i.e., absolute average recall error and average response time; [Figures 1B–1E](#)) were analyzed using standard repeated-measures parametric statistics (e.g., t-test, ANOVA); for these comparisons we report test statistics, p-values, and effect size estimates. For each test, we verified that critical assumptions (i.e., normality, equal variances) were met through visual inspection of the data.

Statistical comparisons – N2pc, decoding performance, and inverted encoding model

The decoding analysis and inverted encoding model we used assume chance-level performance of 0. Likewise, direct comparisons of decoding performance or reconstruction strength across conditions (e.g., pro-cue vs. anti-cue) assume null statistics of 0. Thus, we evaluated task-relevant and task-irrelevant decoding performance by generating null distributions of decoding performance (or differences in decoding performance across conditions) by randomly inverting the sign of each participant's data with 50% probability and averaging the data across participants. This procedure was repeated 10,000 times, yielding a 10,000-element null distribution for each time point (note that, under the central limit theorem, this distribution is guaranteed to be approximately normally distributed with a mean of 0). Finally, we implemented a cluster-based permutation test⁴⁷ with cluster-forming and cluster-size thresholds of $p < 0.05$ to correct for multiple comparisons across time points.