

Pseudomonas aeruginosa countrywide outbreak in hospitals linked to pre-moistened non-sterile washcloths, Norway, October 2021 to April 2022

Kirsten Gravningen¹, Oliver Kacelnik¹, Egil Lingaas², Torunn Pedersen³, Bjørn G Iversen¹, the *Pseudomonas* outbreak group⁴

1. Norwegian Institute of Public Health, Oslo, Norway

2. Department of Infection Prevention, Oslo University Hospital, Oslo, Norway

3. Norwegian National Advisory Unit on Detection of Antimicrobial Resistance, University Hospital of North Norway, Tromsø, Norway

4. The members of the group are listed under Collaborators

Correspondence: Kirsten Gravningen (kirstenmidtun.gravningen@fhi.no)

Citation style for this article:

Gravningen Kirsten, Kacelnik Oliver, Lingaas Egil, Pedersen Torunn, Iversen Bjørn G, the *Pseudomonas* outbreak group. *Pseudomonas aeruginosa* countrywide outbreak in hospitals linked to pre-moistened non-sterile washcloths, Norway, October 2021 to April 2022. *Euro Surveill.* 2022;27(18):pii=2200312. <https://doi.org/10.2807/1560-7917.ES.2022.27.18.2200312>

Article submitted on 08 Apr 2022 / accepted on 04 May 2022 / published on 05 May 2022

In November 2021, a clonal outbreak of *Pseudomonas aeruginosa* of novel sequence type ST3875 was detected in three patients who died of bloodstream infections in one hospital. By 25 April 2022, the outbreak included 339 cases from 38 hospitals across Norway. Initial hospital reports indicate *Pseudomonas* infection as the main contributing cause in seven deaths. In March 2022, the outbreak strain was identified in non-sterile pre-moistened disposable washcloths, used to clean patients, from three lots from the same international manufacturer.

On 19 November 2021, the Norwegian Institute of Public Health (NIPH) was notified by the University Hospital of North-Norway, Tromsø that three patients hospitalised with severe COVID-19 in the intensive care unit had died of bloodstream infections (BSIs) with indistinguishable *Pseudomonas aeruginosa* strains, only a few days apart. Whole genome sequencing (WGS) revealed a novel sequence type. Further cases were identified in several other hospitals thereafter; here we describe the outbreak and preliminary findings.

Timeline of the outbreak

Cases were rapidly detected in hospitals in all four health regions in Norway, strongly indicating a common source of infection [1-5]. *Pseudomonas aeruginosa* is an opportunistic pathogen associated with nosocomial infections that thrives in humid environments [6].

On 18 March 2022, Oslo University Hospital detected *P. aeruginosa* in disposable pre-moistened washcloths after systematic testing of several hundred different products that were used in hospital wards with identified cases. Particular attention was paid to moist


products that had not been terminally sterilised by the manufacturer such as liquid soaps, hand creams, toothpaste and lubricating gels for endoscopic procedures. Samples of the washcloths were cultivated by standard microbiological procedures [7]. On the same day, the hospital notified the NIPH and Norwegian Hospital Procurement Trust about the product type and lot number. The latter immediately alerted all hospitals and the Norwegian distributor that the hospitals should stop using these products. On 19 March, the *P. aeruginosa* isolates were confirmed as the outbreak strain ST3875 by clone-specific PCR and later through WGS. On 21 March, the detection of the probable source of infection was published on the NIPH website [8]. By 25 April, ST3875 had been detected by seven different hospital laboratories in 149 of the 577 washcloths tested from four lots produced on multiple dates [9].

The manufacturer notified their customers in Norway on 21 March. The Norwegian Food Safety Authority (NFSA) banned the contaminated lots from the Norwegian market on 23 and 30 March. The responsible person (RP) for the product in EU/European Economic Area (EEA) was contacted by the NFSA – through the Competent Authority in the country where the RP was based. On 4 April, NFSA notified the EU countries through the EU rapid alert system for unsafe consumer products (RAPEX).

The RP for a cosmetic product throughout the EU/EAA markets is a legal representative who is obliged to ensure that the product is safe and meets regulatory requirements and shall take immediate corrective action (recall or withdrawal) if necessary [10].

FIGURE 1

Timeline of investigation into *Pseudomonas aeruginosa* cases in hospitals, Norway, October 2021–April 2022 (n = 339)



NFSA: Norwegian Food Safety Authority; NIPH: Norwegian Institute of Public Health.

The manufacturer's initial laboratory test reports of the relevant product lots conveyed via the RP to NFSA were negative for *P. aeruginosa*. However, retesting of retained samples confirmed the presence of *P. aeruginosa* in the same lots. As part of the RP internal control, NFSA was able to identify additional product lots that had been incorrectly released by the manufacturer after *P. aeruginosa* was detected and also found discrepancies in other test reports. On 9 April, the NFSA released a warning in Norway against use of all products from this manufacturer.

On 7 April, the RP visited the manufacturer. On 14 April, the findings led to a voluntary recall of all lots of a total of 14 different products from one facility, issued to customers in the EU/EEA. The recall was done in order to ensure the highest level of consumer safety. This decision was supported by the Competent Authority which also notified EU countries through RAPEX.

The complete timeline of events is presented in Figure 1.

Setting

Cases included in the ongoing outbreak are from 15 October 2021 to 25 April 2022.

Outbreak investigation

A national outbreak group coordinated by the NIPH, including professionals from all health regions, was established on 20 January 2022. A confirmed case was defined as a hospitalised patient who was infected or colonised with *P. aeruginosa* ST3875.

The national outbreak group recommended that all hospitals should conduct targeted screening of patients in intensive care units and genotyping of *P. aeruginosa* strains detected in diagnostic and clinical samples. All samples were analysed in local hospital laboratories. In addition, most laboratories retrospectively genotyped *P. aeruginosa* in stored blood culture samples starting 1 January 2021. Initially, *P. aeruginosa* isolates were characterised using WGS or amplified fragment length polymorphism (AFLP) assay [11]. An in-house clone-specific real-time PCR was developed and validated by one hospital laboratory by week 8, 2022, and the PCR specifications were rapidly shared with other laboratories.

NIPH initiated two parallel lines of investigation. For the first line, each case was notified by the hospitals to the NIPH together with a comprehensive record of risk factors, exposures and clinical outcomes. The second line involved compiling a list of products purchased such as mouth swabs, hygiene products, urinary catheters, infusion sets and lidocaine gel during the outbreak period on the wards where the cases were cared for. However, the latter analyses revealed no indication of products that could be a possible source of the outbreak.

Demographic and clinical information

In total, 339 cases from 38 hospitals with a median age of 70 years (interquartile range: 59–78 plus one child) have been identified. The cases included in the outbreak register are described in the Table.

TABLE

Characteristics of cases, outbreak of *Pseudomonas aeruginosa* ST3875 infections in 38 hospitals across all four health regions in Norway, October 2021–April 2022 (n = 339)

Case characteristics	n	%
Sex		
Women	123	36.3
Men	216	63.7
Age (years)		
Median (IQR)	70 (59–78)	
Mean (SD)	67.5 (15.8)	
Health region^a		
Northern	30	8.8
Central	47	13.9
South-eastern	241	71.1
Western	21	6.2
Hospital ward^b		
Intensive care unit	129	38.1
General ward	164	48.4
Emergency ward	10	3.0
Other	36	10.6
Colonisation vs infection		
Colonisation only	65	19.2
Mild to moderate infection	190	56.0
Severe infection ^c	83	24.5
Missing information	1	0.3
<i>Pseudomonas</i> as cause of death^d (n=53)		
Strongly contributing	7	13.2
Possibly contributing	22	41.5
Not contributing	20	37.7
Unknown	4	7.6

IQR: interquartile range; SD: standard deviation.

^a The South-eastern health region is the largest region and provides specialist healthcare to 58% of the population in Norway.

^b Type of ward the patient was staying on when they had the first positive test for *P. aeruginosa* ST3875.

^c The severity of infection was assessed by a physician.

^d The role of *Pseudomonas* in the cause of death was assessed by a physician (the time between death and this assessment varied by patient). Final information is not available for all patients.

The presented information is based on data reported by hospital or infection control staff in each hospital.

Disease severity assessment and the cause of death review were done at the local level by a physician. Fifty-three deaths (16%) have so far been reported and for these, *Pseudomonas* infection was considered the main contributing cause of death in seven cases.

We included the first positive sample for each patient (n = 339). These samples were taken from diverse body sites indicating multiple routes of entry, including 16% blood cultures, 36% urine samples, 22% airway samples, 18% wound samples and 8% from other locations (Figure 2).

After use of the product was stopped between 18 and 20 March (week 11), we observed a rapid decline in

cases over the next 4 weeks until 25 April. However, there may be a reporting delay. Screening is ongoing and the hospitals will continue to report new cases to NIPH.

Genomic epidemiology

Genomic analyses were performed on quality-assessed Illumina-based assemblies by core genome phylogeny (Ridom SeqSphere+ version 8.3) based on 3,867 target genes [12]. Figure 3 illustrates the clonal relatedness of selected clinical strains from the four health regions and one selected washcloth isolate. The analysis showed an outbreak cluster with 0–4 allelic differences and most of the strains, including the one from the washcloth, present in the same node.

Discussion

Our preliminary investigation of the clonal *P. aeruginosa* ST3875 hospital outbreak in Norway revealed that contaminated non-sterile washcloths used for intensive care and other vulnerable patients were the source of the outbreak and have caused a number of infections, many of them severe, and contributed to at least seven deaths. Even though most cases were colonised or had mild urinary tract and wound infections there have been several severe infections including ventilator-associated pneumonia and BSIs among reported cases.


The clinical strains of *P. aeruginosa* ST3875 displayed a wild-type antimicrobial susceptibility profile [13]. However, resistance towards anti-pseudomonas drugs was discovered after antimicrobial treatment of single patients (data not shown). Thus, had it not been for the close proximity of the three initial BSI cases (time and place), the outbreak may have gone undetected for a longer period. The pre-moistened washcloth was distributed and used all over Norway, including in primary healthcare. According to the manufacturer’s website, this product is used in healthcare worldwide.

The washcloth is classified as a non-sterile cosmetic product and does not have to meet the standards for medical devices and medicines. However, every cosmetic manufacturer is responsible for the microbiological safety of its products to ensure that they have been produced under hygienic conditions. *P. aeruginosa* is one of four species (in addition to *Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*) for which cosmetic products should be tested and contamination excluded, according to the Commission Implementing Decision on guidelines on Annex I to Regulation (EC) No 1223/2009 on cosmetic products [10,14,15].

The use of this product is widespread but often not documented in patient records and thus difficult to investigate in traditional epidemiological studies. According to the hospital line lists, 40% of cases had been exposed to disposable washcloths, 6% had not, and 54% reported ‘unknown’ or had missing information. However, one hospital systematically tested

FIGURE 2

Distribution of *Pseudomonas aeruginosa* ST3875 cases by test date (week) for the first positive sample, by case and sample type, 38 hospitals, Norway, October 2021–April 2022 (n = 339)



One sample per patient is included in the figure. For patients who had a positive blood culture sample, this sample is included in the figure, even if the patient had an earlier positive sample from another body site (n = 4). Eleven blood cultures between weeks 41 and 51 were analysed retrospectively as part of the outbreak investigation.

Authors' contributions

KG, OK, EL, TP, BGI, AMA, PE, NOH, SBJ, KSK, NIK, JAK, SAK, PL, DHS, MNU and LEØ designed the outbreak investigation. KG, OK, EL, BGI, AMA, PE, NOH, SBJ, KSK, NIK, JAK, SAK, PL, DHS, MNU and LEØ implemented the investigation and collected data in collaboration with the hospitals and laboratories. EL, TP, AB, BCB, NOH, AI, JJ, SBJ, PL, IHL, MTN, ASu, DHS and Ask were responsible for the laboratory analysis. KG, OK, EL, TP, BGI, PE, GG, SBJ, JAK, BL, MNU and LEØ analysed and interpreted the data. KG and OK wrote the first draft of the manuscript. All authors revised the manuscript and approved the final version.

References

- Iversen BG, Jacobsen T, Eriksen HM, Bukholm G, Melby KK, Nygård K, et al. An outbreak of *Pseudomonas aeruginosa* infection caused by contaminated mouth swabs. *Clin Infect Dis*. 2007;44(6):794-801. <https://doi.org/10.1086/511644> PMID: 17304450
- Chittick P, Russo V, Sims M, Robinson-Dunn B, Oleszkowicz S, Sawarynski K, et al. An outbreak of *Pseudomonas aeruginosa* respiratory tract infections associated with intrinsically contaminated ultrasound transmission gel. *Infect Control Hosp Epidemiol*. 2013;34(8):850-3. <https://doi.org/10.1086/671268> PMID: 23838230
- Becks VE, Lorenzoni NM. *Pseudomonas aeruginosa* outbreak in a neonatal intensive care unit: a possible link to contaminated hand lotion. *Am J Infect Control*. 1995;23(6):396-8. [https://doi.org/10.1016/0196-6553\(95\)90272-4](https://doi.org/10.1016/0196-6553(95)90272-4) PMID: 882117
- Stephenson JR, Heard SR, Richards MA, Tabaqchali S. Gastrointestinal colonization and septicaemia with *Pseudomonas aeruginosa* due to contaminated thymol mouthwash in immunocompromised patients. *J Hosp Infect*. 1985;6(4):369-78. [https://doi.org/10.1016/0195-6701\(85\)90053-2](https://doi.org/10.1016/0195-6701(85)90053-2) PMID: 2868034
- Iversen BG, Eriksen HM, Bø G, Hagestad K, Jacobsen T, Engeset E, et al. *Pseudomonas aeruginosa* contamination of mouth swabs during production causing a major outbreak. *Ann Clin Microbiol Antimicrob*. 2007;6(1):3. <https://doi.org/10.1186/1476-0711-6-3> PMID: 17355630
- Kerr KG, Snelling AM. *Pseudomonas aeruginosa*: a formidable and ever-present adversary. *J Hosp Infect*. 2009;73(4):338-44. <https://doi.org/10.1016/j.jhin.2009.04.020> PMID: 19699552
- International Organization for Standardization (ISO). *Cosmetics – microbiology – identification of Pseudomonas aeruginosa (ISO 2217:2015)*. Geneva: ISO; 2021. Available from: <https://www.iso.org/standard/68312.html>
- The Norwegian Institute of Public Health (NIPH). Utbruddsbakterien funnet i engangs vaskeklut brukt i helsetjenesten. [Outbreak bacterium detected in disposable washcloth used in healthcare]. Oslo: NIPH; 2022. Norwegian. Available from: <https://www.fhi.no/nyheter/2022/utbruddsbakterien-funnet-i-engangs-vaskeklut-brukt-i-helsetjenesten>
- Norwegian Food Safety Authority (NFSA). Trekker flere engangsvaskekluter etter funn av ulovlige bakterier. [Withdrawal of several disposable washcloths after detection of illegal bacteria]. Brumunddal: NFSA; 2022. Norwegian. Available from: https://www.mattilsynet.no/kosmetikk/stoffer_i_kosmetikk/trekker_flere_engangsvaskekluter_etter_funn_av_ulovlige_bakterier.46361
- European Commission. Regulation (EC) No 1223/2009 of the European Parliament and of the Council of 30 November 2009 on cosmetic products. Official Journal of the European Union. 2009; L 342/59. Available from: <http://data.europa.eu/eli/reg/2009/1223/oj>
- Berdal JE, Smith-Erichsen N, Bjørnholt JV, Blomfeldt A, Bukholm G. Does *Pseudomonas aeruginosa* colonization influence morbidity and mortality in the intensive care unit patient? Experience from an outbreak caused by contaminated oral swabs. *Acta Anaesthesiol Scand*. 2006;50(9):1095-102. <https://doi.org/10.1111/j.1399-6576.2006.01044.x> PMID: 16987339
- Jünemann S, Sedlazeck FJ, Prior K, Albersmeier A, John U, Kalinowski J, et al. Updating benchtop sequencing performance comparison. *Nat Biotechnol*. 2013;31(4):294-6. <https://doi.org/10.1038/nbt.2522> PMID: 23563421
- Kahlmeter G, Turnidge J. How to: ECOFFs-the why, the how, and the don'ts of EUCAST epidemiological cutoff values. *Clin Microbiol Infect*. 2022;S1198-743X(22)00101-X. <https://doi.org/10.1016/j.cmi.2022.02.024> PMID: 35218980
- European Commission. Commission Implementing Decision of 25 November 2013 on Guidelines on Annex I to Regulation (EC) No 1223/2009 of the European Parliament and of the Council on cosmetic products. Official Journal of the European Union. 2013; L 315/82. Available from: https://eur-lex.europa.eu/eli/dec_impl/2013/674/oj
- Scientific Committee on Consumer Safety. The SCCS notes of guidance for the testing of cosmetic ingredients and their safety evaluation. 11th revision. SCCS/1628/21. Brussels: European Commission; 2021. Available from: https://ec.europa.eu/health/system/files/2021-04/sccs_o_250_o.pdf

License, supplementary material and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence and indicate if changes were made.

Any supplementary material referenced in the article can be found in the online version.

This article is copyright of the authors or their affiliated institutions, 2022.