

Utilization Outcomes of a Pilot Primary Care Team Redesign

Health Services Research and
Managerial Epidemiology
Volume 5: 1-6
© The Author(s) 2018
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2333392818789844
journals.sagepub.com/home/hme



Alison R. Landrey¹, Valerie S. Harder², Marie B. Sandoval¹,
John G. King³, David S. Ziegelman¹, and Charles D. MacLean¹

Abstract

Objectives: To evaluate the effect of a team-based primary care redesign on primary care, emergency department (ED) and urgent care (UC) utilization, and new patient access to primary care.

Study Design: A retrospective pre–post difference-in-differences analysis of utilization outcomes for patients on a redesigned primary care team compared to a standard primary care group.

Methods: Within a patient-centered medical home, a pilot team was developed comprising 2 colocated “teamlets” of 1 physician, 1 nurse practitioner (NP), 1 registered nurse (RN), and 2 licensed practical nurses (LPNs). The redesigned team utilized physician–NP comanagement, expanded roles for RNs and LPNs, and dedicated provider time for telephone and e-mail medicine. We compared changes in number of office, ED, and UC visits during the implementation year for patients on the redesigned team compared to patients receiving the standard of care in the same clinic. Proportion of new patient visits was also compared between the pilot and the control groups.

Results: There were no differences between the redesign group and control group in per-patient mean change in office visits ($\Delta = -0.04$ visits vs $\Delta = -0.07$; $P = .98$), ED visits ($\Delta = 0.00$ vs $\Delta = 0.01$; $P = .25$), or UC visits ($\Delta = 0.00$ vs $\Delta = 0.05$; $P = .08$). Proportion of new patient visits was higher in the pilot group during the intervention year compared to the control group (6.6% vs 3.9%; $P < .0001$).

Conclusions: The redesign did not significantly impact ED, UC, or primary care utilization within 1 year of follow-up. It did improve access for new patients.

Keywords

emergency visits, practice management, program evaluation, access to care, primary care

Background

Over the past decade, health-care delivery organizations, insurers, and specialty organizations have supported the transition of primary care practices toward the Patient-Centered Medical Home (PCMH) model to address the triple aim of health-care reform: improving quality of care and patient experience while controlling health-care costs.^{1,2,3-7} However, effects of PCMH models have been mixed.⁷⁻¹¹ Shortages of primary care physicians, patient complexity, and competing demands on primary care present growing challenges to the goals of the medical home model.^{2,5} Team-based care is a central feature of the PCMH with potential to address some of these challenges, yet team composition and implementation are variable among medical homes.^{2,10,12-14} Current proposals for primary care redesign call for further work in team-based care and emphasize

expanded use of non-face-to-face visits.^{2,15,16} Although lack of reimbursement for such encounters has been a barrier to innovation in these domains, as health policy and reimbursement shift

¹ Department of Medicine, University of Vermont Medical Center, Burlington, VT, USA

² Departments of Pediatrics and Psychiatry, University of Vermont Medical Center, Burlington, VT, USA

³ Department of Family Medicine, University of Vermont Medical Center, Burlington, VT, USA

Submitted June 26, 2018. Accepted June 26, 2018.

Corresponding Author:

Alison R. Landrey, Department of Medicine, University of Vermont Medical Center, 1 Timber Lane, South Burlington, VT 05403, USA.

Emails: alison.landrey@uvmhealth.org, alison.landrey@gmail.com






Figure 1. Usual care model compared with pilot teamlet-based care model.

toward value-based payment, care models that utilize teams and non-face-to-face visits may be well positioned to address ongoing gaps in access and care.¹⁷⁻²⁰

Concentrating care in a smaller care unit, or “teamlet,” within a larger practice can enhance continuity and improve experience for both providers and patients.¹⁴ Collaboration between a physician and an advanced practice provider (APP) in shared care of a panel of patients is an example of a teamlet that has demonstrated improved quality measures and provider and patient satisfaction.^{21,22} Expanding non-face-to-face access within such teams with established provider–patient relationships may further enhance care delivery.¹⁵ Specifically, use of telephone and online encounters to meet patients’ needs can improve access, efficiency, continuity, and timeliness of visits.²³⁻²⁵

We piloted a teamlet-based primary care model incorporating enhanced non-face-to-face access. The goal of this study was to determine the effect of this pilot on primary care, emergency department (ED), and urgent care (UC) utilization. Because enhanced access and care continuity are associated with decreased ED utilization, we hypothesized that in comparison to a control group within the same clinic, patients attributed to the pilot providers would have decreased ED and UC utilization.^{3,26} A secondary goal of the study was to determine the effect of the pilot on access for new patients.

Methods

Study Setting

The study was conducted at a National Committee for Quality Assurance level III academic PCMH²⁷ that cares for approximately 12 000 adult patients. The practice was closed to new patients, and patients’ ability to see their personal primary care physician for acute or timely follow-up care was often limited; patients were often scheduled with any available provider for acute visits. Providers had no dedicated clinic time for non-face-to-face visits. The majority of the APPs’ clinic time was

spent seeing patients for acute care, preoperative, or preventive care rather than chronic disease management. There were 7 registered nurses (RNs) dedicated to telephone triage. Seven licensed practical nurses (LPNs) and 4 medical assistants (MAs) were paired with providers to perform visit preparation, rooming duties, and panel management. The LPNs and MAs were termed clinical care associates (CCAs). The RNs performed face-to-face patient visits rarely, and LPN visits were limited to vaccine and medication administration.

Intervention

A team-based care pilot was developed within the PCMH with a mission to improve new patient access, care value, and patient access to their personal care team. The pilot was implemented in May 2015 and comprised of 2 teamlets of 5 providers (an MD, an APP, an RN and 2 CCAs, including at least 1 LPN; Figure 1). The team was colocated in 1 hallway, with each MD-APP pair sharing office space. Team meetings occurred 1 to 3 times weekly to coordinate care and iterate process work.^{28,29}

Physicians were given 30 minutes and APPs 90 minutes daily for non-face-to-face patient care. Phone staff were instructed to direct all pilot patient calls for acute concerns to the pilot RN rather than schedule directly with providers and were instructed not to schedule nonpilot patients with pilot providers and vice versa. Efforts were made to accommodate pilot patients for acute visits within the patient’s teamlet rather than the alternate teamlet. Over the first 6 months, a chronic disease comanagement model within the MD-NP pairings was developed iteratively; patients with chronic medical conditions who were seen every 1 to 6 months were scheduled alternately with the APP and the MD every other visit.

Nonprovider staff on the pilot team had expanded roles in managing chronic illness and coordinating care. The CCAs reviewed after visit instructions and checked patients out in the examination room rather than patients checking out at the front desk. Beginning in June 2015, outreach calls to patients who had

Table 1. Patient Demographics for Pilot Group Versus Control Group.

Patient Demographics	Pilot Group, N = 1807	Control Group, N = 4715	P Value
Age, mean (SD)	57.7 (16.2)	59.5 (15.6)	.066
Sex, % female	55.7	58.8	.024
Hypertension, % yes	41.7	40.6	.437
Diabetes, % yes	12.1	10.9	.169
Current smoker, % yes	7.1	6.8	.646
Anxiety/depression, % yes	40.2	34.0	<.001
Insurance type, %			
Commercial	69.2	71.5	.287
Medicare	21.0	19.8	
Medicaid	6.1	5.3	
Other	3.7	3.4	

recently been to the university hospital ED or UC were made by either the teamlet APP or the LPN. The LPN and RN blood pressure, diabetes, and medication reconciliation face-to-face visits began in July 2015, with protocols refined iteratively.

During pilot implementation, the nonpilot providers and staff were aware of the abovementioned interventions but did not participate, thus serving as a control for this study. The APPs on the pilot team and one physician in the control provider group were listed as accepting new patients during the pilot implementation period.

Data Collection and Statistical Analysis

We conducted a retrospective review of electronic health record (EHR) data for all patients attributed to the clinic who had at least 1 visit to the clinic during both a baseline year (May 2014 to April 2015) and the intervention year (May 2015 to April 2016). Patients were attributed to either pilot providers or control providers based on their EHR primary care provider assignment on May 1, 2014. Billing data from visits to the clinic and to the university ED and UC center were collected, all of which used the same EHR. *International Classification of Diseases, Tenth Revision (ICD-10)* codes for common chronic medical conditions were obtained from patient problem lists for baseline characteristics.

We compared demographic characteristics of patients within the pilot group to the control group using simple χ^2 and t tests (Table 1). Difference-in-differences regression analyses were used to compare changes over the 2 time periods for 3 outcomes: number of patient visits to the ED, UC, and clinic, between the pilot team and the control group (Table 2).³⁰ The following model (Equation 1) was fit for each outcome ($Y_{i,t}$):

$$Y_{i,t} = \beta_0 + \beta_1(\text{redesign}_i) + \beta_2(\text{time}_t) + \beta_3(\text{redesign}_i \times \text{time}_t) + e_{i,t}. \quad (1)$$

The patient-level regression models (Equation 1) for each outcome ($Y_{i,t}$) used a difference-in-differences specification to examine the impact of care redesign, where i was an indicator for pilot redesign, and t was an indicator for time period. Each regression also included demographics (age, sex, current smoker, and insurance type in 2014) and active problem list diagnoses (hypertension, diabetes, and anxiety/depression) from the EHR as potential confounders. Due to excess zeros in number of visits (count) data, we used the zero-inflated Poisson (ZIP) regression models.³¹ The ZIP regression model included 2 parts: a Poisson count model and the logit model for predicting excess zeros. All ZIP models were run with the robust option in Stata statistical software version 15.³¹ The ZIP models were compared to standard Poisson models; if there was not a better model fit using ZIP, standard Poisson was used. We used χ^2 analysis to evaluate differences in proportion of new patient visits between the pilot and control groups. This study was approved by the institutional review board.

Results

There were no differences in baseline characteristics between the pilot group ($n = 1807$ patients) and control group ($n = 4715$ patients) with respect to age, insurance type, smoking status, or prevalence of hypertension or diabetes. The pilot group had a significantly smaller proportion of females and a higher proportion of patients with anxiety or depression (Table 1). We found no differences between the pilot group and the control group in per-patient mean change in office visits ($\Delta = -0.04$ visits vs $\Delta = -0.07$; $P = .98$), ED visits ($\Delta = 0.00$ vs $\Delta = 0.01$; $P = .25$), or UC visits ($\Delta = 0.00$ vs $\Delta = 0.05$; $P = .08$; Table 2). There was no difference in baseline proportion of new patient visit between the pilot ($191/6578 = 2.9\%$) and control groups ($507/16962 = 3.0\%$; $P = .729$). The pilot group had a higher proportion of new patient visits ($437/6628 = 6.6\%$) during the intervention period compared to the control group ($661/17063 = 3.9\%$; $P < .001$).

Discussion

Our study adds to the growing body of literature evaluating the impact of primary care redesign efforts. We found that the primary care team pilot did not significantly impact ED, UC, or primary care utilization within 1 year of follow-up when compared to a control group. The pilot team did increase access for new patients, which demonstrates the ability of care teams with expanded nonphysician roles to accommodate increases in panel size. Although we cannot fully separate the effects of the intervention from confounding secular or unmeasured effects, we are confident that this practice redesign was the dominant practice innovation throughout the study.

There are several limitations to this study. Given the non-randomized application of the intervention, there may be unmeasured ways in which the study and control groups

Table 2. Unadjusted Mean Annual Per-Patient Visit Counts Pre- and Postintervention for Pilot Group Versus Control Group.

Visit Counts	Pilot Group Preintervention	Pilot Group Postintervention	Control Group Preintervention	Control Group Postintervention	P Value ^a
Primary care office visits, mean (SD)	2.92 (2.61)	2.86 (2.55)	2.91 (2.45)	2.84 (2.34)	.981
Emergency department visits, mean (SD)	0.141 (0.415)	0.139 (0.428)	0.111 (0.387)	0.123 (0.386)	.251
Urgent care walk in visits, mean (SD)	0.277 (0.757)	0.282 (0.749)	0.253 (0.741)	0.300 (0.844)	.08

^aP value for the difference-in-differences effect between the pilot and control groups from pre to post intervention from Poisson regression models controlling for age, sex, hypertension, diabetes, current smoker, anxiety/depression, and insurance type as potential confounders.

differed. Since the intervention took place within the same clinic as the control group, it is also possible that some of the pilot methods were adopted less formally by members of the control group.³ We were not able to assess the integrity of some aspects of the intervention; for example, we could not quantify non-face-to-face visits in our EHR data. Further, as a pilot study in a single practice, the results may not generalize to other sites.

Given the iterative nature of the pilot team's work, a 1-year duration may not have been enough to observe significant differences in care value.^{32,33} The trend toward fewer UC visits among patients attributed to the pilot team compared with control patients during the implementation year suggests that with longer follow-up, a significant difference may have been found. The small sample size may have further limited our ability to detect differences in utilization between the groups. Two larger studies of pilot medical home redesigns with follow-up over 2 to 4 years did demonstrate improvement in ED and UC utilization.³⁴ Further, the baseline characteristics of our study showed higher rates of anxiety and depression in patients attributed to the pilot group, which could have affected utilization.^{35,36} Additionally, baseline ED and UC rates were low, limiting ability to demonstrate improvement.³⁷⁻³⁹ We were only able to collect ED and UC utilization data from the university hospital; although this methodology was the same during both the baseline and the intervention year and the next closest hospital is 27 miles away, the true ED and UC utilization data are underrepresented in our study.

Other quality measures such as glucose and blood pressure control were not evaluated in our study and could have added meaningful outcome data, although team care innovations showing improvement in these areas usually focus on a specific chronic disease.^{20,22} Our pilot study is more reflective of the mixed outcomes often seen in multifaceted care delivery innovations.^{9,32} We did not design a method for evaluating patient satisfaction with the pilot team model compared to the nonpilot team model, which may have been informative. A qualitative evaluation of a measure of the fourth aim of health-care reform, provider and staff experience with the redesign, was completed and analysis is underway.⁴⁰ Finally, a cost analysis of the redesign was beyond the scope of this study; however, the component of the model with the greatest cost impact is the increased provider time for non-face-to-face visits. This component may only be feasible with alternative payment structures.

Despite these limitations, ED/UC utilization in primary care redesign efforts is an important outcome, and our study adds insight into the difficulties in showing improvements within complex and dynamic practice environments. Future studies of primary care redesign efforts could explore longer follow-up, additional care value measures, patient satisfaction, and focused evaluation of single process improvement cycles.

Conclusion

The pilot primary care practice redesign did not significantly impact ED, UC, or primary care utilization within 1 year of follow-up. The redesign team did accommodate more new patient visits during the implementation year. Our study adds insight into the challenges of demonstrating broad utilization benefits of complex system redesign.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This was a pilot study supported by the University of Vermont Medical Center (UVMHC) in that, in contrast to the standard business model at UVMHC, pilot care providers were not held accountable to any RVU targets during the pilot implementation year.

References

- Berwick DM, Nolan TW, Whittington J. The triple aim: care, health, and cost. *Health Aff (Millwood)*. 2008;27(3):759-769.
- Bodenheimer T, Pham HH. Primary care: current problems and proposed solutions. *Health Aff (Millwood)*. 2010;29(5):799-805.
- Werner RM, Canamucio A, Marcus SC, Terwiesch C. Primary care access and emergency room use among older veterans. *J Gen Intern Med*. 2014;29(suppl 2):S689-S694.
- Hefner JL, Wexler R, McAlearney AS. Primary care access barriers as reported by nonurgent emergency department users: implications for the US primary care infrastructure. *Am J Med Qual*. 2015;30(2):135-140.
- Stange KC, Nutting PA, Miller WL, et al. Defining and measuring the patient-centered medical home. *J Gen Intern Med*. 2010; 25(6):601-612.
- Reddy A, Pollack CE, Asch DA, Canamucio A, Werner RM. The effect of primary care provider turnover on patient experience of

- care and ambulatory quality of care. *JAMA Intern Med.* 2015; 175(7):1157-1162.
7. Rosland AM, Krein SL, Kim HM, et al. Measuring patient-centered medical home access and continuity in clinics with part-time clinicians. *Am J Manag Care.* 2015;21(5):e320-e328.
 8. Friedberg MW, Schneider EC, Rosenthal MB, Volpp KG, Werner RM. Association between participation in a multipayer medical home intervention and changes in quality, utilization, and costs of care. *JAMA.* 2014;311(8):815-825.
 9. Baron RJ. Not all (medical) homes are built alike: some work better than others. *JAMA Intern Med.* 2014;174(8):1358-1359.
 10. Porter ME, Pabo EA, Lee TH. Redesigning primary care: a strategic vision to improve value by organizing around patients' needs. *Health Aff (Millwood).* 2013;32(3):516-525.
 11. Hearld LR, Weech-Maldonado R, Asagbra OE. Variations in patient-centered medical home capacity: a linear growth curve analysis. *Med Care Res Rev.* 2013;70(6):597-620.
 12. Bodenheimer TS, Smith MD. Primary care: proposed solutions to the physician shortage without training more physicians. *Health Aff (Millwood).* 2013;32(11):1881-1886.
 13. Kletke PR, Emmons DW, Gillis KD. Current trends in physicians' practice arrangements. From owners to employees. *JAMA.* 1996; 276(7):555-560.
 14. Bodenheimer T, Willard-Grace R. Teamlets in primary care: enhancing the patient and clinician experience. *J Am Board Fam Med.* 2016;29(1):135-138.
 15. Berry LL, Seiders K, Wilder SS. Innovations in access to care: a patient-centered approach. *Ann Intern Med.* 2003;139(7): 568-574.
 16. Kaplan GS. Health care scheduling and access: a report from the IOM. *JAMA.* 2015;314(14):1449-1450.
 17. Institute of Medicine Committee on Quality of Health Care in A. *Crossing the Quality Chasm: A New Health System for the 21st Century.* Washington, DC: National Academies Press (US). Copyright 2001 by the National Academy of Sciences. All rights reserved; 2001.
 18. Porter ME. A strategy for health care reform—toward a value-based system. *N Engl J Med.* 2009;361(2):109-112.
 19. Wagner EH, Flinter M, Hsu C, et al. Effective team-based primary care: observations from innovative practices. *BMC Fam Pract.* 2017;18(1):13.
 20. Proia KK, Thota AB, Njie GJ, et al. Team-based care and improved blood pressure control: a community guide systematic review. *Am J Prev Med.* 2014;47(1):86-99.
 21. Jones MG, DeCherrie LV, Meah YS, et al. Using nurse practitioner comanagement to reduce hospitalizations and readmissions within a home-based primary care program. *J Healthc Qual.* 2016;39(5):249-258.
 22. Litaker D, Mion L, Planavsky L, Kippes C, Mehta N, Frolkis J. Physician–nurse practitioner teams in chronic disease management: the impact on costs, clinical effectiveness, and patients' perception of care. *J Interprof Care.* 2003;17(3):223-237.
 23. Adamson SC, Bachman JW. Pilot study of providing online care in a primary care setting. *Mayo Clin Proc.* 2010;85(8): 704-710.
 24. Scherger JE. E-mail “visits” can save you time. *Med Econ.* 2004; 81(18):37-38, 41.
 25. Murray M, Bodenheimer T, Rittenhouse D, Grumbach K. Improving timely access to primary care: case studies of the advanced access model. *JAMA.* 2003;289(8):1042-1046.
 26. Christakis DA, Wright JA, Koepsell TD, Emerson S, Connell FA. Is greater continuity of care associated with less emergency department utilization? *Pediatrics.* 1999;103(4 pt 1): 738-742.
 27. Accountable Care Organization Accreditation Levels. 2018. <http://www.ncqa.org/programs/accreditation/accountable-care-organization-aco/aco-accreditation-levels>. Accessed June 19, 2018.
 28. Rodriguez HP, Meredith LS, Hamilton AB, Yano EM, Rubenstein LV. Huddle up!: the adoption and use of structured team communication for VA medical home implementation. *Health Care Manage Rev.* 2015;40(4):286-299.
 29. Rosland AM, Nelson K, Sun H, et al. The patient-centered medical home in the veterans health administration. *Am J Manag Care.* 2013;19(7):e263-272.
 30. Ashenfelter O, Card D. Using the longitudinal structure of earnings to estimate the effect of training programs. *Rev Econ Stat.* 1985;67(4):648-660.
 31. Lambert D. Zero-inflated poisson regression, with an application to defects in manufacturing. *Technometrics.* 1992;34(1):1-14.
 32. Crabtree BF, Chase SM, Wise CG, et al. Evaluation of patient centered medical home practice transformation initiatives. *Med Care.* 2011;49(1):10-6.
 33. Bohmer RM. The hard work of health care transformation. *N Engl J Med.* 2016;375(8):709-711.
 34. Rosenthal MB, Friedberg MW, Singer SJ, Eastman D, Li Z, Schneider EC. Effect of a multipayer patient-centered medical home on health care utilization and quality: the Rhode Island Chronic Care Sustainability Initiative Pilot Program. *JAMA Intern Med.* 2013;173(20):1907-1913.
 35. Goodman RM. Emergency department use associated with primary care office management. *Am J Manag Care.* 2013;19(5): e185-e196.
 36. Blonigen DM, Macia KS, Bi X, Suarez P, Manfredi L, Wagner TH. Factors associated with emergency department use among veteran psychiatric patients. *Psychiatr Q.* 2017;88(4): 721-732.
 37. Morgan SR, Chang AM, Alqatari M, Pines JM. Non-emergency department interventions to reduce ED utilization: a systematic review. *Acad Emerg Med.* 2013;20(10):969-985.
 38. Xu T, Klein EY, Zhou M, Lowenthal J, Sharfstein JM, Peterson SM. Emergency department utilization among the uninsured during insurance expansion in Maryland. *Ann Emerg Med.* 2018. pii: S0196-0644(18)30371-8.
 39. National Center for Health Statistics: Emergency department visits. 2015. <https://www.cdc.gov/nchs/fastats/emergency-department.htm>. Accessed June 19, 2018.
 40. Bodenheimer T, Sinsky C. From triple to quadruple aim: care of the patient requires care of the provider. *Ann Fam Med.* 2014; 12(6):573-576.

Author Biographies

Alison R. Landrey, MD, is an assistant professor of medicine at the Robert Larner, MD College of Medicine at the University of Vermont and she practices general internal medicine at the Adult Primary Care South Burlington practice at the University of Vermont Medical Center.

Valerie S. Harder, PhD, MHS is a biostatistician and epidemiologist. She is an associate professor of psychiatry and pediatrics and is the director of the health services research team at the Vermont Child Health Improvement Program.

Marie B. Sandoval, MD is an associate professor of medicine at the Robert Larner, MD College of Medicine and is the medical director at the Adult Primary Care South Burlington practice at the University of Vermont Medical Center, where she practices general internal medicine.

John G. King, MD is an associate professor and vice chair of family medicine at the Robert Larner, MD College of Medicine. He is also the family medicine residency program director and practices family medicine at the Milton Family Practice at the University of Vermont Medical Center.

David S. Ziegelman, MD is an assistant professor of medicine at the Robert Larner, MD College of Medicine and he practices general internal medicine at the Adult Primary Care South Burlington practice at the University of Vermont Medical Center.

Charles D. MacLean, MDMC is a professor of medicine and associate dean for Primary Care at the Robert Larner, MD College of Medicine at the University of Vermont. He practices general internal medicine at the Adult Primary Care Essex practice at the University of Vermont Medical Center.