

Original Research

Acute Effects of Dry Needling on Myofascial Trigger Points in the Triceps Surae of Ballet Dancers: A Pilot Randomized Controlled Trial

Jennifer A Janowski, DScPT, OCS, FAAOMPT^{1, a}, Deanna M L Phelan-Smith, DPT², Maria N Kroat Brady, DPT², Kelsey L Michels, DPT², Alexandra H Timm, DPT, OCS², Nicole M Boucher, DPT², Kedron D Casteen, BFA², David Village, PT, DHSc¹, Mark D Sleeper, PT, PhD²

¹ Andrews University, ² Northwestern University

Keywords: movement system, temperature, range of motion, force, dry needling, calves

<https://doi.org/10.26603/001c.21475>

International Journal of Sports Physical Therapy

Vol. 16, Issue 2, 2021

Background

There is convincing evidence that dancers suffer injuries to the triceps surae musculature. Research on the immediate effects of dry needling (DN) is limited, and it is important to understand the acute effects of this treatment prior to performance.

Purpose

The purpose of this pilot study was to assess the immediate effects of DN on myofascial trigger points in terms of skin surface temperature, pain, active and passive range of motion, and torque production in the triceps surae of ballet dancers.

Study Design

Randomized, double-blinded pilot study

Methods

Professional ballet dancers that fit inclusion and exclusion criteria (n=11) were randomly assigned to an experimental or control group. The dancers had three pre-determined standard point (SP) measurement spots that were used as a baseline for surface temperature comparisons. The dancers were also palpated for trigger point (TP) spots. Both SP and TP spots were marked for future measurements. The experimental group received DN, while the control group received sham DN (SHAM) to their bilateral calves at the TP spots. Immediately prior to and following treatment, both DN and SHAM groups were tested for skin surface temperature, pain, range of motion, and plantar flexion torque by blinded assessors. Paired t-tests and independent t-tests were performed to examine for differences between groups.

Results

The surface temperature for the TP was higher than the SP measurements prior to intervention (Right calf p= .014; Left calf p= .031). There were no significant changes in VAS scale reported pain and ROM. The plantar flexion torque measurements showed an increase in the DN group of the left calf at the angular velocity of 60 degrees/sec.

Conclusion

This was a unique pilot study examining the acute effects of DN on professional ballet dancers. The results were limited due to low sample size. However, the methodology for

a Corresponding Author:

Jennifer A. Janowski
Athletico Gold Coast
24 E. Chicago Ave
Chicago, IL 60611
jenniferjanowski@yahoo.com

this study and surface temperature results invites future research.

Level of evidence

Level 1b

INTRODUCTION

Dancers often suffer from decreased ability to perform relevé, plié, and jumping due to deficits at the ankle joint.^{1,2} Calf pain from overuse, prevalent among professional ballet dancers, creates these limitations, which prevents the dancer from performing at their full potential.³⁻⁵ Research suggests dry needling (DN) may be effective for immediately decreasing musculoskeletal pain,⁶⁻¹⁷ but the acute results of DN on range of motion (ROM) have been mixed.^{7,8,10,12,14,16,18,19} Currently, evidence regarding the immediate results of DN on gastrocnemius and soleus function is limited.¹⁴ Additionally, only a small number of studies have demonstrated the instant effects of DN on force production.^{7,20-22} There are no studies on DN and surface temperature of the specific muscle treated.

Myofascial pain occurs in patients with MTrPs, defined as a point within a taut skeletal muscle exhibiting an increased response to a stimulus.²³ One hypothesis for the cause of MTrPs is that acetylcholine (ACh) is released excessively at the neuromuscular junction, creating an area of tautness within the muscle.²³ This may cause unremitting stimulation of the sarcomere and may lead to hypoxia.²³⁻²⁵ DN involves the insertion of a thin, filiform needle into MTrPs that elicit a local twitch response affecting the ACh at the neuromuscular junction.²³ DN is also thought to impact blood flow by causing the release of vasoactive substances, which promote vasodilation and increase oxygenation similar to deep tissue massage.^{23,26,27}

Even though researchers have documented increases in blood flow and oxygen saturation to the tissue after DN, the question remains if DN affects the surface temperature at the tissue being treated.^{23,26,28,29} One study demonstrated an increased temperature reading at an acupuncture point after the needle was in this point for 10 minutes.³⁰ Recent research has shown that DN provokes intense vasodilation and temperature increases in measured referral points to the muscle treated.³¹ However, there are no studies that assess the pre- and post-surface temperature of the specific muscle being treated with DN.

Meta-analyses conducted on pain scores indicate that there is a statistically significant improvement in pain ratings immediately after DN.^{9,32} DN has also been shown to reduce pain faster than analgesics following total knee arthroplasty.¹⁵ Instant improvement on pain pressure threshold in the masseter muscle compared to placebo, decreased pain sensitivity in 57 shoulder patients, and a decrease in cervical muscular pain have been demonstrated after DN.^{6,8,12,13} Patients with myofascial pain syndrome demonstrated improvements on the Visual Analog Scale (VAS) after DN vs. sham needles immediately and at four weeks post treatment.¹⁷

Other authors have explored the effects of DN on pain in athletes such as rugby and volleyball.^{10,14,16,18} Mason, Tansey and Westrick reported decreased knee pain after DN in a case report on a dancer with posterior knee pain.¹⁴

These results on athletes might indicate DN as an effective means to treat ballet dancers' calf pain immediately, but DN's impact on performance ability remains unclear.

Amongst dancers, the effect of DN on ROM is especially important. Since one-fourth of injuries in ballet dancers involve the ankle joint, improving dorsiflexion ROM is key to prevention and treatment of acute and overuse injuries.⁴ Adequate triceps surae flexibility has demonstrated increased dorsiflexion ROM, increased force production by improving the ability to generate elastic energy, and decreased excessive pronation upon weight acceptance.^{4,33} In addition, increased dorsiflexion when landing jumps has been suggested to increase ankle plantar flexor pre-stretch, which may improve the utilization of elastic energy and enhance jump function.³⁴ Therefore, improving dorsiflexion ROM of the triceps surae may be important in injury prevention and dancer performance.

Fernandez-De-Las-Penas describes trigger points as taut bands which limit joint mobility.³⁵ Grieve et al^{36,37} performed studies on manual trigger point release to the triceps surae musculature and reported positive results in regards to ROM after intervention. Eftekharsadat, Babaei-Ghazani, and Zeinolabedinzadeh³⁸ performed DN in patients with heel pain and demonstrated no effect in ankle ROM. However, the methodology of this study was different because it pre-determined points in the calf, did not manipulate the needle, and the measurements were taken at 4 weeks post-intervention.³⁸ Additional authors have found immediate increases in shoulder ROM,^{16,39} cervical ROM,^{7,12,18,40} and jaw opening⁸ after DN.

Due to the possibility of increased ROM, DN may impact force and torque production immediately after treatment. Behm and Chaouachi¹ have shown that increased ROM is associated with decreased force production acutely.¹ Prolonged static stretching, which also increases ROM, has been shown to decrease voluntary peak torque and electromyographic amplitude when subjects perform isometric maximal voluntary contractions.⁴¹ Static stretching causes decreased strength production by affecting the length-tension relationship of the muscle secondary to changes in compliance.¹ If DN could have the same lengthening effects as stretching, it may also have a similar potentially detrimental effect on torque production.

However, Ge²¹ found that latent trigger points decrease force production, so it is vital to research the effect DN would have on force production. Dar and Hicks found²⁰ that DN in multifidus increases muscle function immediately. Cerezo-Telléz et al demonstrated an immediate cervical muscle strength in all directions of flexion, extension, rotation, and side bending after cervical DN.⁷ DN has improved muscle endurance of the knee extensors immediately and at four weeks in soccer players.²²

DN might not result in improved outcomes immediately. Huguenin et al¹⁰ found that pain improved with DN, but passive straight leg raise and hip internal rotation remained unchanged in both therapeutic dry needling and placebo

groups. However, Huguenin et al¹⁰ also concluded that they did not choose the appropriate ROM assessment. One RCT of DN demonstrated no improvement in hamstring ROM or knee pain immediately when compared to sham DN.⁴² One meta-analysis states that DN is less effective in treating pain conditions, but may be better at increasing ROM.¹⁹ DN has been demonstrated as less effective than acupuncture for motion-related chronic neck pain.⁴³

The purpose of this study was to assess the immediate effects of DN on myofascial trigger points in terms of skin surface temperature, pain, active and passive range of motion, and plantar flexion torque production in the triceps surae of ballet dancers. Based on the literature review, it was hypothesized that DN would have a significantly greater effect than sham DN on decreasing pain, increasing skin temperature, increasing ankle ROM, in addition to a decreasing in plantar flexion torque immediately following treatment.

METHODS

SUBJECTS

Eleven healthy, full time (25 hours of dance per week) ballet dancers were recruited from a professional ballet company by word of mouth. These dancers were included in the study if they presented with MTRPs in the calf through palpation. Dancers were excluded if they had significant health problems or bleeding disorders, were pregnant, feared needles, or had taken anticoagulant or pain relieving medications within the past 24 hours. Participation in this study was also denied if an individual had undergone DN or acupuncture in the prior four weeks. Prior to testing, the participants read and signed a specific Northwestern University and Andrews University IRB approved consent form. This included a health history form asking about recent injuries and general medical health.

Subjects were randomly assigned into two groups through computerized random number generation by the physical therapist performing the intervention so the assessors were blinded. The experimental group received bilateral DN to the calf, plus passive calf stretching of two repetitions of thirty seconds on a slant board in both knees bent and straight positioning, while the control group received bilateral sham DN (SHAM) in addition to the same slant board calf stretching protocol. All subjects had the intervention performed in a lab room at Northwestern University's Physical Therapy department. A physical therapist provided the intervention, while physical therapy students, blinded to group assignment, obtained pre/post measurements after they had practiced and demonstrated competency in their measurement areas.

MATERIALS

DN was performed using Seirin J-Type 30x30 needles. The 10cm VAS (Visual Analog Scale) was used to assess overall calf pain pre- and post- intervention as reported by participant, and temperature was measured with The Exergen TAT-2000 Series Professional Model Temporal Scanner (Exergen Corporation, Watertown, MA). A goniometer was used to measure the ankle dorsiflexion PROM in CKC



Figure 1: Trigger Points Palpated (Circled)

(closed kinetic chain) and AROM in OKC (open kinetic chain). An isokinetic dynamometer, the Biodex System 3 Pro (Biodex Medical Systems, Shirley, New York), was used to measure calf force.

PROCEDURE

Individuals meeting inclusion criteria were randomly blinded and placed into either the experimental or control group. The subjects were barefoot and wore athletic clothing. Each dancer followed a five-minute, standardized dance-specific warm up video prior to testing.

Initially, baseline skin temperature was measured at three standard points (SP) with a surface thermometer placed on marked spots 10cm below the popliteal fold, 10cm above the calcaneal tuberosity, and at a point in between. A blinded assessor performed and averaged the three measurements to discern an overall calf temperature pre-intervention. Then, a physical therapist certified in DN and having >10-years' experience palpating dancers' calf musculature located five trigger point (TP) spots using palpation. The trigger points were circled with a marker ([Figure 1](#))

After trigger point palpation, another blinded assessor administered a 10cm Visual Analog Scale (VAS) for the overall pain score of the five TP pre-intervention. The same blinded assessor measured all of the subjective calf pain for both the DN and SHAM groups pre- and post-intervention. The blinded assessor used the same script for each subject. The VAS has established test-retest reliability and sensitivity, and change in pain intensity can be easily obtained.⁴⁴ A 50% pain reduction measured by the VAS was determined as statistically significant.⁴⁴ As demonstrated in previous research, participants had access to their pre-test VAS at the

time of the post-test VAS to reduce error.⁴⁴

Next, the original temperature assessor measured surface temperature one more time to the nearest degree (Fahrenheit) at the marked TP and SP locations. The five TP measurements were averaged and the three SP measurements were averaged. This set of surface temperature measurements was performed post-palpation, but pre-intervention, to confirm that the physical therapist's palpation did not affect surface temperature. This second round of SP surface temperature measurements was conducted to confirm that the calf palpation did not change baseline temperature.

Ankle dorsiflexion AROM was measured bilaterally by a single blinded assessor using a standard 6-inch goniometer with the subjects in an open-kinetic chain seated position and for PROM in the closed-kinetic chain (CKC) weight-bearing lunge position. In OKC, the subjects sat over the edge of a plinth with their knees bent approximately 45-60° in order to prevent passive insufficiency of the gastrocnemius.⁴⁵ The subtalar joint was placed in neutral position with the transcondylar axis of the knee in the frontal plane.⁴⁶ Ankle dorsiflexion active range of motion (AROM) was measured in open chain first, using the standard text version.⁴⁷ The neutral position was 0° and ankle motion was recorded to the nearest degree from that position in the dorsal direction.⁴⁵ Subjects were asked to perform maximal active ankle dorsiflexion bilaterally three times and the angle was measured to the nearest degree. All measurements were recorded and the mean of the three was used for analysis.⁴⁶ Only AROM was measured in this position because it is more functional and applicable to ballet while passive ROM could be influenced by human error due to the inconsistency with overpressure.

Dickson et al⁴⁸ found that functional ankle dorsiflexion in modern dancers is best quantified in the weight-bearing lunge position, which has also been previously assessed for intra-rater (ICC=0.97-0.98) and inter-rater (ICC= 0.97-0.99) reliability.² Subjects were asked to stand facing a wall in a lunge position with one knee touching the wall and the foot on the same lower extremity sliding backward to the point of maximum tolerance while the knee remains in contact with the wall and the heel with the floor. Researchers verified that each subject's knee and heel remained in contact with the appropriate surface, and that the knee was aligned over the second toe prior to taking any measurements.⁴⁸ The same bony landmarks were used to align the goniometer as in OKC ankle dorsiflexion AROM. Measurements were recorded three times bilaterally and the mean of the three was used for analysis. (Figure 2)

Maximum muscular torque of the triceps surae was measured bilaterally using a Biodex prior to and after the DN or SHAM intervention. Two assessors blinded to group assignment performed all of these measurements. The participants were barefoot, seated and secured using a double chest strap, quadriceps strap, and two additional support straps fastened over the dorsum of the foot. The hip was placed at 90° flexion, the knee at 30° flexion, and the ankle at an angle of comfort for the participant. The axis of rotation of the Biodex ankle attachment was aligned with the lateral malleolus of the ankle being tested. Prior to testing, the participants warmed up by submaximally plantar- and dorsi-flexing each ankle ten times on the Biodex with no re-



Figure 2: Measurement of Closed-chain Ankle Dorsiflexion (PROM)




Figure 3: Biodex Measurement Set-up

sistance. The subject then performed four maximal concentric contractions plantar flexion at angular velocities of 60°, 90°, and 120°/sec with a two-minute rest period between tests to minimize fatigue. The order of velocities was randomized. The first measurement was eliminated, and the remaining three were averaged for analysis using standard Biodex software in Newton.meters (NM).^{52,46,49} (Figure 3).

After all the pre-tests, the DN was administered to participants in the experimental group, while the control group received SHAM dry needling while lying prone on a plinth. The assessors were not present during the DN or sham DN. The DN was performed in the five previously palpated and circled trigger point locations on each calf using a clean technique with sterile needles. Needles were inserted into

Table 1: Demographic data

SUBJECT NUMBER	GENDER	YEARS PROFESSIONAL DANCING	PRIOR DRY NEEDLING
1	M	12	YES
2	M	3	NO
3	M	6	YES
4	M	3	NO
5	F	6	YES
6	F	12	NO
7	F	15	YES
8	M	13	NO
9	F	15	NO
10	M	17	YES
11	F	4	YES

the triceps surae and repeatedly moved up and down in order to elicit a twitch response. All DN participants had twitch responses during the DN intervention.

The sham needles were prepared following the method adopted by Cotchett et al,⁵⁰ which was originally found to be valid by Tough and colleagues.⁵¹ Prior to treatment, the needle tips were removed with wire cutters.⁵² The end of the needle was filed down to create a blunt surface that when tapped would not pierce the skin.⁵⁰ The sham needle was placed back into its tube and repackaged. To begin each treatment, the skin was disinfected and a prepared sham needle was removed from packaging to stimulate a removal of a real needle.⁵² The sham needle was manipulated using the same technique as real dry needling. First, the applicator was placed on the skin and the dry needling administrator tapped the needle without piercing the skin. The applicator was removed while the administrator held the sham needle in place on the skin's surface and mimicked needle rotation for five seconds, then removed the needle. The sham needle was disposed of in a sharps container to simulate the noise and effects associated with sharps disposal.⁵⁰

Following the intervention, both groups were instructed to stretch on a 20° incline slant-board with straight and bent knees twice for 30 seconds. The choice to stretch after DN or SHAM as part of an intervention was to mimic the common tendency of ballet dancers to stretch after clinical treatment. Immediately following stretching, pain, temperature, ROM and torque were re-measured on both sides.

At the end of the procedures, the participants were debriefed on their treatment and given a contact number to call if they had any further questions or report adverse effects. No adverse effects were reported. Participants in the control group were given the opportunity to receive dry needling once they had completed the experiment but all declined. A summary of the methodology can be found in [Figure 4](#).

DATA ANALYSIS

The raw data collected was used to calculate the independent and dependent variables. SPSS Version 10.0 (SPSS Inc, Chicago, IL) was used to analyze this data. A paired t-test

was performed to compare pre- and post-intervention, as well as independent t-test to compare between the two test groups using a 95% confidence interval. Additionally, non-parametric statistics were run and the outcomes where statistically significant differences were found were the same between nonparametric and parametric analyses. Because of the specific homogenous population, parametric statistics were chosen. A histogram analysis looking for equally distributed variables was also performed. Statistical differences were assessed using the value $p < 0.05$.

RESULTS

Six male and five female professional ballet dancers from the same professional ballet company fit the inclusion/exclusion criteria. The average number of years dancing professionally was 9.6. Six out of eleven of the dancers had used DN prior to the study. This was a uniform sample, as the subjects were similar in age and from the same professional ballet company. Males and females were similarly represented in both groups. ([Table 1](#))

The mean measurements of pain, temperature, ROM and torque were calculated for all subjects prior to intervention. There were no significant differences for any mean measurement comparing right and left calves in temperature, pain, ROM, or force prior to the intervention. ([Table 2](#))

TEMPERATURE

Prior to intervention, the mean surface temperature of the TP measurements was higher than the SP measurements when performing an independent t-test. This statistical significance happened in both the right and left calves (right: $p = .014$; left: $p = .031$). ([Table 3](#)) Using a paired t-test, the mean temperature for both the right and left calves in the SHAM group (right: $p = .008$; left: $p = .008$) and only the right calf in the DN group (right: $p = .048$) showed significance in temperature decrease from pre- to post- intervention. ([Table 4](#))

Table 2: Comparison of means (\pm SD) between right and left sides of dancers prior to intervention

MEASUREMENT	RIGHT DATA	LEFT DATA	p-value (2-tailed)
Trigger point surface temperature	96.9 \pm .42	96.76 \pm .54	.187
VAS pain score	4.38 \pm 1.942	4.17 \pm 1.344	.759
OKC Sitting ROM	11.36 \pm 4.73	10.51 \pm 3.93	.420
CKC Lunge ROM	29.03 \pm 5.133	29.60 \pm 4.89	.620
Torque at 60 degrees/sec	36.79 \pm 10.53	35.47 \pm 11.66	.570
Torque at 90 degrees/sec	34.04 \pm 11.24	36.64 \pm 12.15	.378
Torque at 120 degrees/sec	34.22 \pm 8.47	34.22 \pm 11.85	1.00

Table 3: Mean temperature of three standard points (SP) calf measurements, reported in degrees, and statistical comparisons

MEASUREMENT	RIGHT SIDE DATA	LEFT SIDE DATA
SP pre/post palpation	Pre-palp: 95.51 \pm 1.74 Post-palp: 96.06 \pm 1.07 (t(10)= -1.77, p=.107, 2-tailed)	Pre-palp. 95.32 \pm 1.86 Post-palp. 95.72 \pm 1.57 (t(10)= -1.50, p=.164, 2-tailed)
SP post initial palpation compared to 3 standard points post DN	Post-palp: 95.82 \pm 1.37 Post-DN: 94.99 \pm 1.73 (t(5)=.898, p=.410, 2-tailed)	Post-palp: 95.82 \pm 1.32 Post-DN: 95.39 \pm 1.09 (t(5)=.582, p=.586, 2-tailed)
SP post initial palpation compared to SP post SHAM intervention	Post-palp: 96.35 \pm .57 Post-SHAM: 95.89 \pm .33 (t(4)=2.50, p=.067, 2-tailed)	Post-palp: 95.61 \pm 1.98 Post-SHAM: 95.51 \pm .68 (t(4)=.150, p=.888, 2-tailed)
Temperature difference of SP pre/post intervention, comparing DN and SHAM	Change DN: 1.18 \pm 1.42 Change SHAM: .47 \pm .39 (t(9)=2.02, p=.075, 2-tailed) Equal variances assumed Levene's significance= .000	Change DN: 1.29 \pm 1.17 Change SHAM: 1.07 \pm .83 (t(9)=.342, p=.740, 2-tailed) Equal variances assumed Levene's significance= .176
SP compared to trigger points (TP) post palpation and prior to intervention	SP: 96.06 \pm 1.07 TP: 96.88 \pm .43 (t(10)=-2.96, p=.014, 2-tailed)	SP: 95.72 \pm 1.57 TP: 96.76 \pm .55 (t(10)=-2.52, p=.031, 2-tailed)

Bold indicates significant difference

Table 4: Mean temperature of five trigger points (TP) calf measurements, reported in degrees with DN or SHAM and statistical comparisons

MEASUREMENT	RIGHT SIDE DATA	LEFT SIDE DATA
TP pre and post DN	Pre-DN: 96.86 \pm .23 Post-DN: 96.51 \pm .32 (t(5)= 2.61 , p=.048, 2-tailed)	Pre-DN: 96.72 \pm .45 Post-DN: 96.35 \pm .41 (t(5)=2.03, p=.098, 2-tailed)
TP pre and post SHAM	Pre-SHAM 96.92, \pm .62 Post-SHAM 96.17 \pm .33 (t(4)= 4.82 , p=.008, 2-tailed)	Pre-SHAM 96.81, \pm .70 Post-SHAM 95.97 \pm .47 (t(4)= 4.92 , p=.008, 2-tailed)
Temperature difference of TP comparing DN and SHAM	Change DN: .41, \pm .21 Change SHAM: .74, \pm .34 (t(9)= -1.99, p=.077, 2-tailed) Equal variances assumed Levene's significance= .474	Change DN: .48, \pm .30 Change SHAM: .83 \pm .38 (t(9)= -1.74, p=.116, 2-tailed) Equal variances assumed Levene's significance= .473

Bold indicates significant difference

Table 5: Mean visual analog pain scale score, in centimeters

MEASUREMENT	RIGHT DATA	LEFT DATA
Pre and Post DN	Pre-DN: 4.03±1.10 Post-DN: 3.38±1.68 (t(5)=1.32, p=.246, 2-tailed)	Pre-DN: 4.33±1.06 Post-DN: 3.69±1.95 (t(5)=.669, p=.533, 2-tailed)
Pre and Post SHAM	Pre-SHAM 4.80±2.74 Post-SHAM 4.58±2.74 (t(4)=.242, p=.821, 2-tailed)	Pre-SHAM 3.99±1.74 Post-SHAM 3.95±1.34 (t(4)=.040, p=.970, 2-tailed)
Change of VAS score, comparing DN and SHAM DN	Change DN .958±.906 Change SHAM: 1.44±1.27 (t(9)= -.735, p=.481, 2-tailed) Equal variances assumed Levene's significance= .435	Change DN 1.63±1.63 Change SHAM: 1.94±.591 (t(9)= -.396, p=.701, 2-tailed) Equal variances assumed Levene's significance= .083




Figure 4: Summary methods flow chart

VAS

There were no significant differences in mean VAS reported pain scores for either the SHAM or DN groups pre- to post-intervention. (Table 5)

ROM

In the seated AROM dorsiflexion measurement, the SHAM group had a statistically significant larger change in mean values in only the left calf (left: p= .005) when compared to the DN group. However, when a paired t-test compared

Table 6: Mean OKC dorsiflexion ROM (AROM), measured in a seated position, reported in degrees

MEASUREMENT	RIGHT DATA	LEFT DATA
Pre and Post DN	Pre-DN: 11.28±5.09 Post-DN 12.11±1.49 (t(5)= -.503, p=.636, 2-tailed)	Pre=DN: 11.05±3.48 Post-DN 10.28±3.64 (t(5)=1.94, p=.110, 2-tailed)
Pre and Post SHAM	Pre-SHAM 11.46±4.84 Post-SHAM 11.60±7.46 (t(4)= -.088, p=.934,2-tailed)	Pre-SHAM 9.87±4.77 Post-SHAM 9.73±5.90 (t(4)=.076, p=.943,2-tailed)
Change in Dorsiflexion, Comparing DN and SHAM	DN 2.61±3.03 SHAM 2.27±2.19 (t(9)=.211, p=.837,2-tailed) Equal variances assumed Levene's significance: .399	Change DN .78±.98 Change SHAM: 3.33±1.33 (t(9)= -3.66, p=.005, 2-tailed) Equal variances assumed Levene's significance: .425

Table 7: Mean CKC dorsiflexion ROM (PROM), measured during the lunge, reported in degrees

MEASUREMENT	RIGHT DATA	LEFT DATA
Pre and Post DN	Pre-DN: 28.22±5.26 Post-DN: 29.89±3.99 (t(5)= -1.58, p=.175,2-tailed)	Pre-DN: 29.05±4.98 Post-DN: 28.28±3.68 (t(5)= .629, p=.557,2-tailed)
Pre and Post SHAM	Pre-SHAM: 30.00±5.40 Post-SHAM: 32.80±4.58 (t(4)= -1.26, p=.277, 2-tailed)	Pre-SHAM: 30.27±5.26 Post-SHAM: 31.60±2.52 (t(4)= -.952, p=.395, 2-tailed)
Change in Dorsiflexion, Comparing DN and SHAM	Change DN: 1.67±2.59 Change SHAM: 4.67±2.71 (t(9)= -1.88, p=.093, 2-tailed) Equal variances assumed Levene's significance= .008	Change DN: 2.11±2.13 Change SHAM: 2.40±2.19 (t(9)= -.222, p=.829, 2-tailed) Equal variances assumed Levene's significance=.006

Table 8: Mean Biodex triceps surae torque in Newton.Meters at angular velocity 60 degrees/sec

MEASUREMENT	RIGHT DATA	LEFT DATA
PRE and POST DN	Pre-DN: 38.87±8.47 Post-DN: 45.09±10.80 (t(5)= -2.26, p=.073, 2-tailed)	Pre-DN: 35.61±6.26 Post-DN: 43.20±8.96 (t(5)= -3.09, p=.027,2-tailed)
PRE and POST SHAM	Pre-SHAM: 34.29±13.17 Post-SHAM: 40.83±14.37 (t(4)= -1.97, p=.121,2-tailed)	Pre-SHAM: 35.31±17.05 Post-SHAM: 45.03±19.55 (t(4)= -1.55, p=.197,2-tailed)
Change in Torque, Comparing DN to SHAM	Change DN: 7.23±5.40 Change SHAM: 8.19±4.97 (t(9)= -.304, p=.768,2-tailed) Equal variances assumed Levene's significance= .607	Change DN: 6.86±5.85 Change SHAM: 23.76±8.90 (t(9)= -2.55, p=.156,2-tailed) Equal variances assumed Levene's significance= .478

Bold indicates significant difference.

the pre- and post- intervention measurements of the seated AROM dorsiflexion measurement in the SHAM group on the left calf, there was no statistical significance. (Table 6)

There was no significant difference in the lunge CKC measurement of dorsiflexion between the DN to the SHAM group. (Table 7)

BIODEX TORQUE

There was a statistically significant (p= .027) increase in the torque of the plantar flexors at the 60-degree angular ve-

locity (Table 8) in the DN group left side only when pre- and post- intervention was compared with a paired t-test. In the 90 and 120-degree angular velocity (Table 9 and 10), there was a statistically significant difference at 90 degrees (p= .014) and 120 degrees (p= .001) noted for mean change in torque, when comparing DN and SHAM on the right side only. The SHAM group had a larger mean change than the DN group. This change in the SHAM group was an increase in torque, but it was not statistically significant when comparing pre/post intervention measurements for the 90 and 120-degree angular velocity using a paired t-test.

Table 9: Mean Biodex triceps surae torque in Newton.Meters at angular velocity 90 degrees/sec

MEASUREMENT	RIGHT DATA	LEFT DATA
PRE and POST DN	Pre-DN: 36.32±8.68 Post-DN: 37.38±7.87 (t(5)= -.347, p=.743,2-tailed)	Pre-DN: 38.85±8.77 Post-DN: 36.84±8.87 (t(5)=.839, p=.440,2-tailed)
PRE and POST SHAM	Pre-SHAM: 31.31±14.31 Post-SHAM: 36.63±15.44 (t(4)= -.554, p=.609,2-tailed)	Pre-Sham: 33.99±16.02 Post-Sham: 40.22±15.10 (t(4)= -1.717, p=.161, 2-tailed)
Change in Torque, Comparing DN vs. SHAM	Change DN: 5.31±4.89 Change SHAM: 18.23, ±9.00 (t(9)= -3.042, p=.014,2-tailed) Equal variances assumed Levene's significance= .110	Change DN: 5.22±2.56 Change SHAM: 7.91±6.05 (t(9)= -.994, p=.346,2-tailed) Equal variances assumed Levene's significance= .086

Bold indicates significant difference.

Table 10: Mean Biodex torque in Newton.Meters at angular velocity 120 degrees/sec

MEASUREMENT	RIGHT DATA	LEFT DATA
PRE and POST DN	Pre-DN: 34.61±4.77 Post-DN: 34.18±8.41 (t(5)= .158, p=.881,2-tailed)	Pre-DN: 35.67±9.55 Post-DN: 34.15±5.62 (t(5)=.538, p=.614,2-tailed)
PRE and POST SHAM	Pre-SHAM: 33.76±12.28 Post-SHAM: 35.28±12.30 (t(4)= -.246, p=.818,2-tailed)	Pre-SHAM: 32.48±15.16 Post-SHAM: 40.25±15.23 (t(4)= -2.067, p=.108,2-tailed)
Change in Torque, Comparing DN and SHAM	Change DN: 5.41±2.85 Change SHAM: 12.35, ±1.71 (t(9)= -4.762, p=.001,2-tailed) Equal variances assumed Levene's significance= .122	Change DN: 4.65±4.98 Change SHAM: 9.57±5.62 (t(9)= -1.540, p=.158,2-tailed) Equal variances assumed Levene's significance= .693

Bold indicates significant difference.

DISCUSSION

Baseline measurements of surface temperature, pain, ROM, and torque demonstrated no significant difference between the right and left sides prior to intervention. (Table 2) This is interesting because dancers often have a lateral preference when performing, but this was not seen in the current measurements.¹¹ This is relevant to future studies in dry needling because it demonstrates that perhaps only one side is needed for measurements and intervention.

Surface temperature measurements pre-intervention showed significantly lower SP mean temperature as compared to the TP mean temperature for the right and left sides (p=.014 right, p= .013 left). After intervention, both the right and left calves in the SHAM group, and right calf in the DN group showed a significant difference in mean temperature. The only other study that measured surface temperature with trigger points and dry needling was Skorpyska et al,³¹ and they noticed an increase in surface temperature in muscles distal to the muscle being treated along the same nerve distribution after DN. A larger sample size would continue to investigate temperature responses after dry needling, since current research remains inconclusive in the direction of temperature change.^{30,31} The surface temperature findings are important in that it invites new research because demonstrates a potential nature of trigger

point changes that may occur to MTrPs after DN. The baseline surface temperature is also important knowledge for the researchers that are trying to further understand trigger point physiology.

There were no significant findings for changes in pain which differs from the results of other investigations who demonstrated decreased pain after DN.⁶⁻¹⁷ Perhaps, specific population interprets pain differently, despite having used the uniform script to describe pain. Tajet-Foxell and Rose⁵³ found that dancers had a higher pain threshold in regards to duration of pain but not intensity on a cold pain tolerance test. Due to this, dancers might tolerate the palpation of the trigger points but might rank the post DN palpation of the tender points as higher on a VAS, delineating the results differently than a typical population. Additionally, the examiner asked the subjects to average the pain between all five trigger points. A suggestion for future research would be to have the subjects rank each spot individually for pain, and then calculate a mean measurement. Lastly, immediate post-needling muscle soreness may have contributed to VAS outcomes, as Ziaieifar et al⁵⁴ suggested that their subjects had improved pain intensity after two days instead of immediately after DN of the upper trapezius muscle.

The results of current research varies with trends in increases or decreases in ROM after immediate

DN.^{7,8,10,12,14,16,18} For seated AROM, the SHAM group had a statistically larger change in AROM pre/post intervention on the left side ($p = .005$) as compared to the DN group. The lunge PROM measurements had no statistical difference between the DN and SHAM groups. Since this measure was performed in self-controlled lunging, some of dancers in the DN group might have been sore from the intervention and therefore self-limited their passive motion. Further research with a larger sample size needs to be performed as this study was underpowered, because our power analysis for ROM measurements required a sample size of 48 subjects.

In the 60-degree angular velocity Biodex force, the DN group on the left side only showed a statistically significant increase in torque production. In both the 90 and 120-degree angular velocity tests, the SHAM group had a statistically significant change as compared to the DN group on the right side only, but this was not significant when comparing pre/post intervention measures. To date, there are no known studies that compare DN using measurements of plantar flexion torque immediately after intervention. Since these findings are limited by the size of the study, caution should be used when performing DN prior to dance performance. Further research into the immediate effects on plantar flexion torque should be conducted.

There are several limiting factors in this study. As previously mentioned, this study was underpowered, which may have resulted in Type II errors. In part, this was due to coordinating multiple assessors in addition to the research process requiring 90 minutes per subject. Additionally, some subjects had previous experience with DN and were familiar with the twitch response. This potentially allowed them the ability to guess which treatment group they were assigned to, therefore limiting the subject-related blinding. Two subjects requested to not be dry needled due to concerns of being sore for dance rehearsal, so the SHAM group had two non-blinded subjects, further impacting the validity of the blinded subjects' results.

This pilot study builds the framework for future DN studies, but further research is needed to determine the appropriate instruments to measure the effects of DN in the dance population. The unique methodology used in this study is important to consider when exploring further DN research. The assessors were blinded, reducing rater bias. All assessors were trained physical therapy students, and the same assessor took all measurements. Additionally, this pilot studied used a sham intervention to further blind the subjects.

CONCLUSION

The results of this research were inconclusive for changes in surface temperature, pain, ROM, and torque immediately after dry needling to the triceps surae musculature. However, this is the first blinded and randomized controlled study assessing the acute effects of DN in professional ballet dancers with a novel methodology for conducting further research. This study also presents potential information on the surface temperature of trigger points as compared to baseline temperature measurements. Further research is needed with larger sample sizes in order to fully understand the immediate effects of dry needling in the dance population prior to performance.

STATEMENT OF INSTITUTIONAL BOARD

Northwestern University, Chicago, IL STU00096692

CONFLICTS OF INTEREST

The authors report no conflicts of interest.

Submitted: April 30, 2020 CDT, Accepted: November 08, 2020 CDT



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-SA-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc-sa/4.0> and legal code at <https://creativecommons.org/licenses/by-nc-sa/4.0/legalcode> for more information.

REFERENCES

1. Behm DG, Chaouachi A. A review of the acute effects of static and dynamic stretching on performance. *Eur J Appl Physiol*. 2011;111(11):2633-2651. doi:10.1007/s00421-011-1879-2
2. Bennell K, Talbot R, Wajswelner H, Techovanich W, Kelly D. Intra-rater and inter-rater reliability of a weight-bearing lunge measure of ankle dorsiflexion. *Aust J Physiother*. 1998;44(3):175-180. doi:10.1016/s004-9514(14)60377-9
3. Allen N, Nevill A, Brooks J, Koutedakis Y, Wyon M. Ballet injuries: Injury incidence and severity over 1 year. *J Orthop Sports Phys Ther*. 2012;42(9):781-790. doi:10.2519/jospt.2012.3893
4. Ahonen J. Biomechanics of the foot in dance: A literature review. *J Dance Med Sci*. 2008;12(3):99-108.
5. Thomas KS, Parcell AC. Functional characteristics of the plantar flexors in ballet dancers, folk dancers, and non-dancer populations. *J Dance Med Sci*. 2004;8(3):73-77.
6. Abbaszadeh-Amirdehi M, Nakhostin Ansari N, Naghdi S, Olyaei G, Nourbakhsh MR. Therapeutic effects of dry needling in patients with upper trapezius myofascial trigger points. *Acupunct Med*. 2017;35(2):85-92. doi:10.1136/acupmed-2016-011082
7. Cerezo-Téllez E, Lacomba MT, Fuentes-Gallardo I, Mayoral del Moral O, Rodrigo-Medina B, Gutiérrez-Ortega C. Dry needling of the trapezius muscle in office workers with neck pain: A randomized clinical trial. *J Man Manip Ther*. 2016;24(4):223-232. doi:10.1179/2042618615y.0000000004
8. Fenandez-Carnero, et al. Short-term effects of dry needling of active myofascial trigger points in the masseter muscle in patients with temporomandibular disorders. *J Orofac Pain*. 2010;24:106-112.
9. Gattie E, Cleland JA, Snodgrass S. The effectiveness of trigger point dry needling for musculoskeletal conditions by physical therapists: A systematic review and meta-analysis. *J Orthop Sports Phys Ther*. 2017;47(3):133-149. doi:10.2519/jospt.2017.7096
10. Huguenin L, Brukner PD, McCrory P, Smith P, Wajswelner H, Bennett K. Effect of dry needling of gluteal muscles on straight leg raise: A randomised, placebo controlled, double blind trial. *Br J Sports Med*. 2005;39(2):84-90. doi:10.1136/bjism.2003.009431
11. Kimmerle M. Lateral bias, functional asymmetry, dance training and dance injuries. *J Dance Med Sci*. 2010;14(2):58-66.
12. Koppenhaver S et al. Effects of dry needling to the symptomatic versus control shoulder in patients with unilateral subacromial pain syndrome. *Manual Therapy*. 2016;26:62-69. doi:10.1016/j.math.2016.07.009
13. Llamas-Ramos R, Pecos-Martín D, Gallego-Izquierdo T, et al. Comparison of the short-term outcomes between trigger point dry needling and trigger point manual therapy for the management of chronic mechanics neck pain: a randomized clinical trial. *J Orthop Sports Phys Ther*. 2014;44(11):852-861. doi:10.2519/jospt.2014.5229
14. Mason J, Tansey K, Westrick R. Treatment of subacute posterior knee pain in an adolescent ballet dancer utilizing trigger point dry needling: A case report. *Int J Sports Phys Ther*. 2014;9(1):116-124.
15. Mayoral O, Salvat I, Martín MT, et al. Efficacy of myofascial trigger point dry needling in the prevention of pain after total knee arthroplasty: A randomized, double-blinded, placebo-controlled trial. *Ev Based Complement Altern Med*. 2013;2013:1-8. doi:10.1155/2013/694941
16. Osborne NJ, Gatt IT. Management of shoulder injuries using dry needling in elite volleyball players. *Acupunct Med*. 2010;28(1):42-45. doi:10.1136/aim.2009.001560
17. Tekin L, Akarsu S, Durmuş O, Çakar E, Dinçer Ü, Kıralp MZ. The effect of dry needling in the treatment of myofascial pain syndrome: A randomized double-blinded placebo-controlled trial. *Clin Rheumatol*. 2013;32(3):309-315. doi:10.1007/s10067-012-2112-3
18. Hsieh Y-L, Kao M-J, Kuan T-S, Chen S-M, Chen J-T, Hong C-Z. Dry needling to a key myofascial trigger point may reduce the irritability of satellite MTrPs. *Phys Med Rehabil*. 2007;86(5):397-403. doi:10.1097/phm.0b013e31804a554d
19. Rodriguez-Mansilla, et al. Effectiveness of dry needling on reducing pain intensity in patients with myofascial pain syndrome: A meta-analysis. *J Tradit Chin Med*. 2016;36(1):30001-30003. doi:10.1016/S0254-6272(16)30001-2
20. Dar G, Hicks GE. The immediate effect of dry needling on multifidus muscles' function in healthy individuals. *J Back Musculoskelet Rehabil*. 2016;29(2):273-278. doi:10.3233/BMR-150624

21. Ge H-Y, Arendt-Nielsen L, Madeleine P. Accelerated muscle fatigability of latent myofascial trigger points in humans. *Pain Med*. 2012;13(7):957-964. doi:10.1111/j.1526-4637.2012.01416.x
22. Haser C, Stöggel T, Kriner M, et al. Effect of dry needling on thigh muscle strength and hip flexion in elite soccer players. *Med Sci Sports Exerc*. 2017;49(2):378-383. doi:10.1249/mss.000000000000111
23. Cagnie B, Dewitte V, Barbe T, Timmermans F, Delrue N, Meeus M. Physiologic effects of dry needling. *Curr Pain Headache Rep*. 2013;17(8):348. doi:10.1007/s11916-013-0348-5
24. Bron C, Dommerholt JD. Etiology of myofascial trigger points. *Curr Pain Headache Rep*. 2012;16(5):439-444. doi:10.1007/s11916-012-0289-4
25. Ge H-Y, Fernández-de-las-Peñas C, Yue S-W. Myofascial trigger points: Spontaneous electrical activity and its consequences for pain induction and propagation. *Chinese Medicine*. 2011;6(13):2-7. doi:10.1186/1749-8546-6-13
26. Cagnie B, Barbe T, De Ridder E, Van Oosterwijck J, Cools A, Danneels L. The influence of dry needling of the trapezius muscle on muscle blood flow and oxygenation. *J Manip Phys Ther*. 2012;35(9):685-691. doi:10.1016/j.jmpt.2012.10.005
27. Kalichman L, Vulfsons S. Dry needling in the management of musculoskeletal pain. *J Am Board Fam Med*. 2010;23(5):640-646. doi:10.3122/jabfm.2010.05.090296
28. Jimbo S, Atsuta Y, Kobayashi T, Matsuno T. Effects of dry needling at tender points for neck pain: near-infrared spectroscopy for monitoring muscular oxygenation of the trapezius. *J Orthop Sci*. 2008;13(2):101-106. doi:10.1007/s00776-007-1209-z
29. Sandberg M, Larsson B, Lindberg L-G, Gerdle B. Different patterns of blood flow response in the trapezius muscle following needle stimulation (acupuncture) between healthy subjects and patients with fibromyalgia and work-related trapezius myalgia. *Eur J Pain*. 2005;9(5):497-510. doi:10.1016/j.ejpain.2004.11.002
30. Agarwal-Kozłowski K, Lange A-C, Beck H. Contact-free infrared thermography for assessing effects during acupuncture: A randomized, single-blinded, placebo-controlled crossover clinical trial. *Anesthesiology*. 2009;111(3):632-639. doi:10.1097/aln.0b013e3181b31e24
31. Skorupska E, Rychlik M, Pawelec W, Samborski W. Dry needling related short-term vasodilation in chronic sciatica under infrared thermovision. *J Evid Based Complement Altern Med*. 2015;1-10. doi:10.1155/2015/214374
32. Kietrys DM, Palombaro KM, Azzaretto E, et al. Effectiveness of dry needling for upper quarter myofascial pain: a systematic review and meta-analysis. *J Orthop Sports Phys Ther*. 2013;43(9):620-634. doi:10.2519/jospt.2013.4668
33. Macklin K, Healy A, Chockalingam N. The effect of calf muscle stretching exercises on ankle joint dorsiflexion and dynamic foot pressures, force and related temporal parameters. *The Foot*. 2012;22(1):10-17. doi:10.1016/j.foot.2011.09.001
34. Caruso JF, Coday MA, Ramsey CA, et al. Leg and calf press training modes and their impact on jump adaptations. *J Strength Cond Res*. 2008;22(3):766-772. doi:10.1519/jsc.0b013e31816a849a
35. Fernández-de-las-Peñas C. Interaction between trigger points and joint hypomobility: A clinical perspective. *J Man Manip Ther*. 2009;17(2):74-77. doi:10.1179/106698109790824721
36. Grieve R, Barnett S, Coghill N, Cramp F. Myofascial trigger point therapy for triceps surae dysfunction: A case series. *Man Ther*. 2013;18(6):519-525. doi:10.1016/j.math.2013.04.004
37. Grieve R, Clark J, Pearson E, Bullock S, Boyer C, Jarrett A. The immediate effect of soleus trigger point pressure release on restricted ankle joint dorsiflexion: A pilot randomised controlled trial. *J Bodywork Movement Ther*. 2011;15(1):42-49. doi:10.1016/j.jbmt.2010.02.005
38. Eftekharsadat B, Babaei-Ghazani A, Zeinolabedinzadeh V. Dry needling in patients with chronic heel pain due to plantar fasciitis: A single-blinded randomized clinical trial. *Med J Islam Repub Iran*. 2016;30:401.
39. Passigli S, Plebani G, Poser A. Acute effects of dry needling on posterior shoulder tightness: A case report. *Int J Sports Phys Ther*. 2016;11(2):254-263.
40. Mejuto-Vázquez MJ, Salom-Moreno J, Ortega-Santiago R, Truyols-Domínguez S, Fernández-de-las-Peñas C. Short-term changes in neck pain, widespread pressure pain sensitivity, and cervical range of motion after the application of trigger point dry needling in patients with acute mechanical neck pain: A randomized clinical trial. *J Orthop Sports Phys Ther*. 2014;44(4):252-261. doi:10.2519/jospt.2014.5108

41. Herda T, Ryan E, Costa P, et al. Acute effects of passive stretching and vibration on the electromechanical delay and musculotendinous stiffness of the plantar flexors. *Electromyogr Clin Neurophysiol*. 2010;50(6):277-288.
42. Mason, et al. The effectiveness of dry needling and stretching vs. stretching alone on hamstring flexibility in patients with knee pain: A randomized controlled trial. *Int J Sports Phys Ther*. 2016;11(5):672-683.
43. Irnich D, Behrens N, Gleditsch JM, et al. Immediate effects of dry needling and acupuncture at distant points in chronic neck pain: Results of a randomized, double-blind, sham-controlled crossover trial. *Pain*. 2002;99(1-2):83-89. doi:10.1016/s0304-3959(02)00062-3
44. Williamson A, Hoggart B. Pain: A review of three commonly used pain rating scales. *J Clin Nurs*. 2005;14(7):798-804. doi:10.1111/j.1365-2702.2005.01121.x
45. Russell JA, Kruse DW, Nevill AM, Koutedakis Y, Wyon MA. Measurement of the extreme ankle range of motion required by female ballet dancers. *Foot Ankle Spec*. 2010;3(8):324-330. doi:10.1177/1938640010374981
46. Kenne E, Unnithan VB. Knee and ankle strength and lower extremity power in adolescent female ballet dancers. *J Dance Med Sci*. 2008;12(2):59-65.
47. Norkin CC, White DJ. *Measurement of Joint Motion: A Guide to Goniometry*. 4th ed. Philadelphia, PA: F.A. Davis Company; 2009.
48. Dickson D, Hollman-Gage K, Ojofeitimi S, Bronner S. Comparison of functional ankle motion measures in modern dancers. *J Dance Med Sci*. 2012;16(3):116-125.
49. Feiring DC, Ellenbecker TS, Derscheid GL. Test-retest reliability of the Biodex isokinetic dynamometer. *J Orthop Sports Phys Ther*. 1990;11(7):298-300. doi:10.2519/jospt.1990.11.7.298
50. Cotchett MP, Landorf KB, Munteanu SE, Raspovic A. Effectiveness of trigger point dry needling for plantar heel pain: Study protocol for a randomised controlled trial. *J Foot Ankle Res*. 2011;4(1):5. doi:10.1186/1757-1146-4-5
51. Tough EA, White AR, Richards SH, Lord B, Campbell JL. Developing and validating a sham acupuncture needle. *Acupunct Med*. 2009;27(3):118-122. doi:10.1136/aim.2009.000737
52. Kreiner M, Zaffaroni A, Alvarez R, Clark G. Validation of a simplified sham acupuncture technique for its use in clinical research: A randomised, single blind, crossover study. *Acupunct Med*. 2010;28(1):33-36. doi:10.1136/aim.2009.001735
53. Tajet-Foxell B, Rose FD. Pain tolerance in professional ballet dancers. *Br J Sports Med*. 1995;29(1):31-34. doi:10.1136/bjism.29.1.31
54. Ziaiefar M, Arab AM, Nourbakhsh MR. Clinical effectiveness of dry needling immediately after application on myofascial trigger point in upper trapezius muscle. *J Chiropr Med*. 2016;15(4):252-258. doi:10.1016/j.jcm.2016.08.009