

Two cases of granuloma faciale showing rosettes

Benedetta Sonogo,^{1,2} Simona Sola,³ Giovanni Biondo,⁴ Cesare Massone¹

¹Dermatology Unit and Scientific Directorate, Galliera Hospital, Genoa; ²Dermatology Clinic of Trieste, Maggiore Hospital, University of Trieste; ³Surgical Pathology, Galliera Hospital, Genoa; ⁴Istituto Clinico Sant’Ambrogio, Gruppo Ospedaliero San Donato, Milan, Italy

Abstract

Granuloma faciale (GF) is a rare benign chronic inflammatory dermatosis often difficult to distinguish clinically from other diseases, both inflammatory and neoplastic. Dermoscopy can be a helpful diagnostic tool and indeed several dermoscopic criteria observed in GF have been described in literature. We present two patients affected by GF in which we have observed rosettes.

Correspondence: Benedetta Sonogo, Maggiore Hospital, piazza dell’Ospitale, 1 - 34125 Trieste, Italy.
Tel: +39.0403992056.
E-mail: benedetta.sonogo@gmail.com

Key words: granuloma faciale, dermoscopy, rosettes, inflammation.

Contributions: BS, CM, SS, GB, conceptualization; BS, CM, data curation, resources, writing; CM, supervision, validation. All the authors approved the final version to be published.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Availability data and materials: data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

Consent for publication: the patients in this manuscript have given written informed consent to the publication of their case details.

Received: 22 February 2023.

Accepted: 25 February 2023.

Early view: 14 June 2023.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Dermatology Reports 2023; 15:9696

doi:10.4081/dr.2023.9696

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Introduction

Granuloma faciale (GF) is a rare benign chronic inflammatory dermatosis often difficult to distinguish clinically from other diseases, both inflammatory and neoplastic. Several dermoscopic criteria observed in GF have been described in literature. We present two patients affected by GF in which we have observed rosettes.

Case Report #1

A 50-year-old man presented to our clinic for multiple erythematous plaques on the forehead that developed gradually over a year (Figure 1a). He complained of constant itching at that site.

Otherwise, healthy, he didn't take any medication. The only significant anamnestic note was the diagnosis 20 years ago of pulmonary sarcoidosis, treated and cured with betamethasone over 1 year, since then he has followed bi-annual check-ups.

At clinical examination, multiple erythematous round-oval plaques were present on the forehead. Dermoscopy with polarized light showed an erythematous-pink background, dilated vessels, white streaks, and multiple rosettes of about 0,1-0,2 mm diameter all over the lesions (Figure 1b). The rest of his skin was unaffected. A general physical examination did not reveal any gross alteration.

The main clinical and dermoscopic differential diagnoses were lupus erythematosus tumidus, sarcoidosis, and B-cell lymphoma.

An incisional biopsy of one of the frontal plaques was performed; histology showed an irregular epidermal hyperplasia with hypergranulosis, and under a grenz zone in the reticular dermis a diffuse lymphohistiocytic infiltrate with eosinophils and several telangiectatic vessels without vasculitis or fibrosis (Figure 1c-e). A diagnosis of GF was made.

Routine laboratory investigations, glucose-6-phosphate dehydrogenase deficiency, angiotensin-converting enzyme, and interleukin-6 receptor resulted within normal limits; autoantibody panels including antinuclear antibody and extractable nuclear antigen were negative. He performed a lung computed tomography scan that showed some traces of pulmonary thickening, a result of the previous sarcoidosis. A topic therapy with tacrolimus 0,1% ointment twice daily was prescribed.

Case Report #2

A 56-year-old man with a previous history of psoriasis arthritis treated with topic diclofenac, suffered for years of asymptomatic erythematous maculae of the scalp for which he tried various topical corticosteroid treatments and calcineurin inhibitors such as pimecrolimus without improvement.

The dermatological examination noted several polymorphic

erythematous patches and plaques on both parietal regions and vertex (Figure 2a). Observation with polarized dermoscopy showed a pink-red background, elongated telangiectasias, perifollicular whitish halo, and multiple rosettes of about 0,1-0,2 mm

diameter (Figure 2b). A biopsy showed a dense interstitial, perivascular, and periadnexal mixed lympho-epithelioid infiltrate with eosinophils, and dilatated vessels without vasculitis.




Figure 1. a) Multiple erythematous round-oval plaques were present on the forehead; b) dermoscopy with polarized light showed an erythematous-pink background, dilatated vessels, white streaks and multiple rosettes of about 0,1-0,2 mm diameter all over the lesions; c) irregular epidermal hyperplasia with hypergranulosis, and under a grenz zone in the reticular dermis a diffuse lymphohistiocytic infiltrate (HE, 40 \times); d) interstitial lymphohistiocytic infiltrate with eosinophils (HE, 100 \times); e) higher magnification of the interstitial lymphohistiocytic infiltrate with eosinophils (HE, 200 \times).




Figure 2. a) Several polymorphic erythematous patches and plaques on both parietal regions and vertex; b) dermoscopy showed a pink-red background, elongated telangiectasias, perifollicular whitish halo, and multiple rosettes of about 0,1-0,2 mm diameter.

Discussion

GF is a rare benign chronic inflammatory dermatosis that usually presents as solitary or multiple pink to purple papules or plaques typically located on the face. It is usually seen in middle-aged males.¹ The origin is unknown but different predisposing factors have been described such as actinic exposure, radiation, trauma, and allergy. The main clinical differential diagnoses are cutaneous sarcoidosis, discoid lupus erythematosus, erythema elevatum diutinum, lupus vulgaris, fixed drug eruption, lymphocytoma cutis, and cutaneous lymphomas.²

On dermoscopy, the background could be pink or red/erythematous.^{2,3} Vascular findings described in the literature are: focused and elongated telangiectasias, prominent telangiectasias and linear arborizing vessels in a parallel arrangement, linear branching vessels, dilated vessels, and ectatic vessels.^{1,3,4}

Peculiar criteria of GF are related to follicular abnormalities described dermoscopically by dilated follicular openings, perifollicular whitish halo, and follicular keratotic plugs.^{2,7}

Non-specific criteria are shiny white streaks described sometimes as orthogonal, whitish grayish structureless area, blackened area, aggregation of brown dots and globules, and purpuric spots.^{1-3,5,7} A new finding described by Jardim *et al.* is a yellowish area, histologically described as deposition of hemosiderin in the papillary dermis.⁵

Our cases showed numerous rosettes. Rosettes, also known as four-clod dots, are dermoscopic criteria described as four white dots oriented at the same angle, arranged as a four-leaf clover.⁸ Rosettes have been attributed to an optical effect of cross-polarization by fibrous or horny material into the hair follicle.³⁻⁸ To our knowledge, rosettes have not been observed in GF yet. First described in 2009 by Cuellar *et al.* in association with actinic keratosis, squamous cell carcinoma, and lichen planus-like keratosis, rosettes were later identified in discoid lupus erythematosus, dermatofibroma, scars, molluscum contagiosum, basal cell carcinoma, pseudolymphoma, melanoma, benign nevi.⁸

The main dermoscopic differential diagnoses in our cases included lupus erythematosus tumidus (LET), sarcoidosis, and B-cell lymphoma. LET is characterized by polymorphous vessels and follicular keratotic plugs on a pink-red background.⁹ Sarcoidosis reveals orange structureless areas and white areas with overlying focused linear vessels.⁷ B-cell lymphomas usually display orange structureless areas with unfocused linear vessels with branches (primary cutaneous marginal zone lymphoma) or orange globules along linear and linear-curved unfocused vessels (primary cutaneous follicle-center lymphoma); follicular plugs might be present.¹⁰

Conclusions

GF may be a simulator of many inflammatory and neoplastic diseases. When the clinic overlaps, dermoscopy could be helpful tool. Although rosettes are not specific to facial granuloma, they may be an additional criterion in formulating the diagnostic hypothesis.

References

1. Caldarola G, Zalaudek I, Argenziano G, et al. Granuloma faciale: A case report on long-term treatment with topical tacrolimus and dermoscopic aspects. *Dermatol Ther* 2011;24:508-11.
2. Teixeira DA, Estrozi B, Ianhez M. Granuloma faciale: a rare disease from a dermoscopy perspective. *An Bras Dermatol* 2013;88:97-100.
3. Lallas A, Sidiropoulos T, Lefaki I, et al. Photoletter to the editor: Dermoscopy of granuloma faciale. *J Dermatol Case Rep* 2012;6:59-60.
4. Pampena R, Paolino G, Didona D, et al. Multiple granuloma faciale: a clinical finding from a dermoscopic point a view. *G Ital Dermatol Venereol* 2018;153:439-40.
5. Jardim MML, Uchiyama J, Kakizaki P, Valente NYS. Dermoscopy of granuloma faciale: A description of a new finding. *An Bras Dermatol* 2018;93:587-9.
6. Lallas A, Argenziano G, Apalla Z, et al. Dermoscopic patterns of common facial inflammatory skin diseases. *J Eur Acad Dermatol Venereol* 2014;28:609-14.
7. Errichetti E. Dermoscopy of Inflammatory Dermatoses (Inflammoscopy): An Up-to-Date Overview. *Dermatol Pract Concept* 2019;9:169-80.
8. Haspelslagh M, Noë M, De Wispelaere I, et al. Rosettes and other white shiny structures in polarized dermoscopy: Histological correlate and optical explanation. *J Eur Acad Dermatol Venereol* 2016;30:311-3.
9. Żychowska M, Reich A. Dermoscopic Features of Acute, Subacute, Chronic and Intermittent Subtypes of Cutaneous Lupus Erythematosus in Caucasians. *J Clin Med* 2022;11:4088.
10. Errichetti E, Geller S, Zalaudek I, et al. Dermoscopy of nodular/plaque-type primary cutaneous T- and B-cell lymphomas: A retrospective comparative study with pseudolymphomas and tumoral/inflammatory mimickers by the International Dermoscopy Society. *J Am Acad Dermatol* 2022;86:774-81.