

M₂-TAM subsets altered by lactic acid promote T-cell apoptosis through the PD-L1/PD-1 pathway

TAO SHAN^{1*}, SHUO CHEN^{1*}, XI CHEN¹, TAO WU¹, YI YANG¹, SHUNLE LI¹, JIANCANG MA¹, JING ZHAO², WANRUN LIN³, WEI LI⁴, XIJUAN CUI⁵ and YA'AN KANG⁶

¹Department of General Surgery, The Second Affiliated Hospital of Medical College; ²Department of Chemistry, School of Science, Xi'an Jiaotong University, Xi'an, Shaanxi 710004, P.R. China; ³Department of Pathology, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA; ⁴Department for Population and Development Studies, School of Public Policy and Administration, Xi'an Jiaotong University; ⁵Department of General Surgery, First Affiliated Hospital of Medical College, Xi'an Jiaotong University, Xi'an, Shaanxi 710061, P.R. China; ⁶Department of Surgical Oncology, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

Received January 10, 2020; Accepted July 3, 2020

DOI: 10.3892/or.2020.7767

Abstract. The aim of the study was to investigate the effects of lactic acid on the phenotypic polarization and immune function of macrophages. The human monocyte/macrophage cell line, THP-1, was selected and treated with lactic acid. Immunofluorescence staining, laser confocal microscopy, reverse-transcription polymerase chain reaction (RT-PCR), western blot, siRNA, and ELISA analyses were used to observe changes in the levels of cluster of differentiation (CD)68, CD163, hypoxia inducible factor (HIF)-1 α , and programmed death ligand-1 (PD-L1) as well as those of cytokines, tumor necrosis factor (TNF)- α , interferon (IFN)- γ , interleukin (IL)-12, and IL-10. THP-1 macrophages and T cells were co-cultured *in vitro* to observe the changes in proliferation and apoptosis of T cells. The results showed that, lactic acid (15 mmol/l) significantly upregulated the expression of the macrophage M₂ marker CD163 (P<0.05), cytokines, IFN- γ and IL-10, secreted by M₂-tumor-associated macrophages (TAM, P<0.05), and HIF-1 α and PD-L1 (P<0.05), and down-regulated the expression of cytokines, TNF- α and IL-12, secreted by M₁-TAM (P<0.05). Redistribution of M₂-TAM subsets and PD-L1 expression was reversed after further transfection of THP-1 cells with HIF-1 α siRNA (P<0.05). After co-culturing, T-cell proliferation was inhibited and

apoptosis was promoted. In summary, modulation of lactic acid level can redistribute M₂-TAM subsets and upregulate PD-L1 to assist tumor immune escape. The HIF-1 α signaling pathway may participate in this process, revealing that macrophages, as 'checkpoints' in organisms, are links that connect the immune status and tumor evolution, and can be used as a target in tumor treatment.

Introduction

The development and progression of a tumor depends not only on its proliferation, but also on its interactions with the local microenvironment (1). Tumors have hypoxic stress microenvironments (2). An acidic microenvironment [pH (6.5-6.8) vs. (7.4)] caused by a large amount of anaerobic glycolysis is an important characteristic of malignant tumors, and an important factor in inducing their occurrence, metastasis, and drug resistance (3).

Tumor-associated macrophages (TAMs) are the most important cell subsets in the aforementioned environment, and their numbers and functions are affected by many factors related to the host and tumor (4). TAMs are primarily divided into the following two subtypes; M1-TAM is a part of the classical activation pathway, and M2-TAM is a part of the alternative activation pathway. M1-TAM shows a high expression of interleukin (IL)-12, low expression of IL-10, and plays an important role in antitumor activity (4). M2-TAM has low expression of IL-12, high expression of IL-10, and exerts a tumor-promoting effect. Experiments have shown that TAMs can play an important positive regulatory role in the development and progression of tumors (5). In recent years, with a more thorough understanding of acidic microenvironments, these are now considered a new target for tumor diagnosis and treatment, especially in regulating TAM immune escape (6). Previous studies have shown that lactic acid can maintain the immunosuppressive tumor microenvironment by regulating the transformation of macrophages to the M₂ type and reducing the cytotoxic function of natural killer

Correspondence to: Dr Xi Chen, Department of General Surgery, The Second Affiliated Hospital of Medical College, Xi'an Jiaotong University, No. 157 Xiwu Road 157 W 5th Road, Xi'an, Shaanxi 710004, P.R. China
E-mail: chenxijiaoda@163.com

*Contributed equally

Key words: microenvironment, pancreatic cancer, lactic acid, TAM, HIF-1 α , PD-L1

cells (7,8). Previous findings showed that TAMs can also combine with programmed death-1 (PD-1) expressed on the surface of T cells through programmed death-ligand 1 (PD-L1) and mediate the immune escape of tumor cells through the PD-L1/PD-1 pathway to promote tumor progression (9,10). However, the authenticity of this mechanism is unclear at present.

In this study, by investigating the effects of lactic acid on macrophage redistribution and immune function, the possible mechanism by which tumor cells, including pancreatic cancer cells, release lactic acid through glycolysis to promote immune escape was explored, providing a foundation for new treatment directions.

Materials and methods

Cell cultures and treatments. The human monocyte/macrophage cell line THP-1, human T lymphocyte cell line H9, and BxPC-3 human pancreatic cancer cell lines (American Tissue Type Collection) were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with penicillin (100 U/ml), streptomycin (100 µg/ml), 0.1 mM nonessential amino acids, 0.2 mM glutamine, 1 mM pyruvate, and 10% heat-inactivated fetal bovine serum (FBS) and were incubated in 5% CO₂ humidified atmosphere at 37°C. Cells were grown to 80% confluency prior to treatment.

Primary human pancreatic cells were isolated from the same pancreatic site in pancreatic cancer and non-cancerous partial pancreatectomy specimens, respectively, [tissue samples obtained from areas within 2.0 cm around the tumor were obtained from 45 patients undergoing partial pancreatectoduodenectomy (Whipple resection) for pancreatic cancer at the Department of Hepatobiliary and Pancreatic Surgery, Affiliated Hospital of the Xi'an Jiaotong University] from January 2016 to December 2018. Of the 45 patients, there were 30 men and 15 women. The median age at the time of surgery was 60.5 years (range 42-77 years). All the patients in the study signed informed consent. Briefly, each specimen was collected and transferred to the laboratory. After several washings with sterile phosphate-buffered saline (PBS), 1 cm² sections of tissues were placed into the wells of culture flasks. Once the tissue appeared to adhere to the flasks (5-6 h), DMEM containing 10% FBS was added gently to the tissue sections. Specimens were inspected daily and the medium was exchanged after 24 h for the first time and every third day thereafter. Tissue samples were then removed from the cultures and were transferred to larger tissue culture vessels once they had reached 70% confluency, after approximately 2 weeks. All the cells used for this study were between passages 3 and 5. Ethics approval was obtained from the Human Subjects Committee of the Xi'an Jiaotong University, China. The antibodies against cluster of differentiation CD68 (sc-20060), CD163 (sc-20066), HIF-1α (sc-13515), PD-L1 (sc-293425) and β-actin (sc-58673) were purchased from Santa Cruz Biotechnology.

Proliferation assay. Cell proliferation was determined via the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT; Sigma) uptake method. The cells were seeded (5x10³/well) in 200 µl of DMEM in 96-well plates

and cultured overnight. After treatment with lactic acid [0-20 mmol/l, the concentration was chosen based on a previous article (7)] at 37°C with 5% CO₂, MTT reagent (5 mg/ml) was added for the evaluation of cell growth, and incubation was continued for an additional 4 h. The reaction was terminated with 150 µl dimethyl sulfoxide (DMSO) per well. Absorbance values were determined using an MRX Revelation 96-well multiscanner (Dynex Technologies). The cells cultured in DMEM served as the control. The cell viability index was calculated according to the formula: experimental OD value/control OD value. The experiments were repeated three times.

In situ detection of apoptotic cells. TUNEL assays were performed using the *in situ* cell apoptosis detection kit following the manufacturer's instructions. Briefly, the cells were placed on cover slips and fixed with 4% paraformaldehyde for 30 min at 25°C. The non-specific chromogen reaction, induced by endogenous peroxidase, was inhibited with 3% hydrogen peroxide (H₂O₂) for 10 min. Terminal deoxynucleotidyl transferases (TdT) were used for the incorporation of DNA strand breaks *in situ* for 1 h at 37°C in a humidified box. Positive control slides were treated with DNase and negative control slides were treated with phosphate-buffered saline (PBS) instead of TdT. DNA fragments were stained using 3,3'-diaminobenzidine (DAB) as a substrate for peroxidase, and hematoxylin was used as a counterstain. The apoptotic index was calculated as a ratio of the number of apoptotic cells to the total number of tumor cells on each slide (10 of fields analyzed per sample).

Immunofluorescence assay. Exponentially growing cells were seeded on 25 mm square glass coverslips placed in 35 mm diameter culture dishes. After treatment, the cells were fixed with 4% formaldehyde for 5 min at 25°C, permeabilized with 0.2% solution of Triton X-100 in PBS, and blocked with 2% bovine serum albumin-PBS for 30 min at 25°C. Slides were incubated with anti-CD68 and anti-CD163 overnight. Fluorescent imaging was conducted with a confocal laser scanning microscope (Carl Zeiss MicroImaging, Inc.).

Reverse-transcription polymerase chain reaction. Total RNA was isolated using TRIzol[®] reagent (GIBCO BRL), and the quantities were determined spectrophotometrically. First-strand cDNA was synthesized from 2 µg of total RNA using a RevertAid Kit (Fermentas MBI). The polymerase chain reaction (PCR) primer sets were designed 1) for PD-L1, forward: 5'-CGACATGTGCTAGCATGCTGCTCCTGC-3' and reverse: 5'-CCCTCGAGGCGGCCGCTAGATCC TCTTTC-3'; 2) for CD68, forward: 5'-CACGACGACATG TGGACATTCT-3' and reverse: 5'-TGGGGCAGGAGAAAC TTTGCC-3'; 3) for CD163, forward: 5'-AGCATGGAAGCG GTCTCTGTGATT-3' and reverse: 5'-AGCTGACTCATTCCC ACGACAAGA-3'; and 4) for β-actin, forward: 5'-ATCGTG CGTGACATTAAGGAGAAG-3' and reverse: 5'-AGGAAG GAAGGCTGGAAGAGTG-3'. The PCR conditions included an initial cDNA synthesis reaction at 42°C for 1 h using a RevertAid Kit (Fermentas MBI) followed by a denaturation step for 5 min at 94°C and 22 cycles: 30 sec at 94°C, 30 sec at

55°C, and 30 sec at 72°C. After the last cycle, a final extension was performed at 72°C for 10 min. The housekeeping gene β -actin was used as an internal control.

Western blotting. Briefly, 5×10^5 cells were incubated on ice for 30 min in 0.5 ml of ice-cold whole-cell lysate buffer. Debris were removed by centrifugation at $23 \times g$, for 10 min at 4°C. The protein content of the cells was determined, and the cellular lysates were separated by 10% SDS-PAGE and electro-transferred onto nitrocellulose membranes. After being blocked with 5% non-fat milk in TBST, the membranes were incubated with primary antibodies CD68 (1:500, sc-20060), CD163 (1:500, sc-20066), HIF-1 α (1:500, sc-13515), PD-L1 (1:500, sc-293425) at 4°C overnight, followed by 1:2,000 horseradish peroxidase-conjugated secondary antibody (Santa Cruz, sc-516102) for 2 h. Immunoreactive bands were visualized using an Enhanced Chemiluminescence Kit (Amersham Pharmacia Biotech). The western blot signals were quantified by densitometric analysis using Total Lab Nonlinear Dynamic Image Analysis software (Nonlinear).

siRNA assay. To inhibit the expression of HIF-1 α , siRNA oligos were used (Qiagen). HIF-1 α siRNA target sequence was: 5'-AGGAAGAACTATGAACATAAA-3', and the sequence of the control siRNA was: 5'-UUCUCCGAACGU GUCACGUTT-3'; this selection was based on the results of our previous study (11). The cells ($n=2 \times 10^6$) were transfected with siRNA targeted against HIF-1 α (100 nm/l) or a control siRNA (Qiagen) using Lipofectamine 2000 (Invitrogen). Cells were covered overnight prior to starvation. This was then followed by treatment with lactic acid (15 mmol/l) for 24 h. Finally, the cells were harvested for reverse-transcription polymerase chain reaction (RT-PCR) and western blotting.

Indirect co-culture model. The T cells were added to Petri dishes at a density of 1.5×10^5 /ml; after 24 h, TAM-CM was added, and the cells were cultured for 48 h. Cells in PBS or serum-free medium were used as controls. ELISA and TUNEL assays were used to observe the factor and apoptosis of T cells in each Petri dish. Proteins were extracted from the cells. The co-culture of pancreatic cells, THP-1 or pancreatic cancer cells was performed in the same manner.

Statistical analysis. Each experiment was performed at least three times. Data are provided as means \pm standard deviation and differences were evaluated using Student's t-test and a one-way ANOVA (when the variance was homogeneous, Tukey's test was used as the post hoc test; when the variance was uneven, tamhane test was used as the post hoc test). Homogeneity of variance was determined with Fisher test. $P < 0.05$ was considered statistically significant. All statistical analyses were performed using SPSS Version 13.0 statistical software (SPSS).

Results

Changes in the proliferation of THP-1 macrophages with different concentrations of lactic acid. We first examined the effects of different concentrations of lactic acid (0, 5, 10, 15, and 20 mmol/l) on the proliferation of THP-1 cells




Figure 1. Changes in the proliferation of THP-1 macrophages with different concentrations of lactic acid. THP-1 cell proliferative effects were determined following treatment with different concentrations of lactic acid (0, 5, 10, 15, and 20 mmol/l) for 48 h. Results are representative of three independent experiments. No statistically significant difference between groups was found ($P > 0.05$).

by administering different concentrations for 48 h (7). Cell proliferation did not change significantly after treatment with lactic acid concentrations < 15 mmol/l ($P > 0.05$), whereas it was inhibited at 20 mmol/l, indicating drug toxicity. Therefore, lactic acid concentration of 15 mmol/l was selected for subsequent experiments (Fig. 1).

Lactic acid upregulated the expression of macrophage marker molecules in M_2 macrophages. It has previously been reported that CD68 is a specific molecular marker for macrophages, and CD163 expression is a marker for M_2 -TAMs (4). The ratio of CD163:CD68, thus, represents the proportion of M_2 -TAMs. To clarify whether lactic acid can redistribute M_2 -TAMs, THP-1 macrophages were treated with 15 mmol/l for 48 h. Immunofluorescence analysis revealed that the fluorescence intensity of CD163 in the lactic acid-treated group was significantly higher than that in the blank control group; by contrast, the fluorescence intensity of CD68 did not change considerably (Fig. 2A). RT-PCR and western blot analysis revealed that the ratio of CD163:CD68 in the lactic acid group was significantly higher than that in the blank control group (Fig. 2B and C). Similarly, the levels of cytokines, IFN- γ and IL-10, secreted in the supernatant by M_2 -TAMs of the lactic acid-treated group were higher than those secreted by M_2 -TAMs of the blank control group ($P < 0.05$). Additionally, the levels of cytokines, TNF- α and IL-12, secreted by M_1 -TAMs decreased ($P < 0.05$; Fig. 2D). These results suggested that lactic acid can redistribute M_2 -TAM subsets.

Lactic acid upregulated the expression of PD-L1 protein in M_2 -TAMs. To determine whether lactic acid can affect PD-L1 expression in macrophages, 15 mmol/l lactic acid was used to treat THP-1 macrophages for 48 h. RT-PCR showed that PD-L1 mRNA expression significantly increased in the lactic acid-treated group compared to that in the blank control group ($P < 0.05$; Fig. 3A and B). Similarly, western blot analysis revealed that the expression of PD-L1 protein was significantly upregulated in the lactic acid-treated




Figure 2. Lactic acid upregulated the expression of M₂ macrophage marker molecules in THP-1 induced by lactic acid. (A) CD68 and CD163 protein expression in THP-1 was induced by lactic acid stained with FITC-labeled IgG antibody and analyzed by confocal microscopy. The CD163 fluorescence signal in THP-1 cells induced by lactic acid was higher than that in the control; the fluorescence intensity of CD68 did not change considerably in the two groups (magnification, x400). (B) mRNA expression and quantification of CD68 and CD163 in THP-1 cells induced by lactic acid via RT-PCR. (C) The protein expression and quantification of CD68 and CD163 in THP-1 cells induced by lactic acid were assessed by western blotting. (D) The levels of cytokines IFN- γ , IL-10, TNF- α , and IL-12 secreted in THP-1 cells induced by lactic acid were assessed by ELISA. Data from at least three independent experiments with duplicate determinations were expressed as means \pm SD. *P<0.05 was considered statistically significant. NS, control group; Lactate, lactic acid group.




Figure 3. PD-L1 mRNA and protein expression in THP-1 cells induced by lactic acid was evaluated by RT-PCR and western blotting. (A and B) The mRNA expression and quantification of PD-L1 in THP-1 cells induced by lactic acid was evaluated by RT-PCR. (C and D) The protein expression and quantification of PD-L1 in THP-1 cells induced by lactic acid was evaluated by western blotting. Data from at least three independent experiments with duplicate determinations are expressed as means \pm SD. * $P < 0.05$, statistically significant. NS, control group; Lactate, lactic acid group.

group compared to that in the blank control group ($P < 0.05$; Fig. 3C and D), suggesting that lactic acid may affect tumor immune escape through the PD-L1/PD-1 pathway, after redistributing M₂-TAM subsets.

Effect of the lactic acid-induced redistribution of M₂-TAM subsets on the biological behaviors of T cells. PD-L1 and PD-1 are negative costimulatory molecules that can induce apoptosis of specific cytotoxic T lymphocytes (CTLs) and reduce the sensitivity of tumors to the cytotoxic effect of CTLs, thereby assisting tumor immune escape. To determine whether M₂-TAMs overexpressing PD-L1 following treatment with lactic acid affected the proliferation and apoptosis of T cells, T cells and macrophages were first co-cultured indirectly, then lactic acid/PBS was added to the co-culture (Fig. 4A). M₂-TAMs in the lactic acid treatment group had a significantly lower ability to induce activation and proliferation of T cells than those co-cultured in the PBS control group (Fig. 4B). T-cell apoptosis in the co-cultured group was higher in the lactic acid treatment group than that in the PBS control

group (Fig. 4C; Table I), suggesting that M₂-TAMs reduce the cytotoxic effect of T cells.

HIF-1 α signaling pathway may be the key regulatory mechanism. To determine whether the HIF-1 α signaling pathway was involved in the lactic acid-induced redistribution of M₂-TAM subsets and PD-L1 overexpression, a western blot was used to detect HIF-1 α protein expression in THP-1 macrophages treated with lactic acid for 48 h. HIF-1 α expression was significantly increased in the lactic acid-treated group as compared to that in the blank control group (Fig. 5A and D). This suggests that HIF-1 α plays a regulatory role in the redistribution of M₂-TAM subsets and PD-L1 overexpression. Successful transfection with HIF-1 α siRNA was verified using western blotting (Fig. 5B and E). To further verify the effect of HIF-1 α on M₂-TAM subsets and PD-L1 overexpression, we pre-treated macrophages with HIF-1 α siRNA and treated them with lactic acid (Fig. 5C and F). There was no significant difference in the distribution of M₂-TAM subsets and PD-L1 expression between the HIF-1 α siRNA-treated group and the




Figure 4. Effect of the lactic acid-induced redistribution of M₂-TAM subsets on the biological behaviors of T cells. (A) Schematic representation of the co-culturing technique. (B) The T-cell proliferative effect was lower in the lactic acid treatment co-culture group than in the control group. (C) TUNEL staining showed T-cell apoptosis was higher in the lactic acid treatment indirect co-culture group than that in the control group. *P<0.05 was considered statistically significant. Con, Control co-culture group; Lactate, Lactic acid co-culture group.

Table I. T-cell apoptosis of co-cultured group was higher in the lactic acid treatment group than that in the PBS control group (mean ± SD).

Group	TUNEL-positive cells (apoptotic index) (%)
Control	1.05±0.27
Lactic acid	9.46±0.14 ^a

^aP<0.05 vs. control group.

control group (Fig. 5G-J; Table II). These data indicated that the HIF-1 α signaling pathway participates to some extent in the lactic acid-induced redistribution of M₂-TAM subsets and PD-L1 overexpression.

Redistribution of M₂-TAM subsets and effects on T cell apoptosis after the co-culture of pancreatic cancer cells, macrophages, and T cells. To clarify whether lactic acid released by pancreatic cancer cells redistributes M₂-TAM subsets and regulates immune function, we first detected the lactic acid concentration in the supernatant after the

Table II. T-cell apoptosis of HIF-1 α siRNA-macrophages co-culture^a.

Group	TUNEL-positive cells (apoptotic index) (%)
Control	1.13±0.16
Lactic acid	1.22±0.20 ^b

^aNo significant difference between the lactic acid group and the control group (mean ± SD). ^bP<0.05 vs. control group.

co-culturing of pancreatic cancer cells, macrophages, and T cells using a direct or indirect triple-cell culture model (Fig. 6A). Compared with the normal pancreatic cell culture group, the lactic acid concentration in the pancreatic cancer cell co-culture group significantly increased (P<0.05; Fig. 6D). To determine whether lactic acid released by pancreatic cancer cells redistributed M₂-TAMs, the co-cultured cells were collected to detect CD163/CD68 mRNA expression. RT-PCR results showed that the ratio of CD163:CD68 mRNA was significantly upregulated in the pancreatic cancer cell co-culture group compared with




Figure 5. The HIF-1 α signaling pathway may be the key regulatory mechanism. (A and D) Western blot analysis and RT-PCR show that the HIF-1 α protein expression in THP-1 macrophages in the lactic acid-treated group significantly increased compared with that in the blank control group. (B and E) The HIF-1 α protein expression was successfully reduced using transfection with HIF-1 α siRNA. (C and F) The efficacy of HIF-1 α siRNA for knockdown of HIF-1 α protein was confirmed by western blotting. (G and I) There was no significant difference in PD-L1 expression between the HIF-1 α siRNA-treated group and the control group. (H and J) TUNEL staining showed T-cell apoptosis was not significantly different between the HIF-1 α siRNA-treated indirect co-culture group and the control group. Data from at least three independent experiments with duplicate determinations were expressed as means \pm SD. * P <0.05 was considered statistically significant.

that in the normal pancreatic cell culture group (P <0.05) (Fig. 6E and G). Similarly, RT-PCR results showed that PD-L1 mRNA expression was significantly upregulated in the pancreatic cancer cell co-culture group as compared to that in the normal pancreatic cell culture group (P <0.05) (Fig. 6B and C). Compared with the normal pancreatic cell culture group, the pancreatic cancer cell co-culture group significantly induced T cell apoptosis (P <0.05; Fig. 6H;

Table III). The results of the triple-cell culture suggested that lactic acid released by pancreatic cancer cells can redistribute M₂-TAM subsets and regulate immune escape.

Discussion

Tumor cells consist of an acidic microenvironment (2). Lactic acid, a glycolytic metabolite, is believed to be a key factor in




Figure 6. Redistribution of M₂-TAM subsets and effects on T-cell apoptosis after the triple-cell indirect culture model of pancreatic cancer cells, macrophages, and T cells. (A) Schematic representation of the technique of the triple-cell culture model. (B and C) RT-PCR analysis shows that the PD-L1 mRNA expression in THP-1 cells in the pancreatic cancer cell co-culture group was significantly upregulated compared to that in the normal pancreatic cell co-culture group. (D) ELISA results showed that lactate concentration in the pancreatic cancer cell co-culture group was significantly upregulated compared with that in the normal pancreatic cell co-culture group. (E-G) RT-PCR show that the CD163/CD68 mRNA expression of THP-1 in the pancreatic cancer cell co-culture group was significantly upregulated compared to that in the normal pancreatic cell co-culture group. (H) TUNEL assays show that T-cell apoptosis was higher in the pancreatic cancer cell co-culture group than that in the normal pancreatic cell co-culture group. Data from at least three independent experiments with duplicate determinations are expressed as means ± SD. *P<0.05 was considered statistically significant.

Table III. T-cell apoptosis was higher in the pancreatic cancer cell co-culture group compared with the normal pancreatic cell culture group (mean ± SD).

Group	TUNEL-positive cells (apoptotic index) (%)
Normal cell	1.07±0.62
Cancer cell	7.00±0.52 ^a

^aP<0.05 vs. normal cell group.

redistributing macrophage polarization in the tumor microenvironment (3). Previously, it was shown that a high-lactic acid environment in pancreatic cancer tissue samples is consistent with the distribution of M₂-TAM subsets (7). The results of this *in vitro* experiment confirmed that lactic acid could redistribute M₂-TAM subsets (7), which is consistent with the results of the current study. This suggests that lactic acid may

be an initiating factor for M₂-TAMs to promote tumor progression by their transformation from a ‘good’ macrophage into a ‘bad’ macrophage.

The development and progression of solid tumors are immune regulation processes (12). Various factors in the tumor microenvironment interact with each other, interfering with the balance of the immune system, resulting in the immune escape of tumors (13). According to previous findings, macrophages act as checkpoints by determining whether to initiate an immune response or induce immune tolerance, thus, serving as a bridge between the immune state and tumor evolution (4). The mechanism underlying the M₂-TAM modulation of immune escape has not yet been fully elucidated, and previous findings have shown that PD-L1 may be involved in this process (9). PD-L1 belongs to the B7 family and contains IgV-like, IgC-like, and transmembrane regions, and a cytoplasmic tail (14). The latter is involved in intracellular signal transduction, while the IgV and IgC regions are involved in intercellular signal transduction. The binding of PD-L1 to PD-1 on T cells promotes the phosphorylation of tyrosine in the

ITSM domain of PD-1, causes the dephosphorylation of downstream Syk kinase and phosphoinositide 3-kinase, inhibits the activation of downstream AKT, extracellular signal-activated kinase, and other pathways, and ultimately inhibits the transcription and translation of the genes and cytokines required for T-cell activation (14). These activities play a role in negatively regulating T-cell activity. For example, Dong *et al* (15), based on the results of an *in vitro* experiment utilizing mouse models, reported that activation of the PD-L1/PD-1 signaling pathway may induce the specific apoptosis of CTLs, which reduces the sensitivity of the CTL cytotoxic killing effect and promotes the immune escape of tumor cells. Zhou *et al* (16), found that in patients with liver cancer, TAM also inhibits functions of tumor-infiltrating lymphocytes through the highly expressed PD-L1 on its surface, leading to the escape of tumor cells from immune cell attack. Consistent with this finding, we observed that the high expression of PD-L1 in the TAM of pancreatic cancer was consistent with CD163. Moreover, correlation analysis of clinical factors revealed that the overexpression of PD-L1 in TAM was positively associated with lymph node metastasis and tumor staging. In the analysis of prognostic factors of pancreatic cancer, PD-L1 in TAM may serve as an independent risk factor. Patients with PD-L1 overexpression had a significantly worse prognosis than patients with low expression, suggesting that the PD-L1/PD-1 signaling pathway is involved in the mechanism underlying the M₂-TAM regulation of tumor immune escape (17). However, results of this study still need to be verified through relevant *in vitro* experiments.

The abovementioned results show that modulation of lactic acid level may initiate the polarization of macrophages. Previous findings regarding this mechanism have reported that lactic acid is absorbed and oxidized by macrophage mitochondria through MCT1 into pyruvate (18). The latter inhibits prolyl hydroxylase domain (PHD) activity through direct competition with the PHD substrate, leading to the stable accumulation of HIF-1 α , and subsequently, inducing the transformation of the M₂-TAM subsets (19). In the present study, we found that the expression of HIF-1 α protein significantly increased after the treatment of THP-1 macrophages with lactic acid. There was no significant difference in the distribution of M₂-TAM cell subsets and PD-L1 expression after macrophages were pretreated with HIF-1 α siRNA then treated with lactic acid. This indicates that the HIF-1 α signaling pathway participates to some extent in the redistributing of M₂-TAM subsets by lactic acid and PD-L1 overexpression, which is consistent with the finding that HIF-1 α is involved in the redistribution of M₂-TAM subsets and regulation of the immune function. Moreover, we used a triple-cell culture model consisting of pancreatic cancer cells, macrophages, and T cells to simulate the real microenvironment of pancreatic cancer, and again verified that lactic acid released by glycolysis in pancreatic cancer cells could redistribute M₂-TAMs and further regulate PD-L1/PD-1 signals to mediate T-cell apoptosis, thereby assisting immune escape. Results of the present study demonstrated the significance of the involvement of lactic acid, a key metabolite, in tumor progression. siRNA is an important tool for genetic analysis of mammalian cells. siRNA-induced gene silencing in mammalian cells can effectively inhibit the expression of specific genes. In this experiment, single siRNA

of HIF-1 α corresponding to target genes were prepared and detected. Further research with more siRNAs of HIF-1 α may be useful to obtain more reliable results.

There are some limitations to our study. First, lactic acid alone (without macrophage co-culture) may affect T-cell proliferation and apoptosis, and lactic acid alone was not applied as a control group to exclude this bias. Second, we performed TUNEL assays to assess T-cell apoptosis. However, the results could be further verified if other methods such as annexin-V staining could also be used. Third, we did not evaluate the mechanism for reshaping and immune function of M2-TAM subsets. This remains to be studied in further experiments.

In conclusion, this *in vitro* experiment further explored the hypothesis that tumor cells use lactic acid, a glycolysis product, as a weapon to activate M₂-TAMs via the bystander effect and, thus, further regulate PD-L1/PD-1 signal-mediated immune escape. These results suggest that lactic acid, macrophages, and PD-L1 are intrinsically linked, and provide an important theoretical basis for the treatment of pancreatic cancer.

Acknowledgements

Not applicable.

Funding

This study was supported by the Innovation Talent Promotion Project Foundation of Shaanxi Province (grant no. 2018KJXX-058); the Hospital Fund of the Second Affiliated Hospital of the Health Science Center, Xi'an Jiaotong University [grant no. RC(GG)201708]; and the National Natural Science Foundation of China, NSFC (grant no. 81402583); Natural Science Foundation of Shaanxi Province (grant no. 2019JQ-969); Xi'an Jiaotong University Education Foundation, XJTUEF (grant no. xjj2018141).

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Authors' contributions

TS, SC and XC conceived and designed the experiments. TS, SC, TW and WL performed the experiments. XC, SL, JZ and YY analyzed the data. JM, WL, XC, WL and YK contributed to the conducting the experiments. TS wrote the paper and XC, WL and YK revised the article. All authors approved the final version of the manuscript.

Ethics approval and consent to participate

Ethics approval was obtained from the Human Subjects Committee of the Xi'an Jiaotong University, China. All the patients in the study provided signed informed consent.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Moore A: Cancer ecology: The intracellular interactome makes little sense without the intercellular one. *Bioessays* 40: e1800202, 2018.
- Yan L, Raj P, Yao W and Ying H: Glucose metabolism in pancreatic cancer. *Cancers (Basel)* 11: 1460, 2019.
- Rawat D, Chhonker SK, Naik RA, Mehrotra A, Trigun SK and Koiri RK: Lactate as a signaling molecule: Journey from dead end product of glycolysis to tumor survival. *Front Biosci (Landmark Ed)* 24: 366-381, 2019.
- Najafi M, Goradel NH, Farhood B, Farhood B, Salehi E, Nashtaei MS, Khanlarkhani N, Khezri Z, Majidpoor J, Abouzaripour M, *et al*: Macrophage polarity in cancer: A review. *J Cell Biochem* 120: 2756-2765, 2019.
- Xu JY, Wang WS, Zhou J, Liu CY, Shi JL, Lu PH and Ding JL: The importance of a conjoint analysis of tumor-associated macrophages and immune checkpoints in pancreatic cancer. *Pancreas* 48: 904-912, 2019.
- Bohn T, Rapp S, Luther N, Klein M, Bruehl TJ, Kojima N, Aranda Lopez P, Hahlbrock J, Muth S, Endo S, *et al*: Tumor immunoevasion via acidosis-dependent induction of regulatory tumor-associated macrophages. *Nat Immunol* 19: 1319-1329, 2018.
- Colegio OR, Chu NQ, Szabo AL, Chu T, Rhebergen AM, Jairam V, Cyrus N, Brokowski CE, Eisenbarth SC, Phillips GM, *et al*: Functional polarization of tumour-associated macrophages by tumour-derived lactic acid. *Nature* 513: 559-563, 2014.
- Terren I, Orrantia A, Vitale J, Zenarruzabeitia O and Borrego F: NK cell metabolism and tumor microenvironment. *Front Immunol* 10: 2278, 2019.
- Cai J, Qi Q, Qian X, Han J, Zhu X, Zhang Q and Xia R: The role of PD-1/PD-L1 axis and macrophage in the progression and treatment of cancer. *J Cancer Res Clin Oncol* 145: 1377-1385, 2019.
- Sau S, Alsaab HO, Bhise K, Alzhrani R, Nabil G and Iyer AK: Multifunctional nanoparticles for cancer immunotherapy: A groundbreaking approach for reprogramming malfunctioned tumor environment. *J Control Release* 274: 24-34, 2018.
- Shan T, Cui X, Li W, Lin W, Li Y, Chen X and Wu T: Novel regulatory program for norepinephrine-induced epithelial-mesenchymal transition in gastric adenocarcinoma cell lines. *Cancer Sci* 105: 847-856, 2014.
- Bolli E, D'Huyvetter M, Murgaski A, Berus D, Stangé G, Clappaert EJ, Arnouk S, Pombo Antunes AR, Krasniqi A, Lahoutte T, *et al*: Stromal-targeting radioimmunotherapy mitigates the progression of therapy-resistant tumors. *J Control Release* 314:1-11, 2019.
- Kalinski P and Talmadge JE: Tumor immuno-environment in cancer progression and therapy. *Adv Exp Med Biol* 1036: 1-18, 2017.
- Sun C, Mezzadra R and Schumacher TN: Regulation and function of the PD-L1 checkpoint. *Immunity* 48: 434-452, 2018.
- Dong H, Strome SE, Salomao DR, Tamura H, Hirano F, Flies DB, Roche PC, Lu J, Zhu G, Tamada K, *et al*: Tumor-associated B7-H1 promotes T-cell apoptosis: A potential mechanism of immune evasion. *Nat Med* 8: 793-800, 2002.
- Zhou G, Sprengers D, Boor PPC, Doukas M, Schutz H, Mancham S, Pedroza-Gonzalez A, Polak WG, de Jonge J, Gaspersz M, *et al*: Antibodies against immune checkpoint molecules restore functions of tumor-infiltrating T cells in hepatocellular carcinomas. *Gastroenterology* 153: 1107-1119.e10, 2017.
- Wang L, Ma Q, Chen X, Guo K, Li J and Zhang M: Clinical significance of B7-H1 and B7-1 expressions in pancreatic carcinoma. *World J Surg* 34: 1059-1065, 2010.
- Silva LS, Poschet G, Nonnenmacher Y, Becker HM, Sapcariu S, Gaupel AC, Schlotter M, Wu Y, Kneisel N, Seiffert M, *et al*: Branched-chain ketoacids secreted by glioblastoma cells via MCT1 modulate macrophage phenotype. *EMBO Rep* 18: 2172-2185, 2017.
- Wei L, Zhou Y, Yao J, Qiao C, Ni T, Guo R, Guo Q and Lu N: Lactate promotes PGE2 synthesis and gluconeogenesis in monocytes to benefit the growth of inflammation-associated colorectal tumor. *Oncotarget* 6: 16198-16214, 2015.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.