

relatively low doses of antipsychotics was possible with a mean length of stay of $7,25 \pm 2,86$ days. In two of the cases the delirious content was predominantly marked by the coronavirus itself.

Conclusions: It has been suggested that the intense psychosocial stress associated with a new life-threatening disease and national lockdown restrictions could be triggers for new-onset psychotic disorders. Some authors have reported cases similar to ours, which means that we could be experiencing and increase in the incidence of psychotic disorders due to the exceptional circumstances we are living around the world.

Keywords: COVID-19; pandemic; Acute and transient psychotic disorder; Reactive psychosis

EPP0286

Relationship between fear of COVID-19 and individual factors – a preliminary study

C. Cabaços^{1,2*}, A.T. Pereira², A. Araujo^{1,2}, R. Sousa³ and A. Macedo^{1,2}

¹Psychiatry Department, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal; ²Institute Of Psychological Medicine, Faculty Of Medicine, University of Coimbra, Coimbra, Portugal and ³Uf Coimbra Centro, USF Coimbra Centro, Coimbra, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.694

Introduction: Fear associated to infectious diseases is directly related with their transmission rate, morbidity and mortality. High levels of fear associated with COVID-19 can affect people's ability to act and think rationally. In a time of pandemics, it is essential to understand individual factors that might be associated to higher vulnerability to stress and fear.

Objectives: To analyse: a) correlations between Fear of Covid-19 and clinical and sociodemographic characteristics; b) the mediator role of repetitive negative thinking on the relationship between personality traits and Fear of Covid-19.

Methods: 234 adults (75.6% women; mean age = 29.53 ± 12.51) completed an on-line survey with the Portuguese version of the Fear of Covid-19 Scale (FCV-19S) and other questionnaires to evaluate clinical and sociodemographic characteristics (years of education, perceived physical and mental health and infection by Covid-19), Personality (NEO-FFI-20) and Repetitive negative thinking (PTQ-15). SPSS and Process Macro (Hays, 2020) were used.

Results: FCV-19 mean scores were significantly higher in women and significantly correlated with years of education ($r = -.14$) ($p < .05$). History of previous/current Covid-19 infection did not significantly distinguish FCV-19 scores and they did not correlate with perceived health. FCV-19 correlated significantly with neuroticism and PTQ total and dimensional scores ($r > .20$, $p < .01$). Both Repetitive thinking and Cognitive interference were mediators of the relationship between neuroticism and fear of COVID, even after controlling for gender and education.

Conclusions: This study provides preliminary evidence on individual factors that might be associated to the emotional response to the Covid-19 pandemics, aiming to facilitate public health initiatives to ease people's fears in a near future.

Keywords: COVID-19; Fear of COVID-19; Personality; Repetitive negative thinking

EPP0287

Psychopathological consequences related to COVID-19 infection: The most relevant reactions of the general population

G. Serafini^{1*}, A. Aguglia¹, A. Amerio¹, L. Sher² and M. Amore³

¹Department Of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal And Child Health (dinogmi), University of Genoa, IRCCS Ospedale Policlinico San Martino, Genoa, Italy, Genoa, Italy;

²Psychiatry, Mount Sinai Hospital, New York City, United States of America and ³Department Of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal And Child Health (dinogmi), Dipartimento di Neuroscienze, Università di Genova, Genoa, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.695

Introduction: As a result of the emergence of coronavirus disease 2019 (COVID-19) outbreak caused by acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in the Chinese city of Wuhan, a situation of socio-economic crisis and profound psychological distress rapidly occurred worldwide.

Objectives: This work aimed to comprehensively review the current literature about the impact of COVID-19 infection on the mental health in the general population.

Methods: A detailed review has been conducted in order to identify the main psychopathological consequences related to Covid-19 infection in the general population.

Results: Various psychological problems and important consequences in terms of mental health including stress, anxiety, depression, frustration, uncertainty during COVID-19 outbreak emerged progressively. The psychological impact of quarantine related to COVID-19 infection has been additionally documented together with the most relevant psychological reactions in the general population related to COVID-19 outbreak.

Conclusions: The role of risk and protective factors against the potential to develop psychiatric disorders in vulnerable individuals with Covid-19 infection need to be carefully addressed in the clinical practice.

Keywords: COVID-19 infection; mental health; preventive strategies; Psychological Distress

EPP0288

Portuguese version of the fear of COVID-19 scale – psychometric study

C. Cabaços^{1,2*}, A.T. Pereira², P. Paredes³, S. Almeida³, A. Araujo^{1,2}, R. Sousa⁴ and A. Macedo^{1,2}

¹Psychiatry Department, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal; ²Institute Of Psychological Medicine, Faculty Of Medicine, University of Coimbra, Coimbra, Portugal; ³Faculty Of Medicine, University of Coimbra, Coimbra, Portugal and ⁴Uf Coimbra Centro, USF Coimbra Centro, Coimbra, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.696

Introduction: More than in other conditions, fear is associated with infectious diseases, and is directly associated with its transmission rate, morbidity and mortality. High levels of fear can affect the individual's ability to think clearly, react proportionately and make rational decisions

in the context of COVID-19. Recently, Mertens et al. (2020) developed the Fear of Covid-19 Scale (FCV-19S) to measure this construct.

Objectives: To analyse the psychometric properties of the FCV-19S Portuguese version, namely construct validity, internal consistency and convergent validity.

Methods: A community sample of 234 adults (75.6% women; mean age = 29.53 ± 12.51; range: 16-71) completed an on-line survey with the Portuguese versions of the FCV-19S, the Covid-19 Perceived Risk Scale (CPRS) and the Depression Anxiety Stress Scale (DASS-21). The total sample was randomly divided in two sub-samples: sample A (n=117) was used to perform an exploratory factor analysis/EFA; sample B (n=117) to make a confirmatory factor analysis/CFA.

Results: EFA resulted in one component. CFA revealed that the unifactorial model presented acceptable fit indexes ($X^2/df=3.291$; CFI=.977; GFI=.932; TLI=.919; $p[RMSEA \leq .01]=.091$). Cronbach alpha was $\alpha=.855$. The total score significantly correlated with Covid-19 Perceived Risk ($r=.529$, $p<.01$) and with anxiety from DASS-21 ($r=.132$, $p<.05$).

Conclusions: This study provides preliminary evidence for the validity and reliability of the Portuguese version of FCV-19S, which will be used in an ongoing research project on the relationship between fear of Covid-19, personality, cognitive processes and adherence to public health measures to contain the pandemic.

Keywords: COVID-19; Fear of COVID-19

EPP0289

Coping mechanisms and resilience in psychiatric trainees during COVID-19 pandemic

C.-A. Crisan^{1,2*}, R. Pop¹ and A. Mihai²

¹Psychiatric Clinic 1, Emergency County Hospital Cluj-Napoca, Cluj-Napoca, Romania and ²Neurosciences, Iuliu Hatieganu University of Medicine and Pharmacy Cluj-Napoca, Cluj-Napoca, Romania

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.697

Introduction: The Covid-19 pandemic has a profound impact on all domains of day to day life, forcing individuals to make substantial change in the way of living. Such change is known to cause an important psychological distress, and in some persons evidencing silent disorders among apparently functional individuals. Good coping mechanisms and resilience can be the key to overpass this difficult period.

Objectives: The aim of this study is to evaluate the coping mechanisms and resilience that Romanian psychiatric trainees used during Covid19 pandemic.

Methods: We developed an online questionnaire. We included questions about different socio-demographical variables and about coping mechanisms (using COPE scale), resilience (using Connor-Davidson Resilience Scale) and quality of life (using QoL Scale).

Results: The preliminary data show that staying busy, seeking social support and having a positive mindset are emotion-focused coping strategies present in individuals who overpass easier this period.

Conclusions: The Covid-19 pandemic is creating significant distress and impairment in functioning, but individuals who have good psychological mechanisms and who are more adaptable are less vulnerable during Covid-19 pandemic. Future research should build upon these findings to better understand coping mechanisms during crises and also social policies should be developed to acknowledge the variable needs in adults.

Keywords: coping mechanisms; resilience; Covid19 pandemic; stress

EPP0291

The experience of northern italy in elderly patients with COVID-19 infection and mental disorders

G. Serafini^{1*}, E. Bondi², C. Locatelli² and M. Amore³

¹Department Of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal And Child Health (dinogmi), University of Genoa, IRCCS Ospedale Policlinico San Martino, Genoa, Italy, Genoa, Italy;

²Department Of Psychiatry, Hospital Papa Giovanni XXIII, Bergamo, Italy, Bergamo, Italy and ³Department Of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal And Child Health (dinogmi), Dipartimento di Neuroscienze, Università di Genova, Genoa, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.698

Introduction: In December 2019, the first cases of Corona Virus Disease 2019 (COVID-19) outbreak related to acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection were reported in the Chinese city of Wuhan. European countries experienced a tragic growth in the number of Covid-19 cases although several restrictions have been imposed.

Objectives: The study is aimed to describe the first experience of the Hospital Papa Giovanni XXIII in the city of Bergamo, Northern Italy.

Methods: The most relevant clinical characteristics of aged patients with COVID-19 and mental disorders have been described.

Results: According to the experience of the Hospital Papa Giovanni XXIII, medical departments, after appropriate training of all healthcare workers, have been rapidly converted into specific units aimed at treating patients with COVID-19 infection. Specifically, we directly observed a rapidly growing request of psychiatric interventions in aged patients with COVID-19 infection due to the emergence of severe delirium (mainly hyperkinetic) which was reported in approximately 30–50% of cases increasing with age, psychomotor agitation, anxiety, and depressive symptoms. When compared with younger subjects, we found that subjects aged 65 or above with prolonged hospitalization in our hospital are more vulnerable to: 1) environmental factors (e.g., social isolation and distance from family members, stay in intensive/subintensive units, communication difficulties due to therapeutic devices); 2) individual factors (e.g., COVID-19 possible neurotropic properties, impairments in insight and cognitive dysfunctions, comorbid medical conditions, and use of multiple medications).

Conclusions: The main implications of the present findings have been discussed.

Keywords: COVID-19 infection; mental health; delirium; Psychological Distress

EPP0292

Mental health impact of the COVID-19 pandemic on the front-line health care workers in a tertiary care hospital in rawalpindi, pakistan

U. Zubair^{1*}, S. Mansoor¹ and T. Mansoor²

¹Oak, phoenix care center, Dublin, Ireland and ²Gender Studies, NUMS, islamabad, Pakistan

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.699