



Anti-inflammatory effects of ivy leaves dry extract: influence on transcriptional activity of NFκB

Janka Schulte-Michels¹ · Christina Keksel¹ · Hanns Häberlein¹ · Sebastian Franken¹

Received: 31 January 2018 / Accepted: 2 May 2018 / Published online: 11 May 2018
© The Author(s) 2018

Abstract

EA 575[®] is an ivy leaves dry extract (DER 5-7.5:1, 30% ethanol) used against diseases of the lower respiratory tract associated with productive cough. EA 575[®] improves symptoms associated with chronic inflammatory bronchial conditions. Compared to its bronchospasmolytic and secretolytic properties, the anti-inflammatory effects of EA 575[®] are mostly untried. Therefore, we addressed the question of whether the anti-inflammatory effect of EA 575[®] is due to an impact on the NFκB pathway. NFκB nuclear translocation was visualized by immunofluorescence in J774.2 as well as HEK293 cells. In the latter, a luciferase-based reporter was used to monitor NFκB transcriptional activity. Phosphorylation of RelA and its inhibitor IκB was measured by Western blot analysis. Additionally, changes in the stability of NFκB:IκB complex were shown by protein fragment complementation. Decreased transcriptional activity of NFκB under treatment with EA 575[®] was also shown for a human monocytic as well as a human lung epithelial cell line. EA 575[®] is able to inhibit NFκB transcriptional activity by partially inhibiting its translocation to the nucleus after stimulation with TNFα. Furthermore, phosphorylation of IκBα is reduced while phosphorylation of RelA is enhanced after pre-incubation with EA 575[®], leading to an enhanced stability of NFκB:IκBα complex. EA 575[®] has a regulatory impact on the NFκB pathway, possibly by switching specificity of IKK from IκBα to RelA, resulting in enhanced stability of NFκB:IκBα complex and reduced RelA translocation into the nucleus.

Keywords NFκB · *Hedera helix* · β₂-Adrenergic receptor · Airway inflammation · EA 575[®]

Introduction

EA 575[®] is an ivy leaves dry extract (DER 5-7.5:1, 30% ethanol) used for the improvement of symptoms associated with chronic inflammatory bronchial conditions and acute respiratory tract inflammation accompanied by coughing. Its bronchospasmolytic and secretolytic effects have been proven in clinical studies (Lang et al. 2015). The mechanism of action is based on increased β₂-adrenergic responsiveness of the respiratory tract mediated by inhibition of GRK2-mediated phosphorylation of activated β₂-adrenergic receptors, resulting in decreased internalization of β₂-adrenergic receptors under stimulating conditions (Sieben et al. 2009; Schulte-Michels et al. 2016b). The anti-inflammatory effect of EA 575[®] is less examined. As we have shown previously,

EA 575[®] is able to reduce LPS-induced IL-6 release from murine macrophages (J774.2) (Schulte-Michels et al. 2016a). Besides the findings from cell culture experiments, there is also growing evidence from animal models regarding the anti-inflammatory effect of constituents of *hedera helix* leaves. Rutin and chlorogenic acid reduce the carrageenan-induced paw edema in rats (Selloum et al. 2003; dos Santos et al. 2006). Rutin also inhibits acute as well as chronic arthritis states in rats (Guardia et al. 2001). Through a mouse model of ovalbumin-induced allergic asthma, it was demonstrated that chlorogenic acid reduces the synthesis of IgE and Th2-cell specific cytokines like IL-4 and IL-5 (Kim et al. 2010). Additionally, chlorogenic acid protects mice against LPS-induced lung injury (Zhang et al. 2010).

Hocaoglu et al. (2012) demonstrated that an ivy leaves dry extract showed anti-inflammatory properties in a chronic-induced asthma model in mice (Hocaoglu et al. 2012). Compared to placebo, goblet cell count was significantly lower and basal membrane thickness was reduced in lung histopathologic studies when mice were treated orally with the ivy leaves dry extract. Süleyman and coworkers also

✉ Sebastian Franken
sfranken@uni-bonn.de

¹ Institute of Biochemistry and Molecular Biology, Rheinische Friedrich-Wilhelms-University, Nussallee 11, 53115 Bonn, Germany

found an anti-inflammatory effect for an ivy extract enriched with saponins in cotton pellet and carrageenan-induced acute and chronic inflammation in rats (Süleyman et al. 2003).

An important aspect of nearly all inflammatory processes is the transcription factor NF κ B. Under non-stimulating conditions, NF κ B is kept inactive in the cytosol by complexation with its inhibitor I κ B α . Stimulation of, e.g., macrophages with TNF α or LPS activates the I κ B kinase (IKK), which in turn phosphorylates I κ B α , resulting in a cleavage of the NF κ B:I κ B α complex. The released NF κ B can thus be translocated into the nucleus, where it induces the transcription of different cytokines (Hayden and Ghosh 2012). Due to its relevance to inflammatory processes, we questioned if the reduced IL-6 release from EA 575[®] pre-treated murine macrophages could be caused by an impact on the NF κ B pathway.

In this work, we describe the influence of EA 575[®] on the transcriptional activity of NF κ B in different epithelial and immune system-derived cell lines using a Nanoluc reporter cell system. The nuclear translocation of NF κ B after TNF α -stimulation is examined by immunofluorescence microscopy; the phosphorylation of I κ B α and the stability of the NF κ B:I κ B α complex is investigated using Western blots and a protein fragment complementation assay, respectively. We aim to elucidate the anti-inflammatory mode of action of ivy leaves dry extract EA575[®], and to further understand its beneficial effect on chronic inflammatory bronchial conditions.

Materials and methods

Chemicals

Nano-Glo[®] Luciferase Assay was obtained from Promega (Promega GmbH, Mannheim). Ivy leaves dry extract EA 575[®] (CID-100048) was obtained from Engelhard (Engelhard Arzneimittel GmbH & Co. KG Niederdorfelden).

The reference substances rutin, hederacoside C, hederacoside D, and α -hederin were obtained from Phytolab (Vestenbergsgreuth, Germany).

All other reagents were obtained from Merck (Darmstadt, Germany), if not stated otherwise.

Characterization of EA 575[®] by HPLC analysis

Ivy leaves dry extract EA 575[®] (DER 5-7.5:1, 30% ethanol) (Batch number CID-100048) was obtained from Engelhard Arzneimittel GmbH & Co. KG (Niederdorfelden, Germany). Chemical composition and pharmacological properties of relevant ingredients have been reported previously (Sieben et al. 2009; Greunke et al. 2015; Schulte-Michels et al. 2016a, b). For further characterization, EA 575[®] was dissolved in 50% EtOH in a concentration of 40.8 mg/ml. EA

575[®] was analyzed on a Agilent Series 1200 HPLC system equipped with a degasser (G1322A), a quaternary pump (G1311A), an autosampler (G1329A) and a photodiode array detector (G1315D) using a LiChrospher RP-18 column (5 μ m, 125 \times 4 mm, Merck, Darmstadt). Solvent A was H₂O/acetonitrile (44/2, m/m) adjusted to pH 2.0 with phosphoric acid 85%. Solvent B was acetonitrile. The following linear gradient was used: 0–9 min 0% B, 9–10 min to 6% B, 10–25 min to 15% B, 25–50 min to 60% B, 50–51 min to 100% B, 51–65 min 100% B. Flow rate: 0–50, 51–65 min 1.5 ml/min. Detection: 205 nm. Identification of constituents was carried out by comparison of UV spectra of reference substances and their corresponding retention times. Chromatograms were registered and evaluated using Agilent Chemstation Software Version B.04.01.

Cell culture

The mouse macrophage cell line J774.2 was obtained from Sigma-Aldrich (Cat. Nr.: 85011428). J774.2 cells were cultured in DMEM without phenol red (Life Technologies, Carlsbad, CA) supplemented with 100 units/ml penicillin, 100 μ g/ml streptomycin, 2 mM L-Glutamin (Life Technologies) and 10% fetal calf serum (Life Technologies). Cells were maintained by ten-fold dilutions with fresh medium every 2–3 days.

Human embryonic kidney cells (HEK 293) obtained from DSMZ (No. ACC 305) (Braunschweig, Germany) were cultivated in DMEM medium (Life Technologies, Carlsbad, CA) supplemented with 100 units/ml penicillin, 100 μ g/ml streptomycin, and 10% fetal calf serum. Cells were maintained by threefold dilutions with fresh medium every 2–3 days.

Acute monocytic leukemia cells (THP-1) were obtained from DSMZ (No. ACC 16) and maintained in RPMI medium supplemented with 10% fetal calf serum, 100 units/ml penicillin, 100 μ g/ml streptomycin, 50 μ M β -mercaptoethanol and 5 ml Glutamax 100 \times (Life Technologies GmbH, Darmstadt). THP-1 cells were maintained by three-fold dilutions with fresh medium every 2–3 days.

Human lung epithelium-derived cells (A549) obtained from DSMZ (No. ACC 107) were cultivated in DMEM/F12 medium (Life Technologies, Carlsbad, CA) supplemented with 100 units/ml penicillin, 100 μ g/ml streptomycin, and 10% fetal calf serum. Cells were maintained by three-fold dilutions with fresh medium every 2–3 days.

Immunofluorescence

J774.2 or HEK293 cells were seeded on poly-D-lysine (PDL) coated coverslips in a density of 100,000 cells per well. At a cell density of 90% medium was changed to serum- and antibiotic-free medium. J774.2 cells were incubated with

80 µg/ml EA 575[®] HEK293 cells with 240 µg/ml EA 575[®] overnight. Serum-free medium served as a control. The next morning 100 ng/ml LPS and 25 ng/ml TNFα were added to J774.2 cells and HEK293 cells, respectively, and incubation was continued for another 50 min. Afterwards cells were fixed with 4% formaldehyde in PBS, permeabilized with ice-cold methanol and blocked with 5% BSA in PBS supplemented with 0.3% Triton X-100. Immunostaining was performed with RelA antibody (cell signaling technology, #6956) in a dilution of 1:800 overnight and a Cy3 coupled secondary antibody in a dilution of 1:400 for 1.5 h in the dark. Additionally, cells were incubated with Hoechst 33528 for nuclear staining. Images were taken on a Zeiss Axiovert 200 M with an LCI Plan-Neofluar 63×/1.30 Imm Korr Ph 3 M27 objective and an AxioCamMR.

Generation of plasmids for protein fragment complementation

Generation of protein fragment complementation plasmids, pCDNAHygr_cVenus-RelA and pCDNA zeo_IkBa-nVenus, was already described elsewhere (Yu et al. 2003) and was constructed by GeneArt (Thermo Fisher), according to the mentioned publication.

Generation of NFκB Reporter Plasmid

For the generation of a Nanoluciferase expression construct under control of NFκB binding sequence the pNF-KB-D2EGFP vector (Clontech) was used as a template. Destabilized GFP was removed from the vector by PCR introducing a new XhoI-site (forward primer: 5'TCGGATATCTCGAGC CGGAATTCGGGAAGCTTC-3'; reverse primer: 5'-GTT CAGGGGGAGGTGTG-3'). The open reading frame coding for destabilized Nanoluciferase was cut from pNL1.2[*NlucP*] vector (Promega) using BamHI/XhoI and introduced into the pNF-KB vector using the same restriction sites.

Transfection

HEK 293 cells were transfected by calcium phosphate method. Briefly, cells were seeded in 12 well plates and allowed to attach for at least 24 h. Before transfection medium was changed to 900 µl fresh medium. One µg DNA was mixed with 6.5 µl 2 M CaCl₂ and 50 µl sterile water. The dilution was added dropwise to 2×HBS and after half an hour of resting added to the cells. After 24 h, medium was changed to fresh DMEM containing antibiotic for selection (750 µg/ml geneticin).

THP-1 cells were transfected by Amaxa electroporation technology, Nucleofector[®] II. For each transfection 2× 10⁶ cells were diluted in a self-made appropriate buffer (5 mM KCl, 15 mM MgCl₂, 15 mM HEPES, 150 mM Na₂HPO₄/

NaH₂PO₄ pH 7.2, 50 mM NaCl) with 1 µg DNA and transfected with program T-020. Afterwards, they were transferred to fresh fully supplemented RPMI medium. Twenty-four hours later, medium was changed to full medium containing 700 µg/ml geneticin for selection with medium changes every 2–3 days. After 1 week of selection, surviving cells were tested for expression of the transgene.

A549 cells were transfected using branched polyethyl- enimine (PEI, # 408727, Sigma-Aldrich, Darmstadt, Germany). Therefore, 50,000 cells were seeded into the cavities of a 12-well plate. 2 µg of DNA was diluted into 40 µl of 150 mM NaCl and 3.3 µl of a 1 mg/ml PEI solution was added and the mixture was immediately vortexed for 10 s. After 10 min at RT, the DNA/PEI mixture was applied dropwise to the cells and incubated overnight. The next day medium was changed to fresh DMEM/F12 containing anti- biotic for selection (500 µg/ml geneticin).

Measurement of NFκB activation through a Nanoluciferase-based reporter system

Nanoluciferase expressing cells were seeded in 96-well LumiNunc[™] plates and were allowed to grow for 24 h. Medium was changed to serum-free medium before incubation. Incubation with EA 575[®] was performed with indicated concentrations of EA 575[®] for 8 h, respectively. After 5 h indicated concentrations of TNFα or LPS (THP-1 cells) were added for 3 h. Measurement of luciferase activity was performed using the Nano-Glo[®] Luciferase Assay system (Promega, Mannheim, Germany) according to the manufacturers' instructions.

Western blot analysis


HEK 293 cells were seeded in 12-well plates and allowed to grow to a confluency of 80–90% in full-growth medium. Before incubation, the medium was changed to serum-free medium. Cells were incubated with 240 µg/ml EA 575[®] for 8 h and with 25 ng/ml TNFα for 10 min. After the incubation, cells were harvested with 100 µl per well 10 mM HEPES buffer containing phosphatase inhibitor cocktail (Merck, Darmstadt, Germany) and protease inhibitor cocktail (Sigma-Aldrich, Darmstadt, Germany). Cell solutions were sonified for 15 s and lysed by being forced through a 30 G cannula. To separate cell debris, the solution was centrifuged at 15,000g for 5 min at 4 °C. Supernatant was diluted in 4× sample buffer and 20 µl of the samples was separated in a 10% acrylamide gel with 80 V in stacking and 120 V in resolving gel. Blotting was done with 200 mA per gel for 45 min. For normalization the membrane was stained after blotting with REVERT[™] total protein stain according to the manufacturers' instructions (LI-COR[®]). Intensity was measured on an LI-COR[®]

Odyssey reader in 700 nm channel. Afterwards, the stain was removed and membrane was blocked with Odyssey blocking buffer for 1 h. Immunostaining was performed using phospho-I κ B α antibody (Cells Signaling Technology, #9246) or phospho-RelA antibody (Cell Signaling Technology, #3033), respectively, in a dilution of 1:1000 in Odyssey blocking buffer diluted 1:1 with TBS and 0.1% Tween 20. Afterwards, membrane was washed three times in TBS/Tween 20 and incubated in secondary antibody (LI-COR[®] Goat anti Rabbit-DyLight 800, 1:10,000) diluted in Odyssey blocking buffer 1:1 with TBS, 0.1% Tween 20 and 0.01% SDS for 1 h. Membrane was washed three times in TBS and measured on an LI-COR[®] Odyssey Reader in 800 nm channel. Protein concentration was normalized through the band intensity pI κ B α /p-RelA divided by the normalization factor of REVERT[™] total protein stain.

Protein fragment complementation assay

HEK cells stably expressing I κ B α -nVenus and cVenus-RelA were seeded into a black 96-well Plate (CellCarrier[™], Perkin Elmer, Massachusetts, USA) in total growth medium and allowed to grow to a confluency of about 90%. After reaching confluency, medium was changed to HBSS buffer containing 20 mM HEPES and was pre-incubated with the indicated concentration of EA 575[®] for 5 h. Afterwards, cells were stimulated with 10 ng/ml TNF α and decrease of fluorescence intensity was measured for 3 h every 60 min in a Perkin Elmer EnSight[™] reader with excitation at 515 nm and emission at 550 nm. Additionally, cell confluency was measured in transmission channel. Data were registered and evaluated by Kaleido[™] software.

Fig. 1 HPLC fingerprint analysis of EA 575[®]. Ingredients of the ivy leaves dry extract were separated on an RP-18 column. Main components were identified by comparison to retention times and absorbance of reference standards. The substances identified are rutin, hederacoside C, hederacoside D, and α -hederin



Statistical data evaluation

Statistical data evaluation was performed with One-way ANOVA followed by Tukey multiple comparison test. Results were considered to be significant for p values of at least <0.05 .

Results

EA 575[®] analysis by HPLC

Composition of EA 575[®] (DER 5-7.5:1, 30% ethanol) was characterized by HPLC fingerprint analysis. The following substances could be identified by comparison of UV spectra and retention times of corresponding reference substances: rutin, hederacoside C, hederacoside D, and α -hederin (Fig. 1).

Partial inhibition of NF κ B translocation into the nucleus

As previously shown, pre-incubation with EA 575[®] reduces IL-6 secretion from murine macrophage cell line J774.2 (Schulte-Michels et al. 2016a). To examine whether this reduction is achieved by an inhibition of nuclear translocation of NF κ B, we performed an immunofluorescence assay on NF κ B subunit RelA in J774.2 cells. Incubation with 25 ng/ml TNF α for 50 min showed a considerable shift of NF κ B (detected by Cy3 conjugated secondary antibody fluorescence) from cytosol to the nucleus (Fig. 2b) when compared to cells only incubated in serum-free medium (Fig. 2a). Overnight pre-incubation with 80 μ g/ml EA 575[®] observably reduced the signal located in the nucleus after 50 min of stimulation (Fig. 2c). In parallel, nuclear translocation of NF κ B was examined in human embryonic kidney




Fig. 2 NFκB immunostained in red with anti-RelA primary antibody and Cy3 coupled secondary antibody shown in J774.2 (**a–c**) as well as HEK 293 cells (**d–f**). Nucleus stained in blue with Hoechst 33258. (**a+d**) unstimulated cells, (**b+e**) cells stimulated with 100 ng/ml LPS (J774.2) or 25 ng/ml TNFα (HEK293) for 50 min, and (**c+f**)

cells pre-incubated with 80 μg/ml (J774.2) or 240 μg/ml EA 575[®] (HEK293) overnight and subsequently stimulated with 100 ng/ml LPS (J774.2) or 25 ng/ml TNFα (HEK293) for 50 min. Scale bars correspond to 10 μm. Shown pictures are representative of 3 independent experiments

epithelial cells (HEK 293; Fig. 2d–f). Reduction of translocation was also detected in these cells, even though the effect was not as pronounced as in the macrophage cell line.

Nanoluciferase based NFκB reporter assay

As shown previously by El-Guendy and colleagues, NFκB transcriptional activity can be measured by an NFκB reporter gene assay (El-Guendy and Sinai 2008). Stably transfected HEK 293 Nano-PEST cells showed an almost 25-fold increase in Nanoluciferase activity after stimulation with 25 ng/ml TNFα for 3 h. This increase could be significantly and dose-dependently decreased by 31.0, 34.7, and 42.1% after 8 h of pre-incubation with EA 575[®] concentrations of 40, 80, and 240 μg/ml, respectively (Fig. 3).

Western blot against phosphorylated IκBα and RelA

The disintegration of the NFκB:IκBα complex is induced by the phosphorylation of IκBα mediated by IKKβ, which also phosphorylates NFκB subunit RelA. Therefore, we measured both phosphorylation of IκBα on Ser32 and phosphorylation of RelA at Ser536. Pre-incubation with 240 μg/ml EA 575[®] for 8 h was able to impede TNFα-induced phosphorylation of IκBα on Ser32 significantly (Fig. 4). Phosphorylation of RelA at Ser536 was increased after pre-incubation with 240 μg/ml EA 575[®] and stimulation with TNFα. A significant difference in RelA phosphorylation between ivy




Fig. 3 HEK 293 cells stably transfected with a Nanoluciferase based NFκB reporter plasmid were incubated for 8 h with the indicated amounts of EA 575[®] or solvent alone. After 5 h of pre-incubation cells were stimulated for further 3 h with 25 ng/ml TNFα or left unstimulated. Results represent the mean ± SD ($n \geq 30$, *** $p < 0.001$)

pre-treated and control cells could be seen after 10 min of TNFα incubation (Fig. 5).

Protein fragment complementation assay

Reduced IκBα phosphorylation is thought to be a stabilizing factor for the NFκB:IκBα complex. Therefore, we tested




Fig. 4 Phosphorylation of IκBα on Ser32 before and after stimulation with 25 ng/ml TNFα for 10 min and with or without pre-treatment with 240 μg/ml EA 575[®] for 8 h. Results represent the mean ± SD ($n \geq 8$, $**p < 0.01$)




Fig. 5 Phosphorylation of RelA on Ser536 (p-RelA) in HEK cells after stimulation with 25 ng/ml TNFα for different time periods with or without pre-treatment using 240 μg/ml EA 575[®]. **a** Quantitative analysis of western blot analysis. Results represent the mean ± SD ($n \geq 3$, $*p < 0.05$). **b** Representative result of Western Blot analysis of p-RelA

via a protein fragment complementation assay if pre-incubation with EA 575[®] enhances the stability of NFκB:IκBα under TNFα stimulating conditions. In this assay, binding of IκBα to NFκB results in the complementation of the fluorescent protein Venus. Disintegration of the complex in turn leads to a reduction of the detected Venus fluorescence

(Yu et al. 2003). As shown in Fig. 6, fluorescence intensity decreased by 47.4% after stimulation with 10 ng/ml TNFα for 3 h. After pre-incubation with 160 μg/ml EA 575[®] for 8 h, TNFα-induced fluorescence intensity was lowered to 60.8%, which represents 8.2% of conserved complex.

Effects of EA 575[®] on different cell types

To further evaluate the observed regulatory effect of EA 575[®], we tested THP-1 as a human monocytic cell line, as well as A549 cells derived from human lung epithelium. Cells were stably transfected with the Nanoluciferase-based NFκB reporter. As can be seen in Fig. 7, both cell types showed increased Nanoluciferase levels after application of TNFα, which is significantly reduced when cells were pre-incubated with 240 μg/ml EA 575[®]. Compared to cells incubated with TNFα only, inflammatory response was reduced to 75 and 88% for THP-1 and A549 cells, respectively.

Discussion

Ivy leaves dry extract EA 575[®] is well established for the improvement of symptoms associated with chronic inflammatory bronchial diseases, as well as for the treatment of acute inflammatory respiratory tract infections accompanied by coughing. Molecular mechanisms behind this anti-inflammatory activity of EA 575[®] have not yet been investigated satisfactorily.

EA 575[®] reduces the amount of secreted IL-6 from mouse macrophages (Schulte-Michels et al. 2016a). As can be derived from our work, this reduction is achieved via an




Fig. 6 Fluorescence intensity measurement of HEK 293 cells stably transfected with cVenus-RelA and IκBα-nVenus before and after stimulation with 10 ng/ml TNFα and with or without pre-treatment using 160 μg/ml EA 575[®] for 8 h. Results represent the mean ± SD ($n \geq 23$, $**p < 0.01$, $***p < 0.001$)




Fig. 7 NF κ B activity in THP-1 and A549 Nano-PEST cells after stimulation with 10 ng/ml LPS or 5 ng/ml TNF α , respectively, and with or without pre-treatment using 240 μ g/ml EA 575[®]. Data are shown as mean \pm SD of three independent experiments ($n \geq 16$, *** $p < 0.001$)

impaired nuclear translocation of NF κ B, which we displayed by an immunofluorescence assay against RelA, a subunit of NF κ B. Inhibition of NF κ B nuclear translocation by ivy leaves constituents, e.g., rutin as well as chlorogenic and dicaffeoylquinic acids, was already shown in mast cells and macrophages (Kim et al. 2008; Puangpraphant et al. 2011; Hwang et al. 2014). Using an NF κ B reporter gene assay, we could show that EA 575[®] inhibits dose-dependently the transcriptional activity of NF κ B. Decreased NF κ B activity after administration of ivy constituents was already shown. For example, rutin as well as 3,5- and 4,5-dicaffeoylquinic acid inhibits the NF κ B-dependent iNOS gene transcription, and as a result the synthesis of nitric oxide, whereas chlorogenic acid reduces LPS-induced NF κ B-dependent cyclooxygenase-2 (COX-2) expression in RAW 264.7 cells (Shan et al. 2009; Park et al. 2009; Kazłowska et al. 2010). Furthermore, 3,4-dicaffeoylquinic acid inhibits the phorbol-12-myristate-13-acetate-induced COX-2 expression in RAW264.7 cells by blocking the catalytic activity of JNK/p38 MAP kinases and the activation of C/EBP β and AP-1 (Han et al. 2010). The impairment of nuclear translocation and reduction in transcriptional activity of NF κ B might be caused by an enhanced stability of the NF κ B:I κ B α protein complex. To test this issue, we performed a protein fragment complementation assay where Venus protein fluorescence is achieved upon binding of I κ B α to NF κ B. After stimulation with 10 ng/ml TNF α , preserved Venus fluorescence was significantly higher when cells were pre-incubated with EA 575[®], thereby demonstrating an enhanced stability under TNF α stimulation. Complex stability is regulated by phosphorylation of I κ B α through IKK β , which forms together with IKK α and their regulating subunit IKK γ (also called NEMO) the IKK

complex. Phosphorylation of I κ B α on serine at position 32 (Ser32) leads to its ubiquitination and proteasomal degradation and thereby the release of NF κ B from the complex, permitting it to translocate into the nucleus (Scheidereit 2006). Thus, we were interested in whether the observed effects were due to reduced phosphorylation of I κ B α . A Western blot against I κ B α phosphorylated at Ser32 indicated a reduced phosphorylation at this site, which explains stabilization of the NF κ B:I κ B α complex as shown by protein fragment complementation (see above). Reduced cellular phosphorylation of I κ B α after incubation with natural products also existing in EA 575[®] was already demonstrated, as for chlorogenic acid on hepatic stellate cells (Shi et al. 2013). Besides I κ B α , IKK β also phosphorylates the NF κ B subunit RelA at Ser536. Indeed, we also found an impact on RelA-Ser536 phosphorylation which was increased after pre-incubation of cells with EA 575[®]. The phosphorylation of RelA is much less understood. Besides several phosphorylation sites which seem to have positive effects on RelA transcriptional activity, there is increasing evidence that the phosphorylation of Ser536 by IKK β impairs the nuclear translocation of RelA and thereby diminishes its transcriptional activity (Mattioli et al. 2004; Moreno et al. 2010; Christian et al. 2016). Schröfelbauer et al. showed that the phosphorylation of I κ B α on Ser32 depends on the scaffold function of NEMO and that inhibition of NEMO leads to a switch in its specificity towards RelA at Ser536, which is in good accordance with our observations (Schröfelbauer et al. 2012).

In addition to factors secreted by immune cells, those released by epithelial cells also promote inflammation of lung tissue (Mitchell and O'Byrne 2016, 2017; Green and Turner 2017). Consistent with that, the anti-inflammatory effect of EA 575[®], detected in monocytes/macrophages (THP-1/J774.2) and examined here in detail in HEK cells, can also be observed in A 549 cells, a human lung epithelial cell. The precise target cells of EA 575[®] in the inflamed lung should be the subject of future studies.

Taken together, our results indicate that the previously shown anti-inflammatory properties of EA 575[®] (Schulte-Michels et al. 2016a) result from a regulatory effect on the NF κ B pathway, which is probably due to a specificity switch of IKK β . As discussed above, this would reduce RelA transcriptional activity in a duplexed manner, firstly by stabilizing the complex with its inhibitor and secondly by reducing RelA translocation into the nucleus. Whether EA 575[®] achieves this by manipulating the scaffold function of the regulatory NEMO subunit, leading to stronger association of IKK β to RelA compared to I κ B α , must be subject of further investigations.

Acknowledgements This work was supported by a research grant of Engelhard Arzneimittel GmbH & Co.KG, Niederdorfelden, Germany.

Compliance with ethical standards

Conflict of interest S Franken is an employee of Engelhard Arzneimittel GmbH & Co.KG, Niederdorfelden, Germany. All other authors confirm that there is no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Christian F, Smith EL, Carmody RJ (2016) The regulation of NF- κ B subunits by phosphorylation. *Cells*. <https://doi.org/10.3390/cells5010012>
- dos Santos MD, Almeida MC, Lopes NP, de Souza GEP (2006) Evaluation of the anti-inflammatory, analgesic and antipyretic activities of the natural polyphenol chlorogenic acid. *Biol Pharm Bull* 29:2236–2240. <https://doi.org/10.1248/bpb.29.2236>
- El-Guendy N, Sinai AP (2008) Potential problems inherent in cell-based stable NF-kappaB-GFP reporter systems. *Mol Cell Biochem* 312:147–155. <https://doi.org/10.1007/s11010-008-9730-8>
- Green CE, Turner AM (2017) The role of the endothelium in asthma and chronic obstructive pulmonary disease (COPD). *Respir Res* 18:20. <https://doi.org/10.1186/s12931-017-0505-1>
- Greunke C, Hage-Hülsmann A, Sorkalla T et al (2015) A systematic study on the influence of the main ingredients of an ivy leaves dry extract on the β 2-adrenergic responsiveness of human airway smooth muscle cells. *Pulm Pharmacol Ther* 31:92–98. <https://doi.org/10.1016/j.pupt.2014.09.002>
- Guardia T, Rotelli AE, Juarez AO, Pelzer LE (2001) Anti-inflammatory properties of plant flavonoids. Effects of rutin, quercetin and hesperidin on adjuvant arthritis in rat. *Farmacologia* 56:683–687
- Han EH, Kim JY, Kim HG et al (2010) Inhibitory effect of 3-caffeoyl-4-dicaffeoylquinic acid from *Salicornia herbacea* against phorbol ester-induced cyclooxygenase-2 expression in macrophages. *Chem Biol Interact* 183:397–404. <https://doi.org/10.1016/j.cbi.2009.11.015>
- Hayden MS, Ghosh S (2012) NF- κ B, the first quarter-century: remarkable progress and outstanding questions. *Genes Dev* 26:203–234. <https://doi.org/10.1101/gad.183434.111>
- Hocaoglu AB, Karaman O, Erge DO et al (2012) Effect of *Hedera helix* on lung histopathology in chronic asthma. *Iran J Allergy Asthma Immunol* 11:316–323
- Hwang SJ, Kim Y-W, Park Y et al (2014) Anti-inflammatory effects of chlorogenic acid in lipopolysaccharide-stimulated RAW 264.7 cells. *Inflamm Res* 63:81–90. <https://doi.org/10.1007/s00011-013-0674-4>
- Kazłowska K, Hsu T, Hou C-C et al (2010) Anti-inflammatory properties of phenolic compounds and crude extract from *Porphyra dentata*. *J Ethnopharmacol* 128:123–130. <https://doi.org/10.1016/j.jep.2009.12.037>
- Kim HP, Park H, Son KH et al (2008) Biochemical pharmacology of biflavonoids: implications for anti-inflammatory action. *Arch Pharm Res* 31:265–273. <https://doi.org/10.1007/s12272-001-1151-3>
- Kim H-R, Lee D-M, Lee S-H et al (2010) Chlorogenic acid suppresses pulmonary eosinophilia, IgE production, and Th2-type cytokine production in an ovalbumin-induced allergic asthma: activation of STAT-6 and JNK is inhibited by chlorogenic acid. *Int Immunopharmacol* 10:1242–1248. <https://doi.org/10.1016/j.intimp.2010.07.005>
- Lang C, Rottger-Luer P, Staiger C (2015) A valuable option for the treatment of respiratory diseases: review on the clinical evidence of the Ivy Leaves Dry Extract EA 575(R). *Planta Med* 81:968–974. <https://doi.org/10.1055/s-0035-1545879>
- Mattioli I, Sebald A, Bucher C et al (2004) Transient and selective NF-kappa B p65 serine 536 phosphorylation induced by T cell costimulation is mediated by I kappa B kinase beta and controls the kinetics of p65 nuclear import. *J Immunol* 172:6336–6344
- Mitchell PD, O'Byrne PM (2016) Epithelial derived cytokines in asthma. *Chest*. <https://doi.org/10.1016/j.chest.2016.10.042>
- Mitchell PD, O'Byrne PM (2017) Biologics and the lung: TSLP and other epithelial cell-derived cytokines in asthma. *Pharmacol Ther* 169:104–112. <https://doi.org/10.1016/j.pharmthera.2016.06.009>
- Moreno R, Sobotzik J-M, Schultz C, Schmitz ML (2010) Specification of the NF-kappaB transcriptional response by p65 phosphorylation and TNF-induced nuclear translocation of IKK epsilon. *Nucleic Acids Res* 38:6029–6044. <https://doi.org/10.1093/nar/gkq439>
- Park KH, Park M, Choi SE et al (2009) The anti-oxidative and anti-inflammatory effects of caffeoyl derivatives from the roots of *Aconitum koreanum* R. RAYMOND. *Biol Pharm Bull* 32:2029–2033
- Puangpraphant S, Berhow MA, Vermillion K et al (2011) Dicaffeoylquinic acids in Yerba mate (*Ilex paraguariensis* St. Hilaire) inhibit NF- κ B nucleus translocation in macrophages and induce apoptosis by activating caspases-8 and -3 in human colon cancer cells. *Mol Nutr Food Res* 55:1509–1522. <https://doi.org/10.1002/mnfr.201100128>
- Scheidereit C (2006) I kappa B kinase complexes: gateways to NF-kappa B activation and transcription. *Oncogene* 25:6685–6705. <https://doi.org/10.1038/sj.onc.1209934>
- Schröfelbauer B, Polley S, Behar M et al (2012) NEMO ensures signaling specificity of the pleiotropic IKK β by directing its kinase activity toward I κ B α . *Mol Cell* 47:111–121. <https://doi.org/10.1016/j.molcel.2012.04.020>
- Schulte-Michels J, Runkel F, Gokorsch S, Häberlein H (2016a) Ivy leaves dry extract EA 575[®] decreases LPS-induced IL-6 release from murine macrophages. *Pharmazie* 71:158–161
- Schulte-Michels J, Wolf A, Aatz S et al (2016b) α -Hederin inhibits G protein-coupled receptor kinase 2-mediated phosphorylation of β 2-adrenergic receptors. *Phytomedicine* 23:52–57. <https://doi.org/10.1016/j.phymed.2015.12.001>
- Selloum L, Bouriche H, Tigrine C, Boudoukha C (2003) Anti-inflammatory effect of rutin on rat paw oedema, and on neutrophils chemotaxis and degranulation. *Exp Toxicol Pathol* 54:313–318. <https://doi.org/10.1078/0940-2993-00260>
- Shan J, Fu J, Zhao Z et al (2009) Chlorogenic acid inhibits lipopolysaccharide-induced cyclooxygenase-2 expression in RAW264.7 cells through suppressing NF-kappaB and JNK/AP-1 activation. *Int Immunopharmacol* 9:1042–1048. <https://doi.org/10.1016/j.intimp.2009.04.011>
- Shi H, Dong L, Dang X et al (2013) Effect of chlorogenic acid on LPS-induced proinflammatory signaling in hepatic stellate cells. *Inflamm Res* 62:581–587. <https://doi.org/10.1007/s00011-013-0610-7>
- Sieben A, Prenner L, Sorkalla T et al (2009) Alpha-hederin, but not hederacoside C and hederagenin from *Hedera helix*, affects the binding behavior, dynamics, and regulation of beta 2-adrenergic receptors. *Biochemistry* 48:3477–3482. <https://doi.org/10.1021/bi802036b>
- Süleyman H, Mshvildadze V, Gepdiremen A, Elias R (2003) Acute and chronic antiinflammatory profile of the ivy plant, *Hedera helix*, in rats. *Phytomedicine* 10:370–374. <https://doi.org/10.1078/0944-7113-00260>

Yu H, West M, Keon BH et al (2003) Measuring drug action in the cellular context using protein-fragment complementation assays. *Assay Drug Dev Technol* 1:811–822. <https://doi.org/10.1089/154065803772613444>

Zhang X, Huang H, Yang T et al (2010) Chlorogenic acid protects mice against lipopolysaccharide-induced acute lung injury. *Injury* 41:746–752. <https://doi.org/10.1016/j.injury.2010.02.029>