

# N-(3-Bromo-1,4-dioxo-1,4-dihydro-2-naphthyl)-4-fluoro-N-(4-fluorobenzoyl)-benzamide

Emmanuel S. Akinboye, Ray J. Butcher,\* Dwayne A. Wright, Yakini Brandy and Oladapo Bakare

Department of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA

Correspondence e-mail: rbutcher99@yahoo.com


Received 30 December 2008; accepted 2 January 2009

Key indicators: single-crystal X-ray study;  $T = 200$  K; mean  $\sigma(\text{C}-\text{C}) = 0.007$  Å;  $R$  factor = 0.056;  $wR$  factor = 0.158; data-to-parameter ratio = 13.2.

In the title compound,  $\text{C}_{24}\text{H}_{12}\text{BrF}_2\text{NO}_4$ , synthesized from 2-amino-3-bromo-1,4-naphthoquinone and 4-fluorobenzoyl chloride, the two *p*-fluorophenyl rings are inclined at 73.9 (1) and 73.6 (1)° to the naphthoquinone ring system. The two imido carbonyl O atoms are *anti* to each other, while the fluorophenyl rings are located opposite each other, connected to the imide group in a funnel-like arrangement. This conformation allows the fluorine groups be oriented slightly away from each other. An examination of the packing shows a close intermolecular  $\text{F}\cdots\text{O}$  contact of 2.982 (5) Å and a  $\text{Br}\cdots\text{O}$  contact of 2.977 (4) Å. In addition, the molecules are linked by weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{F}$  interactions.

## Related literature

For similar structures, see: Lien *et al.* (1997); Huang *et al.* (2005); Bakare *et al.* (2003); Akinboye *et al.* (2009); Win *et al.* (2005); Rubin-Preminger *et al.* (2004). For general background, see: Berhe *et al.* (2008).



## Experimental

### Crystal data

$\text{C}_{24}\text{H}_{12}\text{BrF}_2\text{NO}_4$	$V = 1979.88$ (6) Å <sup>3</sup>
$M_r = 496.26$	$Z = 4$
Monoclinic, $P2_1/n$	$\text{Cu } K\alpha$ radiation
$a = 14.5931$ (3) Å	$\mu = 3.30$ mm <sup>-1</sup>
$b = 6.6471$ (1) Å	$T = 200$ (2) K
$c = 20.6324$ (4) Å	$0.53 \times 0.48 \times 0.32$ mm
$\beta = 98.407$ (2)°	

### Data collection

Oxford Diffraction Gemini R diffractometer	7835 measured reflections
Absorption correction: multi-scan ( <i>CrysAlis RED</i> ; Oxford Diffraction, 2007)	3802 independent reflections
$T_{\min} = 0.230$ , $T_{\max} = 0.348$	3362 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.023$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$	289 parameters
$wR(F^2) = 0.158$	H-atom parameters constrained
$S = 1.14$	$\Delta\rho_{\text{max}} = 1.29$ e Å <sup>-3</sup>
3802 reflections	$\Delta\rho_{\text{min}} = -0.51$ e Å <sup>-3</sup>

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}5-\text{H}5\cdots\text{O}1\text{B}^{\text{i}}$	0.95	2.56	3.297 (6)	135
$\text{C}4-\text{H}4\cdots\text{F}1\text{A}^{\text{ii}}$	0.95	2.40	3.266 (6)	151

Symmetry codes: (i)  $-x, -y + 2, -z$ ; (ii)  $x + \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$ .

Data collection: *CrysAlisPro* (Oxford Diffraction, 2007); cell refinement: *CrysAlisPro*; data reduction: *CrysAlisPro*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

RJB acknowledges the NSF MRI program (grant No. CHE-0619278) for funds to purchase an X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2117).

## References

- Akinboye, E. S., Butcher, R. J., Brandy, Y., Adesiyun, T. A. & Bakare, O. (2009). *Acta Cryst. E65*, o24.
- Bakare, O., Ashendel, C. L., Peng, H., Zalkow, L. H. & Burgess, E. M. (2003). *Bioorg. Med. Chem.* **11**, 3165–3170.
- Berhe, S., Kanaan, Y., Copeland, R. L., Wright, D. A., Zalkow, L. H. & Bakare, O. (2008). *Lett. Drug Des. Disc.* **5**, 485–488.
- Huang, L., Chang, F., Lee, K., Wang, J., Teng, C. & Kuo, S. (2005). *Bioorg. Med. Chem.* **6**, 2261–2269.
- Lien, J., Huang, L., Wang, J., Teng, C., Lee, K. & Kuo, S. (1997). *Bioorg. Med. Chem.* **5**, 2111–2120.
- Oxford Diffraction (2007). *CrysAlisPro* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Rubin-Preminger, J. M., Win, T., Granot, Y. & Bittner, S. (2004). *Z. Kristallogr. New Cryst. Struct.* **219**, o323.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Win, T., Yerushalmi, S. & Bittner, S. (2005). *Synthesis*, p. 1631.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o277 [doi:10.1107/S1600536809000117]

## N-(3-Bromo-1,4-dioxo-1,4-dihydro-2-naphthyl)-4-fluoro-N-(4-fluorobenzoyl)benzamide

E. S. Akinboye, R. J. Butcher, D. A. Wright, Y. Brandy and O. Bakare

### Comment

We have developed some imido-substituted 2-chloro-1,4-naphthoquinones with cytotoxic activities on some cancer cell lines (Bakare *et al.*, 2003; Berhe *et al.*, 2008); and have recently reported on the crystal structure of *N*-(3-bromo-1,4-dioxo-1,4-dihydro-naphthalen-2-yl)-2-chloro-*N*-(2-chloro-benzoyl)-benzamide (Akinboye *et al.*, 2009). In continuation of our work, the title compound C<sub>24</sub>H<sub>12</sub>BrF<sub>2</sub>NO<sub>4</sub>, (I), was synthesized as a potential anticancer agent.

The crystal structure shows that the two *p*-fluorophenyl rings are inclined at 73.9 (1) and 73.6 (1) $^{\circ}$  to the naphthoquinone ring. The two imido carbonyl oxygen atoms are anti-to each other, while fluorophenyl rings are placed facing each other and connected to the imide moiety in a funnel-like arrangement. This conformation allowed the fluorine groups in the *para* position of each fluorophenyl ring to be oriented slightly away from each other. An examination of the packing shows a close contact between F1A and O2 at (1/2 -  $x$ , 1/2 +  $y$ , 1/2 -  $z$ ) (2.982 (5) $\text{\AA}$ ) and between C2 and O1B at (1/2 -  $x$ , -1/2 +  $y$ , 1/2 -  $z$ ) (2.977 (4) $\text{\AA}$ ). In addition, the molecules are linked by weak intermolecular C—H $\cdots$ O and C—H $\cdots$ F interactions (Table 1).

### Experimental

To a solution 2-amino-3-bromo-1,4-naphthoquinone (300 mg, 1.19 mmol) in dry THF was added NaH (68.64 mg 2.86 mmol) and the mixture was stirred for 15 minutes. 4-Fluoro-benzoylchloride (0.35 ml, 2.86 mmol) was added thereafter and this mixture was stirred at room temperature for 16–24 hr under argon. The solvent was removed *in vacuo* and the solid residue was dissolved in dichloromethane (40 ml). The resultant solution was washed with water (3  $\times$  15 ml), saturated NaCl solution (2  $\times$  15 ml) and dried over anhydrous magnesium sulfate. The solvent was removed *in vacuo* and the residue recrystallized from ethyl acetate to obtain a yellow solid (391.0 g m). Further recrystallization was carried out in ethanol to furnish the title imide (340.2 mg, 57%).

### Refinement

The methyl H atoms were constrained to an ideal geometry with C—H distances of 0.98  $\text{\AA}$  and  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ , but each group was allowed to rotate freely about its C—C bond. All other H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C—H distances in the range 0.95–1.00  $\text{\AA}$  and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. View of (I) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 20% probability level.

# supplementary materials

---




Fig. 2. View of the packing viewed down the  $a$  axis. Dashed bonds show weak C—H···F interactions as well as the close Br···O intermolecular contact.

## *N*-(3-Bromo-1,4-dioxo-1,4-dihydro-2-naphthyl)-4-fluoro-*N*-(4-fluorobenzoyl)benzamide

### Crystal data

$C_{24}H_{12}BrF_2NO_4$	$F_{000} = 992$
$M_r = 496.26$	$D_x = 1.665 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	$\text{Cu } K\alpha \text{ radiation}$
Hall symbol: -P 2yn	$\lambda = 1.54184 \text{ \AA}$
$a = 14.5931 (3) \text{ \AA}$	Cell parameters from 5844 reflections
$b = 6.6471 (1) \text{ \AA}$	$\theta = 4.0\text{--}73.4^\circ$
$c = 20.6324 (4) \text{ \AA}$	$\mu = 3.30 \text{ mm}^{-1}$
$\beta = 98.407 (2)^\circ$	$T = 200 (2) \text{ K}$
$V = 1979.88 (6) \text{ \AA}^3$	Prism, pale yellow
$Z = 4$	$0.53 \times 0.48 \times 0.32 \text{ mm}$

### Data collection

Oxford Diffraction Gemini R diffractometer	3802 independent reflections
Radiation source: fine-focus sealed tube	3362 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.023$
Detector resolution: 10.5081 pixels $\text{mm}^{-1}$	$\theta_{\text{max}} = 73.6^\circ$
$T = 200(2) \text{ K}$	$\theta_{\text{min}} = 4.0^\circ$
$\varphi$ and $\omega$ scans	$h = -17 \rightarrow 18$
Absorption correction: multi-scan (CrysAlis RED; Oxford Diffraction, 2007)	$k = -8 \rightarrow 5$
$T_{\text{min}} = 0.230$ , $T_{\text{max}} = 0.348$	$l = -25 \rightarrow 25$
7835 measured reflections	

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.056$	H-atom parameters constrained
$wR(F^2) = 0.158$	$w = 1/[\sigma^2(F_o^2) + (0.0639P)^2 + 6.5199P]$
$S = 1.14$	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} < 0.001$

3802 reflections  $\Delta\rho_{\max} = 1.29 \text{ e } \text{\AA}^{-3}$   
 289 parameters  $\Delta\rho_{\min} = -0.51 \text{ e } \text{\AA}^{-3}$   
 Primary atom site location: structure-invariant direct Extinction correction: none  
 methods

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.22095 (3)	0.70099 (8)	0.19146 (2)	0.0444 (2)
F1A	-0.2690 (2)	0.9709 (5)	0.41727 (16)	0.0599 (8)
F1B	0.1251 (3)	0.7110 (6)	0.52944 (16)	0.0752 (11)
O1	0.1976 (2)	0.7543 (6)	0.04619 (17)	0.0448 (8)
O2	-0.1290 (2)	0.8464 (5)	0.12723 (16)	0.0439 (8)
O1A	-0.0683 (2)	0.4826 (5)	0.21656 (17)	0.0479 (8)
O1B	0.0933 (2)	1.0688 (5)	0.24341 (15)	0.0424 (7)
N	0.0163 (2)	0.7707 (6)	0.22161 (18)	0.0358 (8)
C1	0.1109 (3)	0.7445 (7)	0.1341 (2)	0.0330 (9)
C2	0.1213 (3)	0.7515 (6)	0.0634 (2)	0.0335 (9)
C3	0.0344 (3)	0.7515 (6)	0.0152 (2)	0.0336 (9)
C4	0.0391 (3)	0.7337 (7)	-0.0509 (2)	0.0386 (10)
H4A	0.0976	0.7218	-0.0656	0.046*
C5	-0.0414 (4)	0.7333 (6)	-0.0961 (2)	0.0404 (10)
H5A	-0.0380	0.7208	-0.1415	0.048*
C6	-0.1272 (3)	0.7512 (7)	-0.0744 (2)	0.0395 (10)
H6A	-0.1824	0.7503	-0.1051	0.047*
C7	-0.1317 (3)	0.7702 (7)	-0.0086 (2)	0.0375 (10)
H7A	-0.1903	0.7830	0.0059	0.045*
C8	-0.0513 (3)	0.7706 (6)	0.0370 (2)	0.0333 (9)
C9	-0.0579 (3)	0.7970 (6)	0.1070 (2)	0.0343 (9)
C10	0.0287 (3)	0.7658 (6)	0.1550 (2)	0.0324 (9)
C1A	-0.0557 (3)	0.6469 (7)	0.2405 (2)	0.0372 (10)
C2A	-0.1127 (3)	0.7365 (7)	0.2871 (2)	0.0363 (9)
C3A	-0.1389 (3)	0.9372 (7)	0.2822 (2)	0.0389 (10)
H3AA	-0.1194	1.0199	0.2493	0.047*
C4A	-0.1936 (3)	1.0168 (8)	0.3254 (2)	0.0424 (10)
H4AA	-0.2131	1.1532	0.3222	0.051*

## supplementary materials

---

C5A	-0.2187 (3)	0.8922 (9)	0.3732 (2)	0.0457 (11)
C6A	-0.1944 (4)	0.6934 (8)	0.3788 (3)	0.0492 (12)
H6AA	-0.2135	0.6120	0.4122	0.059*
C7A	-0.1415 (3)	0.6137 (8)	0.3348 (2)	0.0444 (11)
H7AA	-0.1248	0.4754	0.3371	0.053*
C1B	0.0646 (3)	0.9152 (7)	0.2643 (2)	0.0354 (9)
C2B	0.0816 (3)	0.8578 (8)	0.3349 (2)	0.0382 (10)
C3B	0.0772 (3)	1.0046 (8)	0.3817 (2)	0.0452 (11)
H3BA	0.0641	1.1397	0.3685	0.054*
C4B	0.0918 (4)	0.9575 (10)	0.4476 (3)	0.0529 (13)
H4BA	0.0880	1.0576	0.4799	0.063*
C5B	0.1121 (4)	0.7597 (10)	0.4649 (3)	0.0544 (14)
C6B	0.1210 (3)	0.6130 (9)	0.4200 (3)	0.0517 (13)
H6BA	0.1377	0.4798	0.4337	0.062*
C7B	0.1051 (3)	0.6619 (8)	0.3541 (2)	0.0430 (11)
H7BA	0.1103	0.5616	0.3220	0.052*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br	0.0270 (3)	0.0572 (4)	0.0469 (3)	0.0056 (2)	-0.00134 (19)	0.0002 (2)
F1A	0.0519 (17)	0.068 (2)	0.0664 (19)	0.0020 (15)	0.0300 (15)	-0.0024 (16)
F1B	0.084 (3)	0.098 (3)	0.0405 (17)	-0.006 (2)	-0.0033 (16)	0.0140 (17)
O1	0.0291 (15)	0.057 (2)	0.0496 (19)	0.0045 (14)	0.0103 (13)	-0.0014 (16)
O2	0.0304 (15)	0.058 (2)	0.0429 (17)	0.0055 (15)	0.0043 (13)	-0.0034 (15)
O1A	0.0466 (19)	0.0395 (19)	0.059 (2)	-0.0065 (15)	0.0130 (16)	-0.0070 (16)
O1B	0.0404 (17)	0.0418 (18)	0.0429 (17)	-0.0065 (14)	-0.0012 (13)	-0.0001 (14)
N	0.0286 (17)	0.042 (2)	0.0366 (19)	-0.0023 (15)	0.0036 (14)	-0.0002 (15)
C1	0.0211 (18)	0.033 (2)	0.043 (2)	0.0010 (16)	-0.0030 (16)	-0.0005 (17)
C2	0.032 (2)	0.025 (2)	0.043 (2)	0.0023 (16)	0.0040 (17)	-0.0023 (17)
C3	0.033 (2)	0.026 (2)	0.041 (2)	0.0018 (17)	0.0048 (17)	0.0015 (17)
C4	0.044 (3)	0.029 (2)	0.043 (2)	0.0020 (19)	0.0075 (19)	0.0007 (18)
C5	0.056 (3)	0.023 (2)	0.041 (2)	0.0026 (19)	0.006 (2)	-0.0002 (17)
C6	0.045 (3)	0.027 (2)	0.043 (2)	-0.0013 (19)	-0.006 (2)	0.0010 (18)
C7	0.031 (2)	0.034 (2)	0.046 (2)	0.0018 (18)	0.0000 (18)	0.0016 (18)
C8	0.030 (2)	0.027 (2)	0.042 (2)	-0.0008 (16)	0.0012 (17)	0.0009 (17)
C9	0.027 (2)	0.033 (2)	0.042 (2)	-0.0026 (17)	0.0021 (17)	0.0000 (17)
C10	0.031 (2)	0.026 (2)	0.040 (2)	-0.0008 (16)	0.0012 (17)	0.0000 (16)
C1A	0.030 (2)	0.042 (3)	0.039 (2)	-0.0005 (19)	0.0017 (17)	0.0016 (19)
C2A	0.027 (2)	0.042 (2)	0.039 (2)	-0.0057 (18)	0.0015 (17)	0.0000 (18)
C3A	0.030 (2)	0.046 (3)	0.040 (2)	-0.0022 (19)	0.0029 (17)	0.0060 (19)
C4A	0.030 (2)	0.046 (3)	0.051 (3)	0.0027 (19)	0.0042 (19)	0.003 (2)
C5A	0.030 (2)	0.062 (3)	0.046 (3)	-0.003 (2)	0.0092 (19)	-0.002 (2)
C6A	0.045 (3)	0.051 (3)	0.055 (3)	-0.005 (2)	0.015 (2)	0.009 (2)
C7A	0.039 (2)	0.041 (3)	0.054 (3)	-0.004 (2)	0.009 (2)	0.004 (2)
C1B	0.0225 (18)	0.042 (3)	0.041 (2)	-0.0030 (17)	0.0015 (16)	-0.0013 (19)
C2B	0.0250 (19)	0.049 (3)	0.040 (2)	-0.0066 (19)	0.0008 (16)	0.000 (2)
C3B	0.036 (2)	0.050 (3)	0.049 (3)	-0.004 (2)	0.003 (2)	-0.002 (2)

C4B	0.046 (3)	0.069 (4)	0.043 (3)	-0.008 (3)	0.004 (2)	-0.007 (2)
C5B	0.040 (3)	0.083 (4)	0.039 (3)	-0.012 (3)	-0.001 (2)	0.010 (3)
C6B	0.035 (2)	0.062 (3)	0.056 (3)	0.002 (2)	-0.002 (2)	0.011 (3)
C7B	0.029 (2)	0.051 (3)	0.048 (3)	0.003 (2)	-0.0004 (18)	0.002 (2)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

Br—C1	1.873 (4)	C9—C10	1.503 (6)
F1A—C5A	1.353 (6)	C1A—C2A	1.486 (6)
F1B—C5B	1.356 (6)	C2A—C3A	1.387 (7)
O1—C2	1.217 (5)	C2A—C7A	1.390 (7)
O2—C9	1.218 (5)	C3A—C4A	1.385 (7)
O1A—C1A	1.201 (6)	C3A—H3AA	0.9500
O1B—C1B	1.207 (6)	C4A—C5A	1.378 (7)
N—C10	1.412 (6)	C4A—H4AA	0.9500
N—C1B	1.419 (6)	C5A—C6A	1.369 (8)
N—C1A	1.433 (6)	C6A—C7A	1.381 (7)
C1—C10	1.340 (6)	C6A—H6AA	0.9500
C1—C2	1.489 (6)	C7A—H7AA	0.9500
C2—C3	1.493 (6)	C1B—C2B	1.492 (6)
C3—C4	1.382 (6)	C2B—C3B	1.381 (7)
C3—C8	1.395 (6)	C2B—C7B	1.390 (7)
C4—C5	1.388 (7)	C3B—C4B	1.380 (7)
C4—H4A	0.9500	C3B—H3BA	0.9500
C5—C6	1.395 (7)	C4B—C5B	1.383 (9)
C5—H5A	0.9500	C4B—H4BA	0.9500
C6—C7	1.374 (7)	C5B—C6B	1.365 (9)
C6—H6A	0.9500	C6B—C7B	1.384 (7)
C7—C8	1.393 (6)	C6B—H6BA	0.9500
C7—H7A	0.9500	C7B—H7BA	0.9500
C8—C9	1.473 (6)		
C10—N—C1B	119.8 (4)	C7A—C2A—C1A	118.6 (4)
C10—N—C1A	117.0 (4)	C4A—C3A—C2A	119.9 (4)
C1B—N—C1A	122.5 (4)	C4A—C3A—H3AA	120.0
C10—C1—C2	122.5 (4)	C2A—C3A—H3AA	120.0
C10—C1—Br	122.5 (3)	C5A—C4A—C3A	118.0 (5)
C2—C1—Br	115.0 (3)	C5A—C4A—H4AA	121.0
O1—C2—C1	121.0 (4)	C3A—C4A—H4AA	121.0
O1—C2—C3	122.0 (4)	F1A—C5A—C6A	118.4 (5)
C1—C2—C3	117.0 (4)	F1A—C5A—C4A	118.4 (5)
C4—C3—C8	120.1 (4)	C6A—C5A—C4A	123.2 (5)
C4—C3—C2	119.9 (4)	C5A—C6A—C7A	118.6 (5)
C8—C3—C2	120.0 (4)	C5A—C6A—H6AA	120.7
C3—C4—C5	120.3 (4)	C7A—C6A—H6AA	120.7
C3—C4—H4A	119.8	C6A—C7A—C2A	119.7 (5)
C5—C4—H4A	119.8	C6A—C7A—H7AA	120.1
C4—C5—C6	119.7 (4)	C2A—C7A—H7AA	120.1
C4—C5—H5A	120.2	O1B—C1B—N	121.2 (4)
C6—C5—H5A	120.2	O1B—C1B—C2B	123.3 (4)

## supplementary materials

---

C7—C6—C5	120.0 (4)	N—C1B—C2B	115.4 (4)
C7—C6—H6A	120.0	C3B—C2B—C7B	119.9 (4)
C5—C6—H6A	120.0	C3B—C2B—C1B	119.0 (4)
C6—C7—C8	120.7 (4)	C7B—C2B—C1B	121.1 (4)
C6—C7—H7A	119.7	C4B—C3B—C2B	120.8 (5)
C8—C7—H7A	119.7	C4B—C3B—H3BA	119.6
C7—C8—C3	119.3 (4)	C2B—C3B—H3BA	119.6
C7—C8—C9	119.6 (4)	C3B—C4B—C5B	117.8 (5)
C3—C8—C9	121.1 (4)	C3B—C4B—H4BA	121.1
O2—C9—C8	123.4 (4)	C5B—C4B—H4BA	121.1
O2—C9—C10	119.2 (4)	F1B—C5B—C6B	118.9 (6)
C8—C9—C10	117.4 (4)	F1B—C5B—C4B	118.3 (5)
C1—C10—N	124.3 (4)	C6B—C5B—C4B	122.8 (5)
C1—C10—C9	120.6 (4)	C5B—C6B—C7B	118.7 (5)
N—C10—C9	115.0 (4)	C5B—C6B—H6BA	120.6
O1A—C1A—N	119.0 (4)	C7B—C6B—H6BA	120.6
O1A—C1A—C2A	124.3 (4)	C6B—C7B—C2B	119.9 (5)
N—C1A—C2A	116.6 (4)	C6B—C7B—H7BA	120.1
C3A—C2A—C7A	120.6 (4)	C2B—C7B—H7BA	120.1
C3A—C2A—C1A	120.8 (4)		
C10—C1—C2—O1	171.7 (4)	C1B—N—C1A—O1A	-150.5 (4)
Br—C1—C2—O1	-8.9 (6)	C10—N—C1A—C2A	-138.8 (4)
C10—C1—C2—C3	-9.3 (6)	C1B—N—C1A—C2A	31.9 (6)
Br—C1—C2—C3	170.1 (3)	O1A—C1A—C2A—C3A	-138.5 (5)
O1—C2—C3—C4	6.7 (7)	N—C1A—C2A—C3A	39.0 (6)
C1—C2—C3—C4	-172.3 (4)	O1A—C1A—C2A—C7A	39.8 (7)
O1—C2—C3—C8	-172.9 (4)	N—C1A—C2A—C7A	-142.8 (4)
C1—C2—C3—C8	8.1 (6)	C7A—C2A—C3A—C4A	0.5 (7)
C8—C3—C4—C5	-0.5 (6)	C1A—C2A—C3A—C4A	178.8 (4)
C2—C3—C4—C5	179.9 (4)	C2A—C3A—C4A—C5A	1.2 (7)
C3—C4—C5—C6	0.1 (7)	C3A—C4A—C5A—F1A	177.4 (4)
C4—C5—C6—C7	0.3 (7)	C3A—C4A—C5A—C6A	-1.8 (7)
C5—C6—C7—C8	-0.3 (7)	F1A—C5A—C6A—C7A	-178.6 (5)
C6—C7—C8—C3	-0.1 (7)	C4A—C5A—C6A—C7A	0.5 (8)
C6—C7—C8—C9	178.3 (4)	C5A—C6A—C7A—C2A	1.3 (8)
C4—C3—C8—C7	0.5 (6)	C3A—C2A—C7A—C6A	-1.8 (7)
C2—C3—C8—C7	-179.9 (4)	C1A—C2A—C7A—C6A	179.9 (4)
C4—C3—C8—C9	-177.8 (4)	C10—N—C1B—O1B	22.6 (6)
C2—C3—C8—C9	1.8 (6)	C1A—N—C1B—O1B	-147.9 (4)
C7—C8—C9—O2	-10.5 (7)	C10—N—C1B—C2B	-154.1 (4)
C3—C8—C9—O2	167.7 (4)	C1A—N—C1B—C2B	35.4 (6)
C7—C8—C9—C10	171.1 (4)	O1B—C1B—C2B—C3B	40.0 (6)
C3—C8—C9—C10	-10.6 (6)	N—C1B—C2B—C3B	-143.5 (4)
C2—C1—C10—N	-176.5 (4)	O1B—C1B—C2B—C7B	-137.4 (5)
Br—C1—C10—N	4.1 (6)	N—C1B—C2B—C7B	39.1 (6)
C2—C1—C10—C9	0.4 (6)	C7B—C2B—C3B—C4B	-3.1 (7)
Br—C1—C10—C9	-179.0 (3)	C1B—C2B—C3B—C4B	179.5 (4)
C1B—N—C10—C1	56.5 (6)	C2B—C3B—C4B—C5B	0.9 (7)
C1A—N—C10—C1	-132.5 (5)	C3B—C4B—C5B—F1B	-179.0 (5)

C1B—N—C10—C9	−120.6 (4)	C3B—C4B—C5B—C6B	2.1 (8)
C1A—N—C10—C9	50.5 (5)	F1B—C5B—C6B—C7B	178.2 (5)
O2—C9—C10—C1	−168.8 (4)	C4B—C5B—C6B—C7B	−2.9 (8)
C8—C9—C10—C1	9.6 (6)	C5B—C6B—C7B—C2B	0.7 (7)
O2—C9—C10—N	8.3 (6)	C3B—C2B—C7B—C6B	2.2 (7)
C8—C9—C10—N	−173.2 (4)	C1B—C2B—C7B—C6B	179.6 (4)
C10—N—C1A—O1A	38.7 (6)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C5—H5A···O1B <sup>i</sup>	0.95	2.56	3.297 (6)	135
C4—H4A···F1A <sup>ii</sup>	0.95	2.40	3.266 (6)	151

Symmetry codes: (i)  $-x, -y+2, -z$ ; (ii)  $x+1/2, -y+3/2, z-1/2$ .

## supplementary materials

---

Fig. 1




Fig. 2

