



Contents lists available at ScienceDirect

Journal of Ginseng Research

journal homepage: <http://www.ginsengres.org>

Research article

Highly regioselective biotransformation of ginsenoside Rb2 into compound Y and compound K by β -glycosidase purified from *Armillaria mellea* mycelia



Min-Ji Kim¹, Jitendra Upadhyaya¹, Min-Sun Yoon², Nam Soo Ryu², Young Eun Song³, Hee-Won Park⁴, Young-Hoi Kim¹, Myung-Kon Kim^{1,*}

¹ Department of Food Science and Technology, Chonbuk National University, Jeonju, Republic of Korea

² Department of Food Science and Biotechnology, Chonbuk National University, Iksan, Republic of Korea

³ Jeollabuk-do Agricultural Research and Extension Services, Iksan, Republic of Korea

⁴ Korea Ginseng Corporation Research Institute, Korea Ginseng Corporation, Daejeon, Republic of Korea

ARTICLE INFO

Article history:

Received 16 January 2017

Received in Revised form

27 June 2017

Accepted 3 July 2017

Available online 8 July 2017

Keywords:

Armillaria mellea
 β -glycosidase
 biotransformation
 compound Y
 ginsenoside Rb2

ABSTRACT

Background: The biological activities of ginseng saponins (ginsenosides) are associated with type, number, and position of sugar moieties linked to aglycone skeletons. Deglycosylated minor ginsenosides are known to be more biologically active than major ginsenosides. Accordingly, the deglycosylation of major ginsenosides can provide the multibioactive effects of ginsenosides. The purpose of this study was to transform ginsenoside Rb2, one of the protopanaxadiol-type major ginsenosides, into minor ginsenosides using β -glycosidase (BG-1) purified from *Armillaria mellea* mycelium.

Methods: Ginsenoside Rb2 was hydrolyzed by using BG-1; the hydrolytic properties of Rb2 by BG-1 were also characterized. In addition, the influence of reaction conditions such as reaction time, pH, and temperature, and transformation pathways of Rb2, Rd, F2, compound O (C-O), and C-Y by treatment with BG-1 were investigated.

Results: BG-1 first hydrolyzes 3-O-outer β -D-glucoside of Rb2, then 3-O- β -D-glucoside of C-O into C-Y. C-Y was gradually converted into C-K with a prolonged reaction time, but the pathway of Rb2 \rightarrow Rd \rightarrow F2 \rightarrow C-K was not observed. The optimum reaction conditions for C-Y and C-K formation from Rb2 by BG-1 were pH 4.0–4.5, temperature 45–60°C, and reaction time 72–96 h.

Conclusion: β -Glycosidase purified from *A. mellea* mycelium can be efficiently used to transform Rb2 into C-Y and C-K. To our best knowledge, this is the first result of transformation from Rb2 into C-Y and C-K by basidiomycete mushroom enzyme.

© 2017 The Korean Society of Ginseng, Published by Elsevier Korea LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Ginseng, the root of *Panax ginseng* Meyer, contains various bioactive components, such as ginsenosides, polyacetylenes, phenolic acids, and polysaccharides [1–3]. Among these compound classes, ginsenosides are known as the major active compounds, and dammarane-type ginsenosides in ginseng are divided into two groups according to attached position and number of hydroxyl groups in dammarane-type skeletons, i.e., protopanaxadiol (PPD)-type and protopanaxatriol-type [3,4]. A few ginsenosides including Rb1, Rb2, and Rc constitute nearly 80–90% of the total ginsenosides of ginseng [5]. However, the absorption

of these major ginsenosides in human intestine is known to be poor because of large molecular size, low water solubility of hydrophobic aglycones, and poor permeability through the cell membrane [6,7]. Meanwhile, the human intestinal bacterial metabolites, i.e., minor ginsenosides, such as compound O (C-O), C-Y, C-Mc, C-Mc1, and C-K, which are nearly absent in unprocessed ginseng, are more easily absorbed into the bloodstream [8,9]. Previous studies reported an attempt to convert major ginsenosides into minor ginsenosides by physical and chemical techniques [10–12], microbial [13,14] or enzymatic transformations [5,14]. In particular, minor ginsenosides could be prepared by crude microbial preparations or purified β -glycoside hydrolases

* Corresponding author. Department of Food Science and Technology, Chonbuk National University, 567 Baekje-daero, Deokjin-gu, Jeonju 54896, Republic of Korea.
 E-mail address: kmyuko@jbnu.ac.kr (M.-K. Kim).

from major ginsenosides, which were easily obtained from unprocessed extract [5,13].

Recently, biological effects of C-K, such as anti-inflammatory, anticarcinogenic, antiangiogenesis, antiaging, antidiabetic, and hepatoprotective activities, have been reported [14,15], but relatively little is known about the bioactivity of C-Y [16]. However, during microbial or enzymatic biotransformation of ginsenosides, it is difficult to stop the hydrolysis of Rb2 at the stage of C-O or C-Y formation because C-O or C-Y can be further transformed into C-K. Furthermore, the application of some ginsenoside-hydrolyzing β -glycosidases has a serious limitation in that they have poor activities against Rb2, which possess α -L-arabinopyranosidic linkage [13]. Therefore, it is very useful to search for Rb2-hydrolyzing β -glycosidases having a good regioselectivity and a high yield. Our attention has been focused on the transformation of major ginsenosides into minor ginsenosides by glycoside hydrolases secreted from basidiomycetes mycelia. In previous studies [17,18], we found that minor ginsenosides, C-K and C-Mc, could be efficiently produced from Rb1 and Rb2 by a crude enzyme extract or purified β -glycosidase from *Armillaria mellea* mycelium. Rb2 has α -L-arabinopyranosidic linkage different from Rb1 (β -D-glucosidic) and Rb3 (α -L-arabinofuranosidic) at the C-20 outer position of the aglycone. Therefore, we report on the biotransformation of Rb2 using β -glycosidase (BG-1) purified from *A. mellea* mycelium.

2. Materials and methods

2.1. Materials

Individual ginsenoside standards and the standard mixture (Rb1, Rb2, Rc, Rd, Rg3, F2, Rh2, C-K) were provided by the Korea Ginseng Corporation Research Institute (Daejeon, South Korea). C-Y and C-O were provided by Professor Tae-Hoo Yi (Department of Oriental Medicinal Materials and Processing College of Life Science, Kyung-Hee University, Yongin, South Korea). Ginsenoside Rb2 was also isolated from white ginseng roots [19]. *p*-Nitrophenol and *p*-nitrophenyl- β -D-glucopyranoside were purchased from Sigma-Aldrich Co. (St. Louis, MO, USA). Silica gel 60 (70–230 mesh) for

column chromatography and silica gel 60 F₂₅₄ TLC plates were purchased from Merck Co. (Darmstadt, Germany). HPLC-grade distilled water and acetonitrile were purchased from J.T. Baker (Center Valley, PA, USA). Other reagents (analytical grade) were purchased from Daejung Chemicals & Metals Co. (Siheung, Gyeonggi-do, South Korea).




Fig. 2. TLC analysis of reaction products of Rb2 and ammonium sulfate (30–80%) precipitates from mushroom mycelia. The reaction mixture contained Rb2 (2.0 mg) and ammonium sulfate (30–80%) precipitate containing 1.5 U as a β -glucosidase in 1.0 mL of 0.2M acetate buffer (pH 4.5), and distilled water to make a final volume of 2.0 mL was incubated for 96 h at 45°C, respectively. After heating for 10 min in boiling water, the reaction mixture was extracted with water-saturated *n*-butanol (2.0 mL \times 2) and concentrated. The residue was dissolved in methanol (1.0 mL). S, ginsenoside standard mixture; AM, *Armillaria mellea*; GL, *Ganoderma lucidum*; PB, *Phellinus baumii*; GA, *Ganoderma applanata*; PO, *Pleurotus ostreatus*.




Fig. 1. Two main pathways of the transformation of ginsenoside Rb2 by microbial glycosidases [13]. ara (p), α -L-arabinopyranosyl; C-K, compound K; C-O, compound O; C-Y, compound Y; Glc, β -D-glucopyranosyl.




Fig. 3. Time course of transformation of Rb2 by β -glycosidase (BG-1) purified from *Armillaria mellea* mycelium. The reaction mixture containing Rb2 (20 mg) and BG-1 (15 U) in 10.0 mL of 0.2M acetate buffer (pH 4.5), and distilled water to make a final volume of 20 mL was incubated for 96 h at 45°C. The reaction mixture (2.0 mL) was taken at different time intervals. After heating for 10 min in boiling water, the reaction mixture was extracted with water-saturated *n*-butanol (2.0 mL \times 2) and concentrated. The residue was dissolved in methanol (1.0 mL). S, ginsenoside standard mixture.

2.2. Production of mushroom mycelia and purification of β -glycosidase from *A. mellea* mycelium

The strains of five mushrooms—*A. mellea* (KACC 50013), *Ganoderma lucidum* (KACC 42231), *Phellinus baumii* (KACC 53719), *Ganoderma applanata* (KACC 53688), and *Pleurotus ostreatus* (KACC 50356)—were provided by the Korean Agricultural Culture Collection (Suwon, Gyeonggi-Do, South Korea). Unless otherwise stated, all procedures for mushroom mycelia production, preparations of crude enzymes (30–80% salt precipitates), and purification of β -glycosidase from *A. mellea* mycelium were performed according to the methods described in our previous studies [17,18].

2.3. Screening of ginsenoside Rb2-hydrolyzing activity using crude enzyme preparations from mushroom mycelia

A reaction mixture containing Rb2 (2.0 mg) as a substrate and each crude enzyme preparation (1.5 U as a β -glucosidase) containing β -glycosidase activity in 1.0 mL of 0.2M sodium acetate buffer (pH 4.5), and distilled water to make a final volume of 2.0 mL was incubated for 96 h at 45°C, respectively. After heating for 10 min in boiling water, the mixture was extracted with water-saturated *n*-butanol (2.0 mL \times 2). The extract was concentrated under reduced pressure, dissolved in methanol (1.0 mL).

2.4. Influence of reaction condition on transformation of ginsenoside Rb2 by BG-1

For investigation of the time course of transformation of Rb2 by BG-1, the reaction mixture containing Rb2 (20 mg), BG-1 enzyme




Fig. 4. HPLC analysis of the transformation products of Rb2 by BG-1 with different reaction times. The composition of the reaction mixture and the reaction conditions are the same as in Fig. 3. The reaction mixture was taken at different time intervals (0, 12, 48, and 96 h). C-K, compound K; C-O, compound O; C-Y, compound Y.

(15 U), 10 mL of 0.2M acetate buffer (pH 4.5), and distilled water to make a final volume of 20 mL was incubated for 96 h at 45°C. The mixture (2.0 mL) during reaction was taken at different time intervals and extracted with water-saturated *n*-butanol (2.0 mL × 2). The extract was concentrated under reduced pressure, dissolved in methanol (1.0 mL), and was subjected to TLC and HPLC analysis. Reaction mixture (2.0 mL) containing individual ginsenoside (Rd, F2, C-O, or C-Y) in methanol (0.2 mL) and BG-1 (1.5 U) in 1.8 mL of 0.1M acetate buffer (pH 4.5) were incubated for 96 h at 45°C. The reaction mixture was extracted with water-saturated *n*-butanol (2.0 mL × 2). The extract was concentrated to dryness under reduced pressure, and the residue was dissolved in methanol (1.0 mL).

The influence of temperature and pH on transformation of Rb2 by BG-1 was investigated under the following conditions: temperature range 30–70°C and pH range 4.0–9.0. Each reaction was incubated for 96 h at 45°C.

2.5. General analytical methods

β -Glucosidase activity was assayed as described by Mfombep et al. [20]. TLC, HPLC, and ultra-high performance liquid chromatography-quadrupole time-of-flight mass spectrometry (UPLC/Q-TOF-MS) analysis were performed as described in our previous studies [17,18]. The relative ratio of the transformation products in the reaction mixtures was calculated from the peak

area percentages in HPLC analysis without considering the detector response factor. HPLC analyses were performed in triplicate, and the results are expressed as the mean \pm standard deviation.

3. Results

3.1. Biotransformation of ginsenoside Rb2 by BG-1

Microbial β -glucosidases hydrolyze Rb2 through two different main pathways as shown in Fig. 1 [13]. One pathway is Rb2 \rightarrow Rd \rightarrow F2 \rightarrow C-K and another pathway is Rb2 \rightarrow C-O \rightarrow C-Y \rightarrow C-K. In preliminary experiments, when crude enzyme preparations (30–80% ammonium sulfate precipitate) obtained from five mushrooms mycelia were incubated with Rb2 as a substrate, high Rb2-hydrolyzing activity was only observed with enzyme preparation from *A. mellea*, whereas other enzyme preparations showed very weak hydrolytic activities against Rb2 (Fig. 2). Therefore, this result demonstrated that *A. mellea* has the potential to be used to produce minor ginsenosides from Rb2.

The sample was taken from the reaction mixture at different time intervals during a 96-h reaction to investigate the transformation pattern of Rb2 by BG-1, and analyzed by TLC and HPLC. The TLC profile (Fig. 3) showed that Rb2 was gradually transformed into three products having different R_f values with those of the authentic standards. Rb2 was transformed into product 1 in the




Fig. 5. MS spectra (negative ion mode) of the transformation products of Rb2 by BG-1. The reaction mixture (2.0 mL) contained Rb2 (2.0 mg) and BG-1 (1.5 U) in 2.0 mL of 0.1M sodium acetate buffer (pH 4.5) and was incubated for 24 h at 45°C. After heating for 10 min in boiling water, the reaction mixture was analyzed by UPLC/Q-TOF-MS. (A) C-O m/z 961.5005 $[M-H + HCOOH]^-$; (B) C-Y m/z 799.4680 $[M-H + HCOOH]^-$; (C) C-K m/z 667.4415 $[M-H + HCOOH]^-$. C-K, compound K; C-O, compound O; C-Y, compound Y.

initial stage of the reaction. After 72–96 h reaction, almost all of the Rb2 and product 1 were transformed into products 2 and 3 (Fig. 4). The mixture after 24 h reaction was analyzed by UPLC/Q-TOF-MS to determine the molecular weights of the products (Fig. 5). Product 1 showed a quasimolecular ion peak at m/z 961.5005 $[M-H + HCOOH]^-$ (molecular formula $C_{47}H_{80}O_{17}$). Product 2 showed a quasimolecular ion peak at m/z 799.4680 $[M-H + HCOOH]^-$ (molecular formula $C_{41}H_{70}O_{12}$), and product 3 showed a quasimolecular ion peak at m/z 667.4415 $[M-H + HCOOH]^-$ (molecular formula $C_{36}H_{62}O_{18}$). HPLC retention times and R_f values on TLC of products were also consistent with those of the authentic standards, C-O, C-Y, and C-K. These results suggested that BG-1 hydrolyzed β -sophorosyl moiety at the C-3 position of Rb2 to form C-Y via C-O, followed hydrolysis for C-K formation at the α -(1 \rightarrow 6)-arabinopyranosidic linkage at the C-20 position with a prolonged reaction time, but the transformation pathway of C-K formation via Rd and F2 from Rb2 was not observed.

3.2. Influence of reaction conditions and substrate specificity

The enzymatic transformation of ginsenosides was highly influenced by reaction conditions such as pH and temperature. When the influence of pH on transformation of Rb2 by BG-1 was





Fig. 6. Influence of pH on the transformation of Rb2 by BG-1. The reaction mixture (2.0 mL), containing Rb2 (2.0 mg) and BG-1 (1.5 U) in 2.0 mL of buffer solution (0.1M), was incubated for 96 h at 45°C at the given pH values (each at 0.1M); acetate (pH 4.0–5.5), sodium phosphate (pH 6.0 and 7.0), Tris-HCl (pH 8.0), and glycine-NaOH buffer (pH 9.0). Experiments were performed in triplicate, and results are shown as mean \pm standard deviation. BG-1, β -glycosidase; C-K, compound K; C-O, compound O; C-Y, compound Y.

investigated at broad pH range (pH 4.0–9.0), C-Y and C-K formation reached their maxima in the range of pH 4.0–4.5 (Fig. 6). These results indicate that the optimum pH range for the transformation of Rb2 by BG-1 is between pH 4.0 and 4.5. Then the influence of temperature was investigated in the range of 30–70°C. The transformation of Rb2 was maximized at 40–55°C (Fig. 7).

The Rb2 molecule contains two β -D-glucopyranosyl moieties at C-3 position and one α -L-arabinopyranosyl and one β -D-glucopyranosyl moiety at C-20 position of aglycone skeleton. Therefore, Rb2 can be hydrolyzed by enzymes via more than one pathway. The results from TLC and HPLC analysis showed that the biotransformation of Rb2 by BG-1 followed one pathway as shown in Figs. 3 and 4. BG-1 attacked the outer and inner β -glucosidic linkage attached to the C-3 position of aglycone to form C-O and C-Y. To investigate whether BG-1 showed the same specificity and




Fig. 7. Influence of temperature on the transformation of Rb2 by BG-1. The reaction mixture (2.0 mL), containing Rb2 (2.0 mg) and BG-1 (1.5 U) in 2.0 mL of 0.1M acetate buffer (pH 4.5), was incubated for 96 h at the given temperatures. Experiments were performed in triplicate, and results are shown as mean \pm standard deviation. BG-1, β -glycosidase; C-K, compound K; C-O, compound O; C-Y, compound Y.




Fig. 8. TLC analysis of transformation products of Rb2, Rd, C-O, F2, and C-Y by BG-1. The reaction mixture (2.0 mL), containing ginsenoside (each 2.0 mg) in methanol (0.2 mL) and BG-1 (1.5 U) in 1.8 mL of 0.1M acetate buffer (pH 4.5), was incubated for 96 h at 45°C, respectively. BG-1, β -glycosidase; C-K, compound K; C-O, compound O; C-Y, compound Y.

selectivity against glycosyl moieties attached to the C-3 and C-20 positions of other PPD-type ginsenosides, C-O, C-Y, Rd and F2 were also used as substrates. C-O was transformed into C-Y by hydrolysis of the glucosyl moiety at the C-3 position. When C-Y was used as a substrate, BG-1 was also gradually transformed C-Y into C-K by hydrolysis of the α -L-arabinopyranosidic linkage at the C-20 position (Fig. 8). But Rd was transformed into C-K by way of F-2, formed by hydrolysis of outer β -glucosyl linkage at the C-3 position (Fig. 9).

4. Discussion

By screening for mushrooms that have Rb2-hydrolyzing activity, we found that a crude enzymatic preparation from *A. mellea* mycelium can efficiently transform Rb2 into C-Y and C-K, whereas crude enzymes prepared from other mushroom mycelia showed very weak transformation activities against Rb2. These results

demonstrate that β -glycosidases from other mushroom mycelia used in this study could hardly hydrolyze the α -(1 \rightarrow 6)-L-arabinopyranosidic linkage at the C-20 position of Rb2. Previous studies reported that contents of minor ginsenosides in red ginseng extracts or its by-products could be enhanced by fermentation with some mushrooms [21–23]. However, the enhanced component in fermented products by mushrooms used in previous studies was Rd that was contained in unprocessed ginseng rather than minor ginsenosides. The medicinal fungus *Codyceps sinensis* was also shown to be capable of producing C-K from Rb1 [24].

Ginsenoside-hydrolyzing enzymes can be divided into four categories, based on the hydrolyzing characteristics of glycosidic linkages of ginsenoside molecules [25,26]. Interestingly, type III enzyme could selectively hydrolyze 3- β -sophorosyl and the outer 20- α -L-arabinopyranosidic linkages of Rb2 but not the inner 20- β -D-glucosidic linkage [26]. BG-1 also attacked the β -sophorosyl




Fig. 9. Proposed transformation pathways of Rb2, C-O, C-Y, Rd and F2 by BG-1. The Rf values on TLC of transformation products were compared with those of authentic standards. BG-1, β -glycosidase; C-K, compound K; C-O, compound O; C-Y, compound Y.

linkage bonded to the hydroxyl group at C-3 and the outer 20- α -L-arabinopyranosidic linkage at C-20 to form C-Y via C-O in Rb2. C-Y was also very slowly transformed into C-K over a prolonged reaction time. However, the use of a large quantity of enzyme (BG-1) may be necessary to rapidly transform C-Y into C-K, because BG-1 enzyme has a weak activity in hydrolyzing the outer 20- α -L-arabinopyranosidic linkage of the aglycone [18]. Meanwhile, when Rd was used as a substrate, BG-1 transformed it into C-K via F2. These results indicate that BG-1 has no hydrolyzing activity of the inner β -glucosidic linkage bonded to the hydroxyl group at C-20 position of Rb2. The transformation characteristics of BG-1 on Rb2 are similar to those of previously published studies on Rb2 transformation by β -glycosidases from gut bacteria [9,27], *Sulfolobus acidocaldarius* [16], *Microbacterium esteromiticum* [28], *Dictyoglomus turgidum* [29], *Penicillium aculeatum* [30], *Penicillium oxalicum* [31], *Sanguibacter keddiei* [32], recombinant *Sphingomonas* sp. [33], and crude enzyme extracts from *Aspergillus niger* and *Aspergillus usamii* [34].

The optimum pH range for transformation of Rb2 into C-Y and C-K by BG-1 was between 4.0 and 4.5. β -Glucosidase activities secreted by white rot fungi such as *A. mellea* exhibited pH optima between 3.5 and 5.0 [20]. The influence of pH on enzymatic transformation of ginsenosides has been extensively investigated using microbial enzymes isolated from various sources. These enzymes capable of transforming PPD-type ginsenosides showed optimal activities in the range of pH 4.0–6.0 [13,14]. BG-1 exhibited potent Rb2-transforming activity from 40°C to 55°C. Generally, the optimum temperatures for ginsenoside-hydrolyzing enzymes from human gut and soil bacteria are in the range of 37–45°C [13,14,33]. In recent years, even though many studies have reported the successful transformation of major ginsenosides into more active minor ginsenosides, such as Rg3, F2, Rh2, C-Mc, C-Y, and C-K, using various microbial enzymes, there have been relatively few studies on the biotransformation of major ginsenosides by β -glycosidases from basidiomycetes mushrooms. Interestingly, our results showed that BG-1 transformed Rb2 into C-Y and C-K via the Rb2 \rightarrow C-O \rightarrow C-Y and C-K pathway.

In conclusion, *A. mellea* β -glycosidase exhibited potent transformation activity against Rb2. The incubation conditions for optimum transformation of Rb2 into C-Y and C-K were a reaction time of 72–96 h, a pH of 4.0–4.5, and a temperature of 40–55°C. The pathway for C-Y and C-K formation from Rb2 was Rb2 \rightarrow C-O \rightarrow C-Y, but the pathway Rb2 \rightarrow G-Rd \rightarrow G-F2 \rightarrow C-K was not observed. C-Y was also slowly hydrolyzed at α -(1 \rightarrow 6)-arabinopyranosidic linkage at the C-20 position to form C-K given a prolonged reaction time. The results of the present study suggest that β -glycosidase purified from *A. mellea* mycelium can be efficiently used to transform Rb2 into C-Y and C-K.

Conflicts of interest

All contributing authors declare that they have no conflicts of interest.

Acknowledgments

This work was supported in part by a Korea Government Scholarship Program (J.U.).

References

- [1] Attele AS, Wu JA, Yuan CS. Ginseng pharmacology: multiple constituents and multiple actions. *Biochem Pharmacol* 1999;58:1685–93.
- [2] Kim DH. Chemical diversity of *Panax ginseng*, *Panax quinquefolium*, and *Panax notoginseng*. *J. Ginseng Res* 2012;36:1–15.

- [3] Christensen LP. Ginsenosides: chemistry, biosynthesis, analysis, and potential health effects. In: Taylor S, editor. *Adv Food Nutr Res*. Amsterdam: Elsevier; 2008. p. 1–99.
- [4] Park JD, Rhee DK, Lee YH. Biological activities and chemistry of saponins from *Panax ginseng* C. A. Meyer. *Phytochem Rev* 2005;4:159–75.
- [5] Jin F, Yu H, Fu Y, An DS, Im WT, Lee ST, da Silva JAT. Biotransformation of ginsenosides (ginseng saponins). *Int J Biomed Pharmacol Sci* 2012;6:33–44.
- [6] Xu QF, Fang XL, Chen DF. Pharmacokinetics and bioavailability of ginsenoside Rb1 and Rg1 from *Panax notoginseng* in rats. *J Ethnopharmacol* 2005;84:187–92.
- [7] Yu K, Chen F, Li C. Absorption, disposition, and pharmacokinetics of saponins from Chinese medicinal herbs: what do we know and what do we need to know more? *Curr Drug Metab* 2012;13:577–98.
- [8] Tawab MA, Bahr U, Karas M, Wurglics M, Schubert-Zsilavecz M. Degradation of ginsenosides in humans after oral administration. *Drug Metab Dispos* 2003;31:1065–71.
- [9] Hasegawa H. Proof of the mysterious efficacy of ginseng: basic and clinical trials: metabolic activation of ginsenoside: deglycosylation by intestinal bacteria and esterification with fatty acid. *J Pharmacol* 2004;95:153–7.
- [10] Kim WY, Kim JM, Han SB, Lee SK, Kim ND, Park MK, Kim CK, Park JH. Steaming of ginseng at high temperature enhances biological activity. *J Nat Prod* 2000;63:1702–4.
- [11] Han BH, Park MH, Han YN, Woo LK, Sankawa U, Yahara S, Tanaka O. Degradation of ginseng saponins under mild acidic conditions. *Planta Med* 1982;44:146–9.
- [12] Chen YJ, Nose M, Ogihara Y. Alkaline cleavage of ginsenosides. *Chem Pharm Bull* 1987;35:1653–5.
- [13] Park CS, Yoo MH, Noh KH, Oh DK. Biotransformation of ginsenosides by hydrolyzing the sugar moieties of ginsenosides using microbial glycosidases. *Appl Microbiol Biotechnol* 2010;87:9–19.
- [14] Yang XD, Yang YY, Ouyang DS, Yang GP. A review of biotransformation and pharmacology of ginsenoside compound K. *Fitoterapia* 2015;100:208–20.
- [15] Wong AST, Che CM, Leung KW. Recent advances in ginseng as cancer therapeutics: a functional and mechanistic overview. *Nat Prod Rep* 2015;32:256–72.
- [16] Noh KH, Oh DK. Production of the rare ginsenosides compound K, compound Y, and compound Mc by a thermostable β -glycosidase from *Sulfolobus acidocaldarius*. *Biol Pharm Bull* 2009;32:1830–5.
- [17] Upadhyaya J, Kim MJ, Kim YH, Ko SY, Park HW, Kim MK. Enzymatic formation of compound-K from ginsenoside Rb1 by enzyme preparation from cultured mycelia of *Armillaria mellea*. *J Ginseng Res* 2016;40:105–12.
- [18] Upadhyaya J, Yoon MS, Kim MJ, Ryu NS, Song YE, Kim YH, Kim MK. Purification and characterization of a novel ginsenoside Rc-hydrolyzing β -glycosidase from *Armillaria mellea* mycelia. *AMB Expr* 2016;6:112. 1–13.
- [19] Sanada S, Kondo N, Shoji J, Tanaka O, Shibata S. Studies on the saponins of ginseng. I. Structures of ginsenoside-Ro, -Rb1, -Rb2, -Rc, and -Rd. *Chem Pharm Bull* 1974;22:421–8.
- [20] Mfombep PM, Senwo ZN, Isikhuemhen OS. Enzymatic activities and kinetic properties of β -glycosidase from selected white rot fungi. *Adv Biol Chem* 2013;3:198–207.
- [21] Bae SH, Lee HS, Kim MR, Kim SY, Kim JM, Suh HJ. Changes of ginsenoside content by mushroom mycelial fermentation in red ginseng extract. *J Ginseng Res* 2011;35:235–42.
- [22] Hsu BY, Lu TJ, Chen CH, Wang SJ, Hwang LS. Biotransformation of ginsenoside Rd in ginseng extraction residue by fermentation with lingzhi (*Ganoderma lucidum*). *Food Chem* 2013;141:4186–93.
- [23] Ryu JS, Lee HJ, Bae SH, Kim SY, Park Y, Suh HJ, Jeong YH. The bioavailability of red ginseng extract of fermented by *Phellinus linteus*. *J Ginseng Res* 2013;37:108–16.
- [24] Wang WN, Yan BX, Xu WD, Qiu Y, Guo YL, Qiu ZD. Highly selective bioconversion of ginsenoside Rb1 to compound K by the mycelium of *Cordyceps sinensis* under optimized conditions. *Molecules* 2015;20:19291–9.
- [25] Wang D, Yu H, Song J, Xu Y, Jin F. Enzyme kinetics of ginsenosidase IV hydrolyzing 6-O-multi-glycosides of protopanaxatriol type ginsenosides. *Process Biochem* 2012;47:133–8.
- [26] Jin XF, Yu HS, Wang DM, Liu TQ, Liu CY, An DS, Im WT, Kim SG, Jin FX. Kinetics of a cloned special ginsenosidase hydrolyzing 3-O-glucoside of multi-protopanaxadiol-type ginsenosides, named ginsenosidase type III. *J Microbiol Biotechnol* 2012;22:343–51.
- [27] Bae EA, Park ST, Kim DH. Constitutive β -glycosidases hydrolyzing ginsenoside Rb1 and Rb2 from human intestinal bacteria. *Biol Pharm Bull* 2000;23:1481–5.
- [28] Quan LH, Jin Y, Wang C, Min JW, Kim YJ, Yang DC. Enzymatic transformation of the major ginsenoside Rb2 to minor compound Y and compound K by a ginsenoside-hydrolyzing β -glycosidase *Microbacterium esteromiticum*. *J Ind Microbiol Biotechnol* 2012;39:1557–62.
- [29] Lee GW, Kim KR, Oh DK. Production of rare ginsenosides (compound Mc, compound Y and aglycon protopanaxadiol) by β -glycosidase from *Dictyoglomus turgidum* that hydrolyzes β -linked, but not α -linked, sugars in ginsenosides. *Biotechnol Lett* 2012;34:1679–86.
- [30] Lee GW, Yoo MH, Shin KC, Kim KR, Kim YS, Lee KW, Oh DK. β -Glucosidase from *Penicillium aculeatum* hydrolyzes exo-, 3-O-, and 6-O- β -glucosides but not 20-O- β -glucoside and other glycosides of ginsenosides. *Appl Microbiol Biotechnol* 2013;97:6315–24.

- [31] Gao J, Xu W, Fang Q, Liang F, Jin R, Wu D, Tai G, Zhou Y. Efficient biotransformation for preparation of pharmaceutically active ginsenoside compound K by *Penicillium oxalicum* sp. 68. *Ann Microbiol* 2013;63:139–49.
- [32] Kim JK, Cui CH, Yoon MH, Kim SC, Im WT. Bioconversion of major ginsenoside Rg1 to minor ginsenoside F1 using novel recombinant ginsenoside hydrolyzing glycosidase cloned from *Sanguibacter keddieii* and enzyme characterization. *J Biotechnol* 2012;161:294–301.
- [33] Wang L, Liu QM, Sung BH, An DS, Lee HG, Im WT. Bioconversion of ginsenosides Rb1, Rb2, Rc, and Rd by novel β -glucosidase hydrolyzing outer 3-O glycoside from *Sphingomonas* sp. 2F2: cloning, expression, and enzyme characterization. *J Biotechnol* 2011;156:125–33.
- [34] Chi H, Kim DH, Ji GE. Transformation of ginsenosides Rb2 and Re from *Panax ginseng* by food microorganisms. *Biol Pharm Bull* 2005;28:2102–5.