



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Case report

First report of severe acute otitis media caused by *Campylobacter rectus* and review of the literature

Risako Kakuta <sup>a,\*</sup>, Hiroshi Hidaka <sup>a</sup>, Hisakazu Yano <sup>b</sup>, Michiko Okamoto <sup>c</sup>, Daiki Ozawa <sup>a</sup>, Shiro Endo <sup>d</sup>, Mitsuo Kaku <sup>d</sup>, Yukio Katori <sup>a</sup>

<sup>a</sup> Department of Otolaryngology, Head and Neck Surgery, Tohoku University Graduate School of Medicine, Sendai, Miyagi, Japan

<sup>b</sup> Department of Microbiology and Infectious Diseases, Nara Medical University, Kashihara, Nara, Japan

<sup>c</sup> Department of Virology, Tohoku University Graduate School of Medicine, Sendai, Miyagi, Japan

<sup>d</sup> Department of Infection Control and Laboratory Diagnostics, Tohoku University Graduate School of Medicine, Sendai, Miyagi, Japan

## ARTICLE INFO

## Article history:

Received 2 March 2016

Received in revised form

30 May 2016

Accepted 1 June 2016

Available online 28 June 2016

## Keywords:

Acute otitis media

*Campylobacter rectus*

16S ribosomal RNA gene

## ABSTRACT

*Campylobacter rectus* is a member of the human oral flora and is associated with periodontal disease. We report the first case of severe acute otitis media (AOM) due to *C. rectus* in a previously healthy 15-year-old boy, which was confirmed by 16S ribosomal RNA gene sequencing. *C. rectus* is a possible causative pathogen of AOM.

© 2016 Japanese Society of Chemotherapy and The Japanese Association for Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

## 1. Introduction

*Campylobacter rectus* is a Gram-negative, anaerobic, motile bacterium that is known to reside in the human oral cavity and is a causative agent of periodontitis [1,2]. *C. rectus* is also detected gastrointestinal tract with ulcerative colitis or Crohn's disease [3–5]. However, little is known regarding clinical relevance and pathogenic potential about this microorganism. In addition, extraoral infections caused by *C. rectus* have rarely been reported.

Acute otitis media (AOM) is one of the most common diseases in pediatric practice. Its pathogenesis involves complex interactions between bacteria, viruses, and the host inflammatory response. *Streptococcus pneumoniae* and *Haemophilus influenzae* colonize the nasopharynx, and are the most frequent causative pathogens of AOM [6]. Conversely, AOM caused by *C. rectus* has not been reported before. To the best of our knowledge, we are reporting the first case of culture negative severe AOM caused by *C. rectus* in a previously healthy adolescent.

## 2. Case report


In January 2014, a previously healthy 15-year-old boy presented to an ear, nose and throat (ENT) clinic with left-sided hearing loss. AOM was diagnosed and was treated with oral cefditoren-pivoxil for 7 days, followed by prulifloxacin for 4 days. He was then referred to the pediatrics department of a general hospital with worsening left otalgia accompanied by headache and hearing loss. After treatment with intravenous ceftriaxone for two days, he was transferred to the ENT department of Tohoku University Hospital for further evaluation.

On examination, his temperature was 37.0 °C. Otoloscopic examination revealed pulsating serous discharge from the left ear, along with circumferential bulging of the posterior wall of the left external auditory canal and thickening of the left tympanic membrane (Fig. 1a). The right ear was normal and there was no nystagmus or facial palsy. Neurological examination was unremarkable, and nuchal rigidity was absent. He had no history of previous ear discharge or significant illnesses. There was no evidence of periodontal disease or other active dental problems. He kept dogs and cats as pets.

Pure tone audiometry demonstrated conductive hearing loss on the left side (pure tone average: 47.5 dB) and normal hearing on the

\* Corresponding author. Department of Otolaryngology, Head and Neck Surgery, Tohoku University Graduate School of Medicine, 1-1 Seiryomachi, Aoba-ku, Sendai, 980-8574, Japan. Tel.: +81 22 717 7304; fax: +81 22 717 7307.

E-mail address: [kakuta-r@med.tohoku.ac.jp](mailto:kakuta-r@med.tohoku.ac.jp) (R. Kakuta).



**Fig. 1.** Tympanic membrane at the time of admission (a) and 12 days after discharge from our hospital (b). Pulsating serous discharge from the left ear, along with circumferential bulging of the posterior wall of the left external auditory canal and thickening of the left tympanic membrane (a). The left ear returned to normal (b).

right side. Laboratory tests revealed a white blood cell count of  $10.0 \times 10^3/\mu\text{l}$  with 79.1% neutrophils. C-reactive protein was increased to 2.2 mg/dl. Results of liver and renal function tests were within normal limits. Cerebrospinal fluid examination and magnetic resonance imaging revealed no evidence of meningitis at the previous hospital. In addition, aerobic culture of middle ear fluid performed at the previous hospital yielded negative results. Because Gram staining of middle ear fluid was negative, anaerobic culture had not been done in previous hospital.

Sampling of middle ear fluid for bacterial culture was done before the initiation of treatment at our hospital. Antimicrobial therapy was commenced with intravenous ampicillin (3.0 g/day). After treatment for 7 days, his symptoms resolved other than slight persistent left otorrhea. Oral amoxicillin (1000 mg/day) therapy was provided for 5 days after discharge. Twelve days after discharge from hospital, the left ear was normal (Fig. 1b) and the patient's symptoms had resolved (pure tone average: 6.3 dB).

Culture of middle ear secretions was done on sheep blood agar plates (Nissui Pharmaceutical Co., Tokyo, Japan) and chocolate agar (Kyokuto Pharmaceutical Co., Tokyo, Japan) under 5% CO<sub>2</sub>, on bromothymol blue lactose agar (Nippon Becton Dickinson Co. Ltd., Tokyo, Japan) and chromagar candida (Kanto Chemical Co., Inc., Tokyo, Japan) at 35 °C under aerobic conditions, on brucella HK agar (Kyokuto Pharmaceutical Co.) at 35 °C under anaerobic conditions (10% H<sub>2</sub>, 10% CO<sub>2</sub>, with 80% N<sub>2</sub>) for 48 h, and on HK semisolid agar (Kyokuto Pharmaceutical Co.) at 35 °C under aerobic conditions for 7 days. All of these cultures were negative.

### 3. Microbiological studies

#### 3.1. 16S ribosomal RNA gene sequencing and bacterial studies

To identify the causative organism, 16S ribosomal RNA (16S rRNA) gene sequencing was performed. Bacterial DNA was extracted from middle ear fluid using the QIAamp DNA Minikit (Qiagen, Hilden, Germany), after which polymerase chain reaction (PCR) amplification and 16S rRNA gene sequencing were performed as described previously [7]. We used the EzTaxon-e Database for sequence analysis (<http://eztaxon-e.ezbiocloud.net/>). The bacterial 16S rRNA gene sequence detected in middle ear fluid was 99.06% (1371/1385 bp) identical to that of the type strain *C. rectus* (ATCC33238, accession number: L04317), while concordance with other *Campylobacter* species was less than 97.27%. Thus, identification of *C. rectus* was confident. For investigation of the major pathogens of AOM, PCR was also performed for *S. pneumoniae*, *Haemophilus* spp., and *Streptococcus pyogenes* [8–10]. Results for all of these bacteria were negative.

#### 3.2. Viral studies

To determine the association of AOM with respiratory viral infection in this patient, viral studies were performed for the following viruses: influenza A/B virus, parainfluenza virus, adenovirus, cytomegalovirus, respiratory syncytial virus, enterovirus, rhinovirus, coronavirus, herpes simplex virus, human herpesvirus 6, human herpesvirus 7, varicella zoster virus, Epstein–Barr virus, human metapneumovirus, human bocavirus, WU virus, and KI virus. Viral nucleic acids were extracted from samples using the PureLink Viral RNA/DNA MiniKit (Invitrogen, Carlsbad, USA) according to the manufacturer's instructions. To synthesize complementary DNA, the final extract was used as the template with Moloney Murine Leukemia Virus and random hexamers (Invitrogen, Carlsbad, USA). Next, PCR or real-time PCR was performed as described previously with partial modification [11–16]. All of the viruses investigated were negative.

### 4. Discussion

To our knowledge, this is the first report of AOM caused by *C. rectus*. AOM is often considered to be a bacterial infection, but its pathogenesis actually involves complex interactions among viruses and bacteria [6]. In most children who develop AOM, viral infection of the upper respiratory tract initiates the cascade of events that finally leads to this condition [17]. Inflammation of the nasopharynx and Eustachian tube caused by viruses may allow bacteria to infect the middle ear. In the present patient, *C. rectus* could not be isolated by conventional culture at clinical laboratory in our hospital. However, we identified *C. rectus* by 16S rRNA gene sequencing and excluded the major bacterial and viral pathogens of AOM by PCR, RT-PCR, or real-time PCR. As a result, *C. rectus* was found to be the causative organism of severe AOM in this patient. *C. rectus* is difficult to culture and identify. It requires anaerobic conditions for optimal isolation, although the reported composition of the atmosphere needed for successful culture varies among authors [1]. Mahlene et al. obtained the isolate at the same culture conditions with us after 7 days incubation [1]. The supposable reasons why we could not obtain the isolate in spite of the same culture condition of the literature might be attributed to the prior administration of antimicrobial or short time cultivation time. Currently, identification of *C. rectus* is based on analysis of 16S rRNA. For clinical diagnosis of *C. rectus* infections, routine methods allowing accurate identification by microbiological laboratories are needed.

It has been reported that *C. rectus* is typically isolated from the oral cavity, and it was found in 50–94.6% of oral samples from children [2,18]. Familial transmission of oral bacteria is generally accepted [18] and it had been recently suggested that oral bacteria could also be transmitted between humans and their companion dogs. Kato et al. reported that 92.3% of pet dogs analyzed in Japan had *C. rectus* in their oral flora [19]. In addition, Yamasaki et al. found that 66.7% of companion dogs and 21.0% of their owners were positive for *C. rectus* in oral samples [20]. *C. rectus* was detected in fecal samples of healthy or diarrheic pet dogs [21], and the fecal-oral transmission might be one of the possible route. Because the present patient had dogs, it may be suggested that the origin of *C. rectus* infection was his companion pets. The route of spread was suspected to be from the oral cavity via the Eustachian tube to the middle ear. To clarify the relationship between *C. rectus* and AOM, further clinical and epidemiological studies among this organism, human and pet animals are needed.

Only seven cases of *C. rectus* infection have been reported in the English literature (Table 1). The present patient had several different characteristics from the previous 7 cases. First, all of the previous patients were adults, while this patient was an adolescent. Second,

**Table 1**  
Cases of *Campylobacter rectus* infection.

Author (year)	Age/Sex <sup>a</sup>	Diagnosis	Underlying diseases/ Risk factors	Specimen/Culture result	Identification method	Treatment	Outcome
1. Spiegel (1984) [23]	62/M	Left chest wall abscess	Poor oral hygiene, Alcoholism	Tissue/Positive	Biochemical features	Drainage, penicillin	Cured
2. Han (2005) [24]	32/F	Breast abscess	Lymphoma, neutropenia, Nipple piercing	Pus/Positive	16S rRNA gene analysis	Drainage, Vancomycin, Clindamycin, Aztreonam	Cured
3. de Vries (2008) [25]	24/M	Vertebral abscess	Meningoradiculitis of unknown cause	Pus/Positive	16S rRNA gene analysis	Hemilaminectomy, Clindamycin	Cured
4. Mahlen (2009) [1]	66/M	Buccal mucosal abscess	Smoking (30 years), Poor dental care	Pus/Positive	16S rRNA gene analysis	Drainage, Amoxicillin/Clavulanate	Cured
5. Lam (2011) [22]	41/F	Ruptured intracranial mycotic aneurysm with subdural empyema	Hyperglycemia	Blood, pus/Positive	16S rRNA gene analysis	Percutaneous arterial Embolization, Drainage, Vancomycin, Ceftriaxone, Metronidazole	Died
6. Lam (2011) [22]	64/M	Necrotizing soft tissue infection of right thigh	Poor oral hygiene	Tissue/Positive	16S rRNA gene analysis	Above-knee amputation, Surgical debridement, Amoxicillin/Clavulanate	Cured
7. Lam (2011) [22]	56/M	Right thoracic empyema	Poor oral hygiene, Hyperglycemia	Pus/Positive	16S rRNA gene analysis	Drainage, Amoxicillin/Clavulanate	Cured
8. This case	15/M	Acute otitis media	Kept dogs and cats	Middle ear fluid/Negative	16S rRNA gene analysis	Ampicillin, oral amoxicillin	Cured

<sup>a</sup> M: male, F: female.

the previous patients had underlying diseases, while this patient had no predisposing factors except for keeping pets. Third, culture was positive in all of the previous patients, while it was negative in this case, although this may have been related to prior antimicrobial therapy. We previously reported that 16S rRNA gene analysis was useful in patients with culture negative infection [7], and we employed this technique to identify the microorganism in the middle ear fluid of this patient. Although 7/8 cases, including this patient, responded to surgery and/or antimicrobial therapy, it is noteworthy that one patient died of ruptured intracranial mycotic aneurysm. Thus, *C. rectus* infections can occur in previously healthy young people and are occasionally fatal, and *C. rectus* is a potential causative pathogen of culture negative AOM. Therefore, further investigation of this microorganism in patients with AOM seems to be warranted.

Treatment of complicated infections due to *C. rectus* usually requires surgical drainage with prolonged antibiotic therapy [22]. There are limited antimicrobial susceptibility data for *C. rectus*, and there is no standard antibiotic regimen for infections due to this bacterium. One study showed that *C. rectus* was sensitive to imipenem, levofloxacin, amoxicillin-clavulanic acid, ceftiofloxacin, clindamycin, and metronidazole, with beta-lactamase activity not being detected [22]. The present patient recovered after receiving intravenous ampicillin and oral amoxicillin, so *C. rectus* may have been susceptible to penicillin in this case.

In conclusion, we described the first patient with AOM due to *C. rectus* infection detected by 16S rRNA gene analysis. This case suggests that *C. rectus* may be a possible causative pathogen of AOM. Accordingly, we need to perform culture while taking *C. rectus* into consideration in patients with AOM, and methods for its accurate identification by clinical microbiological laboratories are required. For adequate treatment of *C. rectus* infection, antibiotic selection guidelines and surveillance of its antimicrobial susceptibility profile are also required. Further clinical and epidemiological studies will be needed to clarify the relationship between *C. rectus* and AOM.

Nucleotide sequence accession number. The obtained sequence of the *Campylobacter rectus* isolate was deposited as DDBJ accession number LC147456.

All parts of the present study were performed in accordance with the guidance of the Declaration of Helsinki. Informed consent was obtained from the patient, and the review board of Tohoku University Hospital approved publication of this case report.

## Conflict of interest

There is no conflict of interest.

## References

- Mahlen SD, Clarridge 3rd JE. Oral abscess caused by *Campylobacter rectus*: case report and literature review. *J Clin Microbiol* 2009;47:848–51.
- Man SM. The clinical importance of emerging *Campylobacter* species. *Nat Rev Gastroenterol Hepatol* 2011;25(8):669–85.
- Mukhopadhyay I, Thomson JM, Hansen R, Berry SH, El-Omar EM, Hold GL. Detection of *Campylobacter concisus* and other *Campylobacter* species in colonic biopsies from adults with ulcerative colitis. *PLoS One* 2011;6:e21490.
- Man SM, Zhang L, Day AS, Leach ST, Lemberg DA, Mitchell H. *Campylobacter concisus* and other *Campylobacter* species in children with newly diagnosed Crohn's disease. *Inflamm Bowel Dis* 2010;16:1008–16.
- Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM. Global epidemiology of *Campylobacter* infection. *Clin Microbiol Rev* 2015;28:687–720.
- Pettigrew MM, Gent JF, Pyles RB, Miller AL, Nokso-Koivisto J, Chonmaitree T. Viral-bacterial interactions and risk of acute otitis media complicating upper respiratory tract infection. *J Clin Microbiol* 2011;49:3750–5.
- Kakuta R, Yano H, Kanamori H, Shimizu T, Gu Y, Hatta M, et al. *Helicobacter cinaedi* infection of abdominal aortic aneurysm, Japan. *Emerg Infect Dis* 2014;20:1942–5.
- Ubukata K, Asahi Y, Yamane A, Konno M. Combinational detection of autolysin and penicillin-binding protein 2B genes of *Streptococcus pneumoniae* by PCR. *J Clin Microbiol* 1996;34:592–6.
- Hasegawa K, Yamamoto K, Chiba N, Kobayashi R, Nagai K, Jacobs MR, et al. Diversity of ampicillin-resistance genes in *Haemophilus influenzae* in Japan and the United States. *Microb Drug Resist* 2003;9:39–46.
- Sunaoshi K, Nakayama E, Kobayashi R, Suzuki E, Tajima T, Ubukata K. Antibiotic susceptibility and T type identification of *Streptococcus pyogenes* isolated from pediatric outpatients with pharyngotonsillitis. *Jpn J Chemother* 2004;52:401–7.
- National Institute of Infectious Diseases. Manual of pathogenic organism. 2014. <http://www.nih.go.jp/niid/images/lab-manual/Influenza2014.pdf>.
- Suzuki A, Lupisan S, Furuse Y, Fuji N, Saito M, Tamaki R, et al. Respiratory viruses from hospitalized children with severe pneumonia in the Philippines. *BMC Infect Dis* 2012;12:267.
- Casas I, Avellon A, Mosquera M, Jabado O, Echevarria JE, Campos RH, et al. Molecular identification of adenoviruses in clinical samples by analyzing a partial hexon genomic region. *J Clin Microbiol* 2005;43:6176–82.
- Malasao R, Okamoto M, Chaimongkol N, Imamura T, Tohma K, Daput I, et al. Molecular characterization of human respiratory syncytial virus in the Philippines, 2012–2013. *PLoS One* 2015;10:e0142192.
- Tanaka T, Kogawa K, Sasa H, Nonoyama S, Furuya K, Sato K. Rapid and simultaneous detection of 6 types of human herpes virus (herpes simplex virus, varicella-zoster virus, Epstein-Barr virus, cytomegalovirus, human herpes virus 6A/B, and human herpes virus 7) by multiplex PCR assay. *Biomed Res* 2009;30:279–85.
- Matsuzaki Y, Mizuta K, Takahata E, Okamoto M, Itagaki T, Katsushima F, et al. Comparison of virus isolation using the Vero E6 cell line with real-time RT-

- PCR assay for the detection of human metapneumovirus. *BMC Infect Dis* 2010;10:170.
- [17] Marom T, Nokso-Koivisto J, Chonmaitree T. Viral-bacterial interactions in acute otitis media. *Curr Allergy Asthma Rep* 2012;12:551–8.
- [18] Umeda M, Miwa Z, Takeuchi Y, Ishizuka M, Huang Y, Noguchi K, et al. The distribution of periodontopathic bacteria among Japanese children and their parents. *J Periodontol Res* 2004;39:398–404.
- [19] Kato Y, Shirai M, Murakami M, Mizusawa T, Hagimoto A, Wada K, et al. Molecular detection of human periodontal pathogens in oral swab specimens from dogs in Japan. *J Vet Dent* 2011;28:84–9.
- [20] Yamasaki Y, Nomura R, Nakano K, Naka S, Matsumoto-Nakano M, Asai F, et al. Distribution of periodontopathic bacterial species in dogs and their owners. *Arch Oral Biol* 2012;57:1183–8.
- [21] Chaban B, Ngeleka M, Hill JE. Detection and quantification of 14 *Campylobacter* species in pet dogs reveals an increase in species richness in feces of diarrheic animals. *BMC Microbiol* 2010;10(10):73.
- [22] Lam JY, Wu AK, Ngai DC, Teng JL, Wong ES, Lau SK, et al. Three cases of severe invasive infections caused by *Campylobacter rectus* and first report of fatal *C. rectus* infection. *J Clin Microbiol* 2011;49:1687–91.
- [23] Spiegel CA, Telford G. Isolation of *Wolinella recta* and *Actinomyces viscosus* from an actinomycotic chest wall mass. *J Clin Microbiol* 1984;20:1187–9.
- [24] Han XY, Tarrand JJ, Rice DC. Oral *Campylobacter* species involved in extraoral abscess: a report of three cases. *J Clin Microbiol* 2005;43:2513–5.
- [25] de Vries JJ, Arents NL, Manson WL. *Campylobacter* species isolated from extra-oro-intestinal abscesses: a report of four cases and literature review. *Eur J Clin Microbiol Infect Dis* 2008;27:1119–23.