



Cite this article: Marco A. 2014 Sex-biased expression of microRNAs in *Drosophila melanogaster*. *Open Biol.* **4**: 140024.
<http://dx.doi.org/10.1098/rsob.140024>

Received: 11 February 2014

Accepted: 10 March 2014

Subject Area:

genomics/bioinformatics

Keywords:

evolution, sex, gene birth, demasculinization

Author for correspondence:

Antonio Marco

e-mail: amarco.bio@gmail.com

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsob.140024>.

Sex-biased expression of microRNAs in *Drosophila melanogaster*

Antonio Marco

School of Biological Sciences, University of Essex, Colchester CO4 3SQ, UK

1. Summary

Most animals have separate sexes. The differential expression of gene products, in particular that of gene regulators, is underlying sexual dimorphism. Analyses of sex-biased expression have focused mostly on protein-coding genes. Several lines of evidence indicate that microRNAs, a class of major gene regulators, are likely to have a significant role in sexual dimorphism. This role has not been systematically explored so far. Here, I study the sex-biased expression pattern of microRNAs in the model species *Drosophila melanogaster*. As with protein-coding genes, sex-biased microRNAs are associated with the reproductive function. Strikingly, contrary to protein-coding genes, male-biased microRNAs are enriched in the X chromosome, whereas female microRNAs are mostly autosomal. I propose that the chromosomal distribution is a consequence of high rates of de novo emergence, and a preference for new microRNAs to be expressed in the testis. I also suggest that demasculinization of the X chromosome may not affect microRNAs. Interestingly, female-biased microRNAs are often encoded within protein-coding genes that are also expressed in females. MicroRNAs with sex-biased expression do not preferentially target sex-biased gene transcripts. These results strongly suggest that the sex-biased expression of microRNAs is mainly a consequence of high rates of microRNA emergence in the X chromosome (male bias) or hitchhiked expression by host genes (female bias).

2. Introduction

Sexual dimorphism is prevalent in animal species. Sexual phenotypic differences are the consequence of a differential expression of genes between males and females [1]. During the past decade, high-throughput transcript analyses have identified many genes with a sex-biased expression pattern [2–4]. For instance, the *Drosophila* gene *paired* is expressed at a higher level in adult males than in females [5], and it encodes a transcription factor involved in the development of male accessory glands [6]. Indeed, other transcription factors have been identified as sex-biased genes [5,7], indicating that transcriptional gene regulation is tightly linked to sexual dimorphism. Post-transcriptional regulators may also have an impact in sexual dimorphism. MicroRNAs are short endogenous regulatory RNA molecules that are involved in virtually all studied biological processes [8,9]. Recently, differences between male and female microRNA expression profiles have been observed [10–12], suggesting that microRNAs have a role in sexual differentiation.

The study of sex-biased expression of gene products in the model species *Drosophila melanogaster* has produced a number of insightful observations. First, male-biased genes evolve faster than non-biased genes [3,13–15]. Second, the X chromosome is depleted of male-biased genes and enriched for female-biased genes [2,3]. On the other hand, evolutionarily novel genes tend to be X-linked and highly expressed in males [16–23]. These observations suggest a movement




Figure 1. Sex-biased microRNAs in *Drosophila melanogaster*. (a) Heatmap of cross-correlations of all expression datasets analysed. Different experiments are hierarchically clustered. (b) Smear plot of mature microRNA sequences. Grey lines indicate a twofold difference in expression levels between males and females. Red dots are microRNAs with a statistically significant differential expression. (c) MicroRNA transcripts with sex-biased expression, average fold change of their products and their chromosomal location. (d) Frequency plot of sex biases in expression levels for autosomes and the X chromosome.

of male-biased genes from the X chromosome to the autosomes, a process known as demasculinization of the X [2,20,24]. Therefore, sex-biased expression is an important factor affecting the evolutionary fate of protein-coding genes. Likewise, sex-biased expression should have an impact in microRNA evolution. However, this effect may be different to that observed for protein-coding genes as proteins and microRNAs differ in their evolutionary dynamics. For instance, gene duplication is the main mechanism by which novel protein-coding genes emerge, whereas a majority of microRNAs emerge by de novo formation within existing transcripts (reviewed in [25,26]). Consequently, novel microRNAs are more likely to be lost than protein-coding genes in a short evolutionary period. Although microRNAs have been extensively studied in *Drosophila* [27–29], the effect of sex-biased expression in microRNA evolution remains largely unexplored. Here, I investigate whether the sexual profile of microRNA expression resembles that of protein-coding genes, and how sex-biased expression affects differently the evolutionary dynamics of protein-coding genes and microRNAs.

3. Results

3.1. Sex-biased expression of *Drosophila* microRNAs

To characterize which microRNAs have a sex-biased expression pattern in *D. melanogaster*, 13 different small RNA sequencing

experiments (including males, females, embryos, ovaries and testes) were cross-compared (see §5). Figure 1a shows the correlation among the expression profiles for all experiments, indicating that the female and male pairs of profiles are highly correlated, despite coming from independent experiments. Thus, pairs of male and female profiles were used as biological replicates to calculate differential expression between sexes. A total of 476 mature microRNAs (two sequences per microRNA precursor) were analysed. Of them, 28 and 37 mature microRNAs showed a significant expression bias in males and females, respectively (figure 1b; see §5). Table 1 includes details of sex-biased microRNAs and their fold change. The expression levels for all analysed mature microRNAs are available in the electronic supplementary material, table S1. As only reads mapping to a single microRNA were taken into account, removing reads mapping to multiple sites may influence our analysis. Hence, I compared the expression levels resulting from unique reads and from multiple matching reads. Four microRNA families were affected by multiple matching reads: mir-983, mir-281, mir-276 and mir-2. The first three did not show any differential expression between sexes. However, the fourth one included two microRNAs, mir-2a-1 and mir-13b-2, which are female-biased. To avoid biases due to multiple matches, the mir-2 family was removed from the subsequent analyses.

MicroRNA precursors potentially encode for two mature products (so-called 3 prime and 5 prime products). In agreement with this, many of the sex-biased microRNAs are pairs

Table 1. MicroRNA mature sequences with sex-biased expression, and fold change shown in parentheses.

female-biased		male-biased		
mir-989-5p (9.8)	mir-995-5p (3.4)	mir-13b-2-5p (1.7)	mir-985-3p (8.8)	mir-978-3p (4.7)
mir-994-5p (8.7)	mir-313-3p (3.2)	mir-314-5p (1.7)	mir-976-3p (7.7)	mir-959-5p (4.5)
mir-989-3p (8.7)	mir-310-3p (3.1)	mir-306-5p (1.6)	mir-991-3p (7.2)	mir-960-3p (4.2)
mir-994-3p (8.2)	mir-279-5p (3.0)	mir-79-3p (1.4)	mir-977-5p (7.2)	mir-961-5p (4.1)
mir-318-3p (8.2)	mir-995-3p (2.9)	mir-996-5p (1.3)	mir-978-5p (6.8)	mir-303-5p (4.1)
mir-310-5p (7.8)	mir-79-5p (2.9)	mir-308-5p (1.1)	mir-4966-5p (6.6)	mir-984-5p (3.4)
mir-318-5p (6.7)	mir-9c-3p (2.5)	mir-996-3p (1.1)	mir-973-5p (5.9)	mir-959-3p (3.2)
mir-92a-3p (5.7)	mir-9c-5p (2.5)	mir-184-3p (1.0)	mir-975-5p (5.8)	mir-963-5p (3.0)
mir-313-5p (5.1)	mir-9b-5p (2.4)	mir-279-3p (1.0)	mir-982-5p (5.7)	mir-303-3p (3.0)
mir-92b-3p (5.0)	mir-312-5p (2.3)		mir-997-5p (5.5)	mir-964-5p (2.8)
mir-312-3p (4.3)	mir-9b-3p (2.2)		mir-972-3p (5.3)	mir-960-5p (2.5)
mir-311-3p (3.9)	mir-92a-5p (2.2)		mir-961-3p (5.1)	mir-2a-1-3p (1.5)
mir-92b-5p (3.6)	mir-998-3p (1.9)		mir-977-3p (5.0)	mir-993-3p (1.3)
mir-311-5p (3.4)	mir-2a-1-5p (1.8)		mir-992-3p (4.8)	mir-12-5p (1.1)

derived from the same precursor (table 1). Additionally, microRNAs are frequently clustered in the genome, and these clusters of microRNAs are often transcribed in a single RNA molecule (reviewed in reference [26]). Indeed, sex-biased microRNAs are frequently clustered, and nearly all of the microRNAs in a cluster show a consistent sex-biased expression (table 1 and figure 1c). Therefore, the observed bias in mature microRNA production is primarily a consequence of the sex-biased expression of their transcripts.

3.2. Male-biased microRNAs are preferentially located in the X chromosome and expressed in the testes

Figure 1c shows microRNA transcripts with sex-biased expression and their chromosomal distribution. Contrary to the observation for protein-coding genes, microRNAs expressed in males tend to be located in the X chromosome. By contrast, all female-biased microRNAs are located in autosomes, which is again the opposite observation to that which has been made for protein-coding genes. Figure 1d further explores the relationship between sex-chromosome location and sex-biased expression. The frequency distribution of fold change in expression for autosomal microRNAs shows three peaks, one large peak of unbiased expression and two smaller ones of male- and female-biased expression. However, the distribution of X-linked microRNAs is bimodal (figure 1d): they are either unbiased or highly expressed in males. Thus, male-biased microRNAs and the X chromosome are closely associated.

To further understand what it means to be sex-biased expressed, the expression profile of biased microRNAs was explored. Figure 2 plots a hierarchical tree of sex-biased expressed microRNAs and their relative expression levels in testes, ovaries and early embryos. Most male-biased microRNAs are highly expressed in the testes. This indicates that production of microRNAs in males is largely associated with the germline and the reproduction function. This is consistent with figure 1a in which adult samples were poorly correlated with young samples, perhaps because young individuals have not yet developed fully functional gonads.

3.3. Female-biased microRNAs are expressed in ovaries and early embryos

The expression profile in figure 2 shows that female-biased microRNAs fall into three distinct groups. First, a group of female-biased microRNAs are expressed in the somatic stem cells in the ovary, showing that microRNAs are important for the maintenance of stem cells in the ovary, in agreement with previous findings [30]. Second, some female-biased microRNAs are highly expressed in ovaries. This suggests that these microRNAs are important for the formation and maturation of *Drosophila* eggs.

Interestingly, a third group of female-biased microRNAs do not appear to be present in the ovary and they are highly expressed in young embryos (figure 2). These eggs were originally collected up to 1 h after laying [27,31], indicating that these young embryos have not yet started to have zygotic transcription [32]. This suggests that these microRNAs may be maternally deposited by the mother into the unfertilized eggs (oocytes). As a matter of fact, ongoing work in the laboratory has shown that the microRNAs mir-92a and mir-92b, and the mir-310/mir-311/mir-312/mir-313 cluster are abundant in *Drosophila* unfertilized eggs [33]. From these analyses, I conclude that both male- and female-biased microRNAs are mostly associated with the reproductive function.

3.4. Intronic female-biased microRNAs are associated with host gene expression

It may be possible that microRNA transcription pattern is associated with the transcription profile of their neighbouring protein-coding genes. In general, as shown in table 2, both expression patterns were not significantly associated (11 of 19 microRNA transcripts have the same expression bias as their closest neighbouring gene; $p = 0.32$, binomial test). In particular, there are eight microRNA transcripts with male-biased expression, and only three of their respectively closest genes show a similar bias. In the case of female-biased microRNA transcripts, eight




Figure 2. Expression profile of sex-biased microRNAs. Hierarchical clustering and heatmap of microRNAs with sex-biased expression. Z-scores were scaled across rows. Green colour indicates an overexpression in a given tissue/sample with respect to the other samples (columns). SSC, somatic stem cells.

of 11 have their closest gene with a female-biased expression pattern. A closer inspection to the data reveals that this bias is produced mostly by microRNAs hosted within protein-coding genes (overlapping transcripts). Indeed, all six genes hosting microRNAs with female-biased expression are themselves expressed more highly in females than in males ($p = 0.016$). This shows that female-biased expression of microRNAs is highly associated with the production of microRNAs from introns of sex-biased expressed protein-coding genes.

3.5. Evolutionary origin of sex-biased microRNAs

There are two possible ways a gene may become sex-biased. First, a gene can acquire sex-biased expression. Second, a new gene appears (either de novo or by the duplication of an existing gene) having from the very beginning a sex-biased expression. Figure 3 shows the evolutionary origin of sex-biased microRNAs. Most male-biased microRNAs emerged within the *Drosophila* lineage, with only two exceptions:

mir-993 and mir-283/304/12. These are indeed the least biased of all of the microRNAs. Thus, microRNAs with a strong male bias are evolutionarily young. By contrast, the evolutionary origin of female-biased microRNA families is diverse, and there are both old and young microRNAs. Among the old microRNAs, we have the mir-92, mir-184 and mir-9 families, which are conserved even in chordates. Interestingly, there are no *D. melanogaster*-specific microRNAs with a clear female-biased expression (contrary to the case of male-biased microRNAs). There are, however, two female-biased microRNAs which appeared in the *Drosophila* genus lineage: mir-314 and the mir-310–mir-313 cluster.

3.6. Targets of sex-biased microRNAs

Do sex-biased microRNAs also target sex-biased expressed gene transcripts? To explore this question, three different target prediction algorithms were used: TargetScan, miRanda and DIANA-microT (see §5). MicroRNAs were binned by their




Figure 3. Evolutionary origin of sex-biased microRNA transcripts. Phylogenetic tree of *Drosophila melanogaster* and other animal groups. MicroRNAs emerging at a given lineage were shown over the relevant branches. For microRNA clusters, only the first microRNA is shown in the figure over the branch at which the oldest microRNA emerged. Red microRNAs are female-biased and blue are male-biased.

Table 2. Sex-biased microRNA transcripts and their closest neighbouring genes.

microRNA/cluster	fold change ^a	distance to closest gene ^b	closest gene	fold change ^a
mir-972–mir-979	–5.2	overlapping	Grip84	2.3
mir-982–mir-984	–3.5	overlapping	CG3626	0.4
mir-959–mir-964	–2.8	overlapping	CG31646	–2.6
mir-283–mir-12	–1.1	overlapping	Gmap	–0.3
mir-985	–6.7	19 344	disco	–1.5
mir-997	–5.3	737	D1	1.2
mir-2498–mir-992	–4.7	817	CG32532	0.0
mir-993	–1.2	11 072	Ama	2.0
mir-92a–mir-92b	4.6	overlapping	jigr1	2.6
mir-995	3.0	overlapping	cdc2c	3.9
mir-9c–mir-9b	2.2	overlapping	grp	3.1
mir-184	1.1	overlapping	CG44206	0.0
mir-308	1.0	overlapping	RpS23	1.2
mir-998–mir-11	0.4	overlapping	E2f	1.4
mir-989	8.6	2739	Rcd1	0.9
mir-994–mir-318	8.3	249	lrp-1B	–0.3
mir-310–mir-313	4.1	1624	gsm	–0.7
mir-279–mir-996	1.1	2892	Ef1gamma	1.4
mir-314	0.2	182	Tim13	–7.5

^aLogarithm of fold change between male and female expression levels.

^bIn nucleotides.

bias level, and the expression bias of their targets was plotted in figure 4. These boxplots show that there is no tendency of sex-biased microRNAs to target sex-biased transcripts, at least not as a global pattern. I further explored the targets of seven *Drosophila*-subgroup-specific male-biased microRNAs. For two of them, two of the three prediction algorithms detected a significant association with sex-biased transcripts: mir-985 has a tendency to target female-biased genes, whereas mir-997 significantly targets male-biased genes (electronic supplementary material, table S2). The other associations were not significant and/or supported by only one prediction algorithm. Finally, I investigated whether recently emerged male-biased microRNAs also target evolutionarily young genes. I calculated the ratio between *Drosophila*-specific and conserved targeted genes for the targets predicted for the three above-mentioned algorithms. Four of seven studied microRNAs showed a tendency to

target more conserved genes than expected by chance for at least two algorithms (electronic supplementary material, table S3), among them mir-985. The results here described rely heavily on target prediction algorithms and, therefore, should be taken with caution. However, they suggest that newly emerged microRNAs can potentially target conserved genes, altering regulatory relationships that have been conserved throughout evolution.

4. Discussion

In this study, I have shown that sex-biased microRNAs are mainly associated with the reproductive function: male microRNAs are expressed in testes, and female microRNAs are abundant in ovaries and oocytes. However, their evolutionary




Figure 4. Expression bias of microRNAs and their targets. Box plots of expression bias of gene transcripts targeted by microRNAs with no (0), moderate (−5/5) and large (−10/10) sex-biased expression. Targets are shown for three different target prediction algorithms: (a) TargetScan, (b) DIANA-microT and (c) miRanda.

origin is different. Male-biased microRNAs tend to be evolutionarily young (dipteran/*Drosophila*-specific; figure 3) and they often emerge in the X chromosome. Contrary to microRNAs, male-biased protein-coding genes appear to be generally under-represented in the X chromosome in flies [2,3], and a movement of male genes out of the X, or demasculinization of the sex chromosome, has been suggested [2,20]. However, novel genes tend to be X-linked, and male-expressed and older genes may have moved outside the X chromosome [2,3,17,21]. An enrichment in the X chromosome for microRNAs with male-biased expression has also been reported in mammals [12,34–36].

A careful dissection of the evolutionary origin of male-biased genes in *Drosophila* demonstrated that de novo originated genes tend to be X-linked and male-biased, and that there may also be an ongoing demasculinization process in the X chromosome [23]. In addition the study suggested that this demasculinization may also be happening in microRNAs. Their analysis showed that there is about a 12-fold enrichment of evolutionarily young microRNAs in the X chromosome with respect to autosomes. For conserved microRNAs, the enrichment is less than twofold. However, when taking into account that multiple microRNAs may come from the same transcript (figure 1c), the figures are different: 3.5- and 1.8-fold enrichment for young and conserved microRNAs, respectively. These differences are small, and evidence for demasculinization in microRNAs is not supported.

There is an ongoing debate in the scientific literature about sex chromosome demasculinization. Although demasculinization has been generally considered one of the prominent features of *Drosophila* X chromosome evolution, recent work shows that the observed paucity of male-biased genes in the X chromosome may be artefactual [37–39]. Indeed, several groups suggest that demasculinization does not happen in *Drosophila* and propose that there is no global meiotic sex chromosome inactivation (MSCI) [40,41]. The movement of male-biased genes out of the X chromosome is often explained as a response to MSCI. This discussion is not settled, and evidence both for demasculinization and for MSCI is still reported [23,42–44]. Interestingly, most X-linked microRNAs escape MSCI [45]. These observations imply that X-chromosome demasculinization caused by MSCI might not happen during microRNA evolution. Even if there is an ongoing demasculinization process affecting protein-coding genes, microRNAs seem not to be affected.

Female microRNAs are generally older than male-biased microRNAs, and they are frequently encoded within other female-expressed genes. For instance, mir-995 is highly expressed in females (figure 1) and it is associated with oocytes

(figure 2). This microRNA is encoded within the first intron of *cdc2c*, a gene involved in cell proliferation during development [46]. Hence, the presence of mir-995 in oocytes may be a by-product of the host gene expression. In addition, mir-995 can be identified in the same intron of the orthologous *cdc2c* gene in other insects [47], showing a deep conservation of the microRNA/host gene association. Interestingly, mir-92a is encoded within a gene (*jigr1*) whose product is maternally deposited in the oocyte [48], and the microRNA is highly expressed in oocytes (figure 2) and detected in unfertilized eggs [33]. The presence of mir-92a in the developing egg may be a by-product of being intronic to a sex-biased expressed gene. As a matter of fact, mir-92a is associated with leg morphological differences between *Drosophila* species [49], a role (in principle) unrelated to any function in the early developing egg.

Among the microRNAs with a female-biased expression pattern, there are microRNAs associated with the gametic function. Recently, mir-989 has been discovered to be involved in cell migration in the ovary [50]. Indeed, the 3' arm of mir-989 is highly expressed in ovaries (figure 2). The analysis of female mutants also reveals that mir-9c (present in ovaries; figure 2) is somewhat involved in the control of the number of germ cells [51]. Predictably, other female-biased microRNAs here reported, such as mir-994/318, could have a role in gametic function. Strikingly, the mir-310/311/312/313, which is female-expressed (and probably maternally deposited in the egg), is involved in the development of male gonads [52]. This emphasizes that genes with sex-biased expression can also have other functions, even in the opposite sex.

We recently characterized sex-biased microRNAs in the parasitic *Schistosoma mansoni* and reported that one of the microRNA clusters (mir-71/mir-2) has two copies, one in the sexual chromosome with no detectable bias and another copy in an autosome with sex-biased expression. The duplication of the cluster happened more or less at the same time as sexual dimorphism appeared in this genus (*Schistosoma*). We suggested that this may be a case of escaping sex conflict, in which genes involved in sex dimorphism tend to be out of the X chromosome [10,53]. However, this is likely to be an exception to the rule in microRNAs, as their evolutionary dynamics is primarily dominated by high levels of emergence and a low probability of non-tandem duplication.

In summary, I conclude that sex-biased expression of microRNAs is a consequence of a high rate of microRNA de novo emergence. Novel microRNAs tend to appear in the X chromosome and to be expressed in the testes. Conversely, male-biased microRNAs are evolutionarily young and also show a high rate of loss. On the other hand, many

female-biased microRNA emerged within the intron of female-biased host genes. They are generally conserved suggesting that the female gametic function may be more constrained, and purifying selection could eliminate emerging microRNAs impairing ovary/oocyte development. This scenario suggests that positive/adaptive selection may have no more than little contribution to determining the sex-biased expression of microRNAs.

5. Methods

Drosophila melanogaster microRNA sequences are from miRBase version 19 [54]. Expression datasets were downloaded from Gene Expression Omnibus at <http://www.ncbi.nlm.nih.gov/geo/>, with accession numbers: GSM286602 and GSM399107 (adult males); GSM286603 and GSM399106 (adult females); GSM280082 (ovaries); GSM909277 and GSM909278 (testes); GSM385822 and GSM385744 (ovary somatic sheet); GSM180330 and GSM286613 (early embryos); GSM609223 and GSM609224 (young males and females) [31,55–59]. Reads from these experiments were mapped to *D. melanogaster* microRNA hairpins with BOWTIE v. 0.12.7 [60], allowing no mismatches nor multiple matches. Differential expression of microRNAs was estimated with EDGER [61]. In short, read counts were first normalized with the trimmed mean of M-values (TMM) method [62]. Then, the variation within samples was estimated by fitting the expression pattern to a negative binomial distribution. Sex-biased microRNAs were detected by an exact test controlling the false discovery rate [63]. Additionally, the analysis was repeated with a more general method implemented in DESEQ [64], but the results remained overall the same. Expression data for *Drosophila* genes were obtained

from modENCODE, available at www.flybase.org [58,65]. All statistical analyses and figures 1 and 2 were done with R [66].

Evolutionary age of microRNAs and microRNA families was estimated as previously described [67]. In brief, I compiled microRNA sequences with detectable similarity to *D. melanogaster* microRNAs with BLAST [68], using a sensitive set of parameters to detect homologous microRNAs ($-w 4$, $-q -3$, $-r +2$). I also included additional sequences described elsewhere to ensure that all known microRNA families with a common evolutionary origin are taken into account [54,67,69–71]. MicroRNA hairpins were aligned with CLUSTALX v. 2.0 [72], manually refining the alignments with RALEE [73], and phylogenetic trees were reconstructed using the neighbour-joining and maximum-likelihood routines with default parameter as implemented in MEGA 5 [74]. MicroRNA age estimates were also compared with those obtained by Mohammed *et al.* [75] using a different approach: they analysed whole genome alignments of 12 *Drosophila* genomes [76]. Age estimations were fully congruent between both datasets.

MicroRNA targets were retrieved from our previous study [77]. In short, 3'-UTRs were downloaded from FlyBase (genome version BDGP 5.25), and the microRNA targets were predicted with three different programs based on different algorithmic approaches: TargetScan [78], DIANA-microT [79] and miRanda [80], with default parameters. The evolutionary conservation of targeted genes was inferred from the gene family tree available at TreeFam 9 [81].

Acknowledgements. I am greatly indebted to Steve Dorus, Kirill Borziak and two anonymous reviewers for useful comments and critical reading of the manuscript. I also thank Maria Vibrationovski and Colin Meiklejohn for pointing me to relevant papers, and to Haldane's Sieve (<http://haldanessieve.org>) for promoting the dissemination and discussion of preprints (including the arXiv version of this work).

References

- Mank JE. 2009 Sex chromosomes and the evolution of sexual dimorphism: lessons from the genome. *Am. Nat.* **173**, 141–150. (doi:10.1086/595754)
- Parisi M, Nuttall R, Naiman D, Bouffard G, Malley J, Andrews J, Eastman S, Oliver B. 2003 Paucity of genes on the *Drosophila* X chromosome showing male-biased expression. *Science* **299**, 697–700. (doi:10.1126/science.1079190)
- Ranz JM, Castillo-Davis CI, Meiklejohn CD, Hartl DL. 2003 Sex-dependent gene expression and evolution of the *Drosophila* transcriptome. *Science* **300**, 1742–1745. (doi:10.1126/science.1085881)
- Kobayashi S *et al.* 2006 Comparison of gene expression in male and female mouse blastocysts revealed imprinting of the X-linked gene, *Rhox5/Pem*, at preimplantation stages. *Curr. Biol.* **16**, 166–172. (doi:10.1016/j.cub.2005.11.071)
- Arbeitman MN, Fleming AA, Siegal ML, Null BH, Baker BS. 2004 A genomic analysis of *Drosophila* somatic sexual differentiation and its regulation. *Development* **131**, 2007–2021. (doi:10.1242/dev.01077)
- Xue L, Noll M. 2002 Dual role of the Pax gene *paired* in accessory gland development of *Drosophila*. *Development* **129**, 339–346.
- Yang X, Schadt EE, Wang S, Wang H, Arnold AP, Ingram-Drake L, Drake TA, Lusis AJ. 2006 Tissue-specific expression and regulation of sexually dimorphic genes in mice. *Genome Res.* **16**, 995–1004. (doi:10.1101/gr.5217506)
- Krol J, Loedige I, Filipowicz W. 2010 The widespread regulation of microRNA biogenesis, function and decay. *Nat. Rev. Genet.* **11**, 597–610. (doi:10.1038/nrg2843)
- Axtell MJ, Westholm JO, Lai EC. 2011 Vive la différence: biogenesis and evolution of microRNAs in plants and animals. *Genome Biol.* **12**, 221. (doi:10.1186/gb-2011-12-4-221)
- Marco A, Kozomara A, Hui JHL, Emery AM, Rollinson D, Griffiths-Jones S, Ronschaugen M. 2013 Sex-biased expression of microRNAs in *Schistosoma mansoni*. *PLoS Negl. Trop. Dis.* **7**, e2402. (doi:10.1371/journal.pntd.0002402)
- Zhang X, Zheng Y, Jagadeeswaran G, Ren R, Sunkar R, Jiang H. 2012 Identification and developmental profiling of conserved and novel microRNAs in *Manduca sexta*. *Insect Biochem. Mol. Biol.* **42**, 381–395. (doi:10.1016/j.ibmb.2012.01.006)
- Mishima T *et al.* 2008 MicroRNA (miRNA) cloning analysis reveals sex differences in miRNA expression profiles between adult mouse testis and ovary. *Reproduction* **136**, 811–822. (doi:10.1530/REP-08-0349)
- Zhang Z, Hambuch TM, Parsch J. 2004 Molecular evolution of sex-biased genes in *Drosophila*. *Mol. Biol. Evol.* **21**, 2130–2139. (doi:10.1093/molbev/msh223)
- Pröschel M, Zhang Z, Parsch J. 2006 Widespread adaptive evolution of *Drosophila* genes with sex-biased expression. *Genetics* **174**, 893–900. (doi:10.1534/genetics.106.058008)
- Baines JF, Sawyer SA, Hartl DL, Parsch J. 2008 Effects of X-linkage and sex-biased gene expression on the rate of adaptive protein evolution in *Drosophila*. *Mol. Biol. Evol.* **25**, 1639–1650. (doi:10.1093/molbev/msn111)
- Nurminsky DI, Nurminskaya MV, Aguiar DD, Hartl DL. 1998 Selective sweep of a newly evolved sperm-specific gene in *Drosophila*. *Nature* **396**, 572–575. (doi:10.1038/25126)
- Levine MT, Jones CD, Kern AD, Lindfors HA, Begun DJ. 2006 Novel genes derived from noncoding DNA in *Drosophila melanogaster* are frequently X-linked and exhibit testis-biased expression. *Proc. Natl Acad. Sci. USA* **103**, 9935–9939. (doi:10.1073/pnas.0509809103)

18. Begun DJ, Lindfors HA, Kern AD, Jones CD. 2007 Evidence for *de novo* evolution of testis-expressed genes in the *Drosophila yakuba/Drosophila erecta* clade. *Genetics* **176**, 1131–1137. (doi:10.1534/genetics.106.069245)
19. Chen S-T, Cheng H-C, Barbash DA, Yang H-P. 2007 Evolution of *hydra*, a recently evolved testis-expressed gene with nine alternative first exons in *Drosophila melanogaster*. *PLoS Genet.* **3**, e107. (doi:10.1371/journal.pgen.0030107)
20. Sturgill D, Zhang Y, Parisi M, Oliver B. 2007 Demasculinization of X chromosomes in the *Drosophila* genus. *Nature* **450**, 238–241. (doi:10.1038/nature06330)
21. Zhou Q *et al.* 2008 On the origin of new genes in *Drosophila*. *Genome Res.* **18**, 1446–1455. (doi:10.1101/gr.076588.108)
22. Metta M, Schlotterer C. 2008 Male-biased genes are overrepresented among novel *Drosophila pseudoobscura* sex-biased genes. *BMC Evol. Biol.* **8**, 182. (doi:10.1186/1471-2148-8-182)
23. Zhang YE, Vibranovski MD, Krinsky BH, Long M. 2010 Age-dependent chromosomal distribution of male-biased genes in *Drosophila*. *Genome Res.* **20**, 1526–1533. (doi:10.1101/gr.107334.110)
24. Jiang Z-F, Machado CA. 2009 Evolution of sex-dependent gene expression in three recently diverged species of *Drosophila*. *Genetics* **183**, 1175–1185. (doi:10.1534/genetics.109.105775)
25. Campo-Paysaa F, Sémon M, Cameron RA, Peterson KJ, Schubert M. 2011 microRNA complements in deuterostomes: origin and evolution of microRNAs. *Evol. Dev.* **13**, 15–27. (doi:10.1111/j.1525-142X.2010.00452.x)
26. Marco A, Ninova M, Griffiths-Jones S. 2013 Multiple products from microRNA transcripts. *Biochem. Soc. Trans.* **41**, 850–854. (doi:10.1042/BST20130035)
27. Ruby JG, Stark A, Johnston WK, Kellis M, Bartel DP, Lai EC. 2007 Evolution, biogenesis, expression, and target predictions of a substantially expanded set of *Drosophila* microRNAs. *Genome Res.* **17**, 1850–1864. (doi:10.1101/gr.6597907)
28. Stark A, Kheradpour P, Parts L, Brennecke J, Hodges E, Hannon GJ, Kellis M. 2007 Systematic discovery and characterization of fly microRNAs using 12 *Drosophila* genomes. *Genome Res.* **17**, 1865–1879. (doi:10.1101/gr.6593807)
29. Berezikov E *et al.* 2011 Deep annotation of *Drosophila melanogaster* microRNAs yields insights into their processing, modification, and emergence. *Genome Res.* **21**, 203–215. (doi:10.1101/gr.116657.110)
30. Jin Z, Xie T. 2007 Dcr-1 maintains *Drosophila* ovarian stem cells. *Curr. Biol.* **17**, 539–544. (doi:10.1016/j.cub.2007.01.050)
31. Chung W-J, Okamura K, Martin R, Lai EC. 2008 Endogenous RNA interference provides a somatic defense against *Drosophila* transposons. *Curr. Biol.* **18**, 795–802. (doi:10.1016/j.cub.2008.05.006)
32. Lawrence PA. 1992 *The making of a fly: the genetics of animal design*. London, UK: Blackwell Scientific.
33. Marco A. In preparation. Maternal microRNAs in *Drosophila* unfertilized oocytes.
34. Zhang R, Peng Y, Wang W, Su B. 2007 Rapid evolution of an X-linked microRNA cluster in primates. *Genome Res.* **17**, 612–617. (doi:10.1101/gr.6146507)
35. Guo X, Su B, Zhou Z, Sha J. 2009 Rapid evolution of mammalian X-linked testis microRNAs. *BMC Genomics* **10**, 97. (doi:10.1186/1471-2164-10-97)
36. Li J, Liu Y, Dong D, Zhang Z. 2009 Evolution of an X-linked primate-specific microRNA cluster. *Mol. Biol. Evol.* **27**, 671–683. (doi:10.1093/molbev/msp284)
37. Meisel RP, Malone JH, Clark AG. 2012 Disentangling the relationship between sex-biased gene expression and X-linkage. *Genome Res.* **22**, 1255–1265. (doi:10.1101/gr.132100.111)
38. Meiklejohn CD, Presgraves DC. 2012 Little evidence for demasculinization of the *Drosophila* X chromosome among genes expressed in the male germline. *Genome Biol. Evol.* **4**, 1007–1016. (doi:10.1093/gbe/evs077)
39. Lindsley DL, Roote J, Kennison JA. 2013 Anent the genomics of spermatogenesis in *Drosophila melanogaster*. *PLoS ONE* **8**, e55915. (doi:10.1371/journal.pone.0055915)
40. Meiklejohn CD, Landeen EL, Cook JM, Kingan SB, Presgraves DC. 2011 Sex chromosome-specific regulation in the *Drosophila* male germline but little evidence for chromosomal dosage compensation or meiotic inactivation. *PLoS Biol.* **9**, e1001126. (doi:10.1371/journal.pbio.1001126)
41. Mikhaylova LM, Nurminsky DI. 2011 Lack of global meiotic sex chromosome inactivation, and paucity of tissue-specific gene expression on the *Drosophila* X chromosome. *BMC Biol.* **9**, 29. (doi:10.1186/1741-7007-9-29)
42. Vibranovski MD, Lopes HF, Karr TL, Long M. 2009 Stage-specific expression profiling of *Drosophila* spermatogenesis suggests that meiotic sex chromosome inactivation drives genomic relocation of testis-expressed genes. *PLoS Genet.* **5**, e1000731. (doi:10.1371/journal.pgen.1000731)
43. Vibranovski MD, Zhang YE, Kemkemer C, Lopes HF, Karr TL, Long M. 2012 Re-analysis of the larval testis data on meiotic sex chromosome inactivation revealed evidence for tissue-specific gene expression related to the *Drosophila* X chromosome. *BMC Biol.* **10**, 49. (doi:10.1186/1741-7007-10-49)
44. Gao G *et al.* In press. A long term demasculinization of X-linked intergenic noncoding RNAs in *Drosophila melanogaster*. *Genome Res.* (doi:10.1101/gr.165837.113)
45. Song R, Ro S, Michaels JD, Park C, McCarrey JR, Yan W. 2009 Many X-linked microRNAs escape meiotic sex chromosome inactivation. *Nat. Genet.* **41**, 488–493. (doi:10.1038/ng.338)
46. Lehner CF, O'Farrell PH. 1990 *Drosophila* cdc2 homologs: a functional homolog is coexpressed with a cognate variant. *EMBO J.* **9**, 3573–3581.
47. Marco A, Hui JHL, Ronshaugen M, Griffiths-Jones S. 2010 Functional shifts in insect microRNA evolution. *Genome Biol. Evol.* **2**, 686–696. (doi:10.1093/gbe/evq053)
48. Fisher B *et al.* 2012 *BDGP insitu homepage*. See <http://insitu.fruitfly.org/cgi-bin/ex/insitu.pl>.
49. Arif S *et al.* 2013 Evolution of mir-92a underlies natural morphological variation in *Drosophila melanogaster*. *Curr. Biol.* **23**, 523–528. (doi:10.1016/j.cub.2013.02.018)
50. Kugler J-M, Chen Y-W, Weng R, Cohen SM. 2013 miR-989 Is required for border cell migration in the *Drosophila* ovary. *PLoS ONE* **8**, e67075. (doi:10.1371/journal.pone.0067075)
51. Kugler J-M, Chen Y-W, Weng R, Cohen SM. 2013 Maternal loss of miRNAs leads to increased variance in primordial germ cell numbers in *Drosophila melanogaster*. *G3* **3**, 1573–1576. (doi:10.1534/g3.113.007591)
52. Pancratov R *et al.* 2013 The miR-310/13 cluster antagonizes β -catenin function in the regulation of germ and somatic cell differentiation in the *Drosophila* testis. *Development* **140**, 2904–2916. (doi:10.1242/dev.092817)
53. Marco A, Hooks K, Griffiths-Jones S. 2012 Evolution and function of the extended miR-2 microRNA family. *RNA Biol.* **9**, 242–248. (doi:10.4161/ma.19160)
54. Kozomara A, Griffiths-Jones S. 2011 miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Res.* **39**, D152–D157. (doi:10.1093/nar/gkq1027)
55. Ruby JG, Jan C, Player C, Axtell MJ, Lee W, Nusbaum C, Ge H, Bartel DP. 2006 Large-scale sequencing reveals 21U-RNAs and additional microRNAs and endogenous siRNAs in *C. elegans*. *Cell* **127**, 1193–1207. (doi:10.1016/j.cell.2006.10.040)
56. Czech B *et al.* 2008 An endogenous small interfering RNA pathway in *Drosophila*. *Nature* **453**, 798–802. (doi:10.1038/nature07007)
57. Lau N, Robine N, Martin R, Chung W-J, Niki Y, Berezikov E, Lai E. 2009 Abundant primary piRNAs, endo-siRNAs, and microRNAs in a *Drosophila* ovary cell line. *Genome Res.* **19**, 1776–1785. (doi:10.1101/gr.094896.109)
58. Roy S *et al.* 2010 Identification of functional elements and regulatory circuits by *Drosophila* modENCODE. *Science* **330**, 1787–1797. (doi:10.1126/science.1198374)
59. Toledano H, D'Alterio C, Czech B, Levine E, Jones DL. 2012 The let-7–Imp axis regulates ageing of the *Drosophila* testis stem-cell niche. *Nature* **485**, 605–610. (doi:10.1038/nature11061)
60. Langmead B, Trapnell C, Pop M, Salzberg SL. 2009 Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25. (doi:10.1186/gb-2009-10-3-r25)
61. Robinson MD, McCarthy DJ, Smyth GK. 2010 edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140. (doi:10.1093/bioinformatics/btp616)
62. Robinson MD, Oshlack A. 2010 A scaling normalization method for differential expression

- analysis of RNA-seq data. *Genome Biol.* **11**, R25. (doi:10.1186/gb-2010-11-3-r25)
63. Benjamini Y, Hochberg Y. 1995 Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. B Methodol.* **57**, 289–300. (doi:10.2307/2346101)
 64. Anders S, Huber W. 2010 Differential expression analysis for sequence count data. *Genome Biol.* **11**, R106. (doi:10.1186/gb-2010-11-10-r106)
 65. McQuilton P, St Pierre SE, Thurmond J. 2012 FlyBase 101: the basics of navigating FlyBase. *Nucleic Acids Res.* **40**, D706–D714. (doi:10.1093/nar/gkr1030)
 66. R Development Core Team. 2004 *R: A language and environment for statistical computing*. Vienna, Austria: R Development Core Team.
 67. Marco A, Ninova M, Ronshaugen M, Griffiths-Jones S. 2013 Clusters of microRNAs emerge by new hairpins in existing transcripts. *Nucleic Acids Res.* **41**, 7745–7752. (doi:10.1093/nar/gkt534)
 68. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997 Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389–3402. (doi:10.1093/nar/25.17.3389)
 69. Sempere LF, Cole CN, McPeck MA, Peterson KJ. 2006 The phylogenetic distribution of metazoan microRNAs: insights into evolutionary complexity and constraint. *J. Exp. Zool. B Mol. Dev. Evol.* **306**, 575–588. (doi:10.1002/jez.b.21118)
 70. Wheeler BM, Heimberg AM, Moy VN, Sperling EA, Holstein TW, Heber S, Peterson KJ. 2009 The deep evolution of metazoan microRNAs. *Evol. Dev.* **11**, 50–68. (doi:10.1111/j.1525-142X.2008.00302.x)
 71. Nozawa M, Miura S, Nei M. 2010 Origins and evolution of microRNA genes in *Drosophila* species. *Genome Biol. Evol.* **2**, 180–189. (doi:10.1093/gbe/evq009)
 72. Larkin MA *et al.* 2007 CLUSTALW and CLUSTALX version 2.0. *Bioinformatics* **23**, 2947–2948. (doi:10.1093/bioinformatics/btm404)
 73. Griffiths-Jones S. 2005 RALEE: RNA alignment editor in Emacs. *Bioinformatics* **21**, 257–259. (doi:10.1093/bioinformatics/bth489)
 74. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011 MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731–2739. (doi:10.1093/molbev/msr121)
 75. Mohammed J, Flynt AS, Siepel A, Lai EC. 2013 The impact of age, biogenesis, and genomic clustering on *Drosophila* microRNA evolution. *RNA* **19**, 1295–1308. (doi:10.1261/ma.039248.113)
 76. Clark AG *et al.* 2007 Evolution of genes and genomes on the *Drosophila* phylogeny. *Nature* **450**, 203–218. (doi:10.1038/nature06341)
 77. Marco A, MacPherson JI, Ronshaugen M, Griffiths-Jones S. 2012 MicroRNAs from the same precursor have different targeting properties. *Silence* **3**, 8. (doi:10.1186/1758-907X-3-8)
 78. Lewis BP, Burge CB, Bartel DP. 2005 Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* **120**, 15–20. (doi:10.1016/j.cell.2004.12.035)
 79. Maragkakis M *et al.* 2009 Accurate microRNA target prediction correlates with protein repression levels. *BMC Bioinformatics* **10**, 295. (doi:10.1186/1471-2105-10-295)
 80. Enright A, John B, Gaul U, Tuschl T, Sander C, Marks D. 2003 MicroRNA targets in *Drosophila*. *Genome Biol.* **5**, R1. (doi:10.1186/gb-2003-5-1-r1)
 81. Ruan J *et al.* 2007 TreeFam: 2008 update. *Nucleic Acids Res.* **36**, D735–D740. (doi:10.1093/nar/gkm1005)