

RESEARCH ARTICLE

Uniparental and transgressive expression of α -zeins in maize endosperm of o2 hybrid lines

Silvana Castelli¹, Iride Mascheretti¹, Cristian Cosentino, Barbara Lazzari, Raul Pirona, Aldo Ceriotti¹, Angelo Viotti*, Massimiliano Lauria¹*

Istituto di Biologia e Biotecnologia Agraria, CNR, Via Alfonso Corti, Milano, Italy

¹ These authors contributed equally to this work.

* viotti@ibba.cnr.it(AV); lauria@ibba.cnr.it(ML)



Abstract

The α -zein gene family encodes the most abundant storage proteins of maize (*Zea mays*) endosperm. Members of this family are expressed in a parent-of-origin manner. To characterize this phenomenon further, we investigated the expression of a subset of α -zein polypeptides in reciprocal crosses between o2 lines that were characterized by a simplified α -zein pattern. Maize lines that suppressed the expression of α -zeins when used as female parents were identified. The suppression was cross-specific, occurring only when specific genetic backgrounds were combined. Four α -zein sequences that were sensitive to uniparental expression were isolated. Molecular characterization of these α -zeins confirmed that their expression or suppression depended on the genetic properties of the endosperm tissue instead of their parental origin. DNA methylation analysis of both maternally and paternally expressed α -zeins revealed no clear correlation between this epigenetic marker and parent-of-origin allelic expression, suggesting that an additional factor(s) is involved in this process. Genetic analyses revealed that the ability of certain lines to suppress α -zein expression was unstable after one round of heterozygosity with non-suppressing lines. Interestingly, α -zeins also showed a transgressive expression pattern because unexpressed isoforms were reactivated in both F2 and backcross plants. Collectively, our results suggest that parent-of-origin expression of specific α -zein alleles depends on a complex interaction between genotypes in a manner that is reminiscent of paramutation-like phenomena.

OPEN ACCESS

Citation: Castelli S, Mascheretti I, Cosentino C, Lazzari B, Pirona R, Ceriotti A, et al. (2018) Uniparental and transgressive expression of α -zeins in maize endosperm of o2 hybrid lines. PLoS ONE 13(11): e0206993. <https://doi.org/10.1371/journal.pone.0206993>

Editor: Craig Eliot Coleman, Brigham Young University, UNITED STATES

Received: May 11, 2018

Accepted: October 23, 2018

Published: November 15, 2018

Copyright: © 2018 Castelli et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: Research was supported by a grant from MIUR, FIRB project no. RBNE01TYZF, and a special grant from MIUR and CNR for the Epigenomics Flagship Project (EPIGEN). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Introduction

Parental dependency of genetic traits frequently results from unequal contributions of genetic information from male and female gametes. Examples include traits that are encoded by mitochondria and chloroplast genes, Y chromosome-linked genes, transposable elements, and biased chromosome transmission [1,2]. Alternatively, parent-of-origin effects on gene expression can result in specific gene silencing of one of the parent alleles: a phenomenon that is known as genomic imprinting [3]. In plants, genomic imprinting was originally discovered in maize, in which certain alleles of the *r1* locus, which encodes a transcription factor that is

Competing interests: The authors have declared that no competing interests exist.

involved in seed pigmentation, are preferentially expressed upon maternal transmission [4]. Since then, several other examples of imprinted genes have been reported in maize, including specific alleles of the zein-regulator locus *dzr1* [5] and of the α -tubulins [6]. Although the functional relevance of the imprinted expression of these maize genes remains unknown, subsequent studies have demonstrated that genomic imprinting plays a key role in the normal development of the endosperm, the specialized seed tissue that supports the embryo during germination [7]. Indeed, a number of genes cause abnormal seed development in a parent-of-origin manner when they become mutated [8,9,10,11].

High-throughput genomic and post-genomic approaches, particularly RNA-Seq, have provided a more precise genome-wide survey of imprinted genes in hybrid endosperms of different plant species [12,13,14,15], including maize [16,17,18,19]. Collectively, these investigations have enabled the identification of a discrete number of genes that exhibit imprinted expression profiles. Interestingly, only a small subset of imprinted genes are conserved among plant species, and even within the same species there are examples of allelic variations in imprinting [17,20,21]. A possible explanation for this finding is that a plant's endosperm, which is the tissue where genomic imprinting principally occurs in plants, is characterized by an epigenetic asymmetry of parental genomes: the maternal genome is hypomethylated compared to its paternal counterpart [15,17,22,23,24,25]. It has been proposed that genome-wide demethylation of the endosperm is the result of the female epigenetic reprogramming process that occurs in the central cell, the progenitor of the endosperm, and that preferentially targets transposable elements. Demethylation of these elements results in the production of small interfering RNAs that in turn are suspected to reinforce transposon silencing in the adjacent egg cell, which contributes genetic material to the next generation [22,25]. This observation has led to the idea that genes near transposable elements might become indirect targets of transposon reprogramming, resulting in imprinted expression. Imprinted genes that are conserved among species were positively selected because of their adaptive advantage to endosperm development. Differently, imprinted genes that are confined to a specific species or ecotype/variety likely reflect allelic variations of transposon-associated genes that might either lack functional relevance or lead to the production of novel seed phenotypes that are species-specific [17,20,21].

Parent-of-origin effects on gene expression have also been reported for some alleles of the α -zeins [26]. These genes encode the most abundant fraction of the storage protein component of the maize endosperm and form a large gene family that can be subdivided into 4 subfamilies: *z1A*, *z1B*, *z1C*, *z1D* [27].

The *z1A*, *z1B* and *z1D* genes encode the light-type proteins (zLs) with an apparent mass of 19-kDa, whereas the *z1C* genes encode the heavy-type proteins (zHs) with an apparent mass of 22 kDa.

Although α -zeins can be easily characterized by SDS-PAGE, the study of parent-of-origin effects at these loci is complicated. For example, under optimal SDS-PAGE conditions, the 22- and 19-kDa proteins can be further divided, into five size subclasses: zein heavy-type 1 and 2 (zH1 and zH2) and zein light type 1, 2 and 3 (zL1, zL2 and zL3) [28], and each subclass is composed of several polypeptides, as determined by 2D-gel analysis (IEF followed by SDS-PAGE) [28].

A previous study [29] showed that different inbred lines carrying mutant *opaque-2* (*o2*) alleles, which is the major transcription factor of 22-kDa zeins [30], have a simplified zH pattern. We took advantage of these lines to perform a more accurate investigation of the uniparental expression of these genes in maize endosperm. Our results showed that uniparental expression of α -zeins depends on a specific combination of parental genetic backgrounds. DNA methylation, an epigenetic mark that is often associated with imprinting expression [3], did not show an obvious correlation with expression, suggesting that additional factors are

involved in the parent-of-origin effects that occur at these loci. Genetic analyses revealed that the ability of certain lines to suppress zHs expression is unstable after one round of heterozygosity with non-suppressing lines. The overall results support the idea that parent-of-origin expression of specific α -zein alleles depend on a complex interaction between genotypes in a manner that is reminiscent of paramutation-like phenomena.

Materials and methods

Plant material

Maize *o2* mutant lines (Table 1) were characterized and described in Ciceri et al. [29]. Plants were grown either in the greenhouse or in the field, and seeds were harvested at maturity or at 20 days after pollination (DAP). Immature seeds were frozen in liquid nitrogen and stored at -80°C.

Zein extraction, electrophoresis analysis and peptide sequence determination

Zeins were analyzed by SDS-PAGE and 2D-gel analysis (2D, IEF followed by SDS-PAGE) as reported in Lund et al. [26]. Specifically, 100 mg of flour was mixed with 400 μ l of 70% ethanol and 2% of 2-mercaptoethanol, and the mixture was kept in agitation at room temperature overnight. Alternatively, zein extraction was conducted without 2-mercaptoethanol. By

Table 1. zH1 pattern of inbred lines and reciprocal crosses as determined by SDS-PAGE analysis.

	Genotype	zH1 pattern
Inbred lines	NYRo2It (No2It)	+
	W22o2It	+
	Bianchio2It	+
	33o2It	+
	Rossmanno2R (Ro2R)	-
	W23o2R	-
	Mo17o2R (Mo2R)	-
Reciprocal crosses	No2It x Ro2R (NR)	+
	Ro2R x No2It (RN)	-
	No2It x W23o2R	+
	W23o2R x No2It	-
	No2It x Mo2R (NM)	+
	Mo2R x No2It (MN)	+
	W22o2It x Ro2R	+
	Ro2R x W22o2It	-
	W22o2It x W23o2R	+
	W23o2R x W22o2It	-
	Bianchio2It x Ro2R	+
	Ro2R x Bianchio2It	+
	33o2It x Ro2R	+
	Ro2R x 33o2It	+
	Bianchio2It x Mo2R	+
	Mo2R x Bianchio2It	+
	Bianchio2It x W23o2R	+
W23o2R x Bianchio2It	+	

<https://doi.org/10.1371/journal.pone.0206993.t001>

omitting this reagent, at room temperature, the protein extract was mainly composed by the α -zeins, whereas the other zein classes were less represented in the protein extract. This approach reduced the presence of 27-kDa γ zein, thus favoring detection α -zeins, especially in SDS-PAGE experiments run on a short electrophoresis apparatus. To obtain large amounts of a single polypeptide for sequence determination, pieces of gel containing each spot were extracted from a 2D gel and rerun via SDS-PAGE. Proteins were transferred from the SDS-PAGE gel to the Immobilon PVDF transfer membrane (Millipore) in transfer buffer that was composed of 20% methanol, 25 mM Tris, and 192 mM glycine at pH 8.3. Sequences at the amino-terminus were determined by automatic sequencing using an ABI-Perkin Elmer Mod. 477A (Primm).

Nucleic acid extraction and amplification

Total DNA and RNA from 20-DAP endosperms of various genotypes were extracted by using CTAB and Trizol (ThermoFisher), respectively. RNA samples (10 μ g) were treated with one unit of DNase I RNase-free (Ambion) to remove DNA contamination according to the manufacturer's instructions. Subsequently, 2 μ g of the treated RNA was used for first-strand cDNA synthesis with SuperScript II (Invitrogen) and 500 ng of oligo(dT)18 primer at 42°C for 1 h in a final volume of 25 μ L. PCR was performed by using ~40 ng of genomic DNA and 5 μ L of the reverse transcript (RT). For quantitative reverse transcription PCR (qRT-PCR), the cDNA reaction was diluted six-fold, and 3 μ L of the diluted solution was used for the analysis using the iTaq Universal SYBR Green Super mix (Bio-Rad). The comparative CT method ($\Delta\Delta$ CT method) was employed for the relative quantification of gene expression, and the *GRMZM2G023418* gene was used as the endogenous reference. This gene encodes for eukaryotic translation initiation factor that is stably expressed across different maize tissues [31]. To confirm primer specificity, PCR products corresponding to each of the studied α -zeins were cloned into pGEM-T Easy vectors (Promega) and sequenced. Primers and PCR conditions are reported in S1 Table.

Construction of *z1C* zein mini library and characterization of clones

cDNAs obtained from *No2It* and *No2It* x *Ro2R* (NR) endosperms were used to amplify *z1C* sequences using *z1CF* and *z1CR* primers (see S1 Table) that contained *Hind*III restriction sites. The amplification products were analyzed on a 1% agarose gel and then purified using a Wizard PCR Preps kit (Promega). After a reaction with T4 DNA polymerase in the presence of 100 μ M of each of the dNTPs, the fragments were cleaved with *Hind*III, treated at 65°C for 15 min, and inserted by ligation into a modified pBluescriptKS (-) vector (pBSL2A3) that was pre-digested with *Sma*I and *Hind*III. Plasmids were transformed into the XL1B *E. coli* strain (Stratagene) by electroporation. Single transformed colonies were grown overnight in LB medium, and plasmid DNA was extracted using a Wizard Minipreps DNA purification system Kit (Promega). *z1C* recombinant plasmids were characterized by digestion with the restriction enzymes *Pvu*II, *Hinc*II, *Pst*I and *Hha*I, all of which are diagnostic for their presence or absence in the coding sequences of α -zeins. The digested DNA was subjected to gel electrophoresis. Clones of interest were sequenced on both strands to confirm *z1C* identity.

Methylation assays

For each sample, 500 ng of DNA was digested overnight with either a methylation-sensitive or the methylation-dependent *Mcr*BC restriction enzyme in 50 μ L of reaction mix. For the *Msp*JI methylation-dependent restriction enzyme, the digestion was incubated for 3 h in 50 μ L of reaction mix. Semi-quantitative PCR (PlatinumTaq DNA Polymerase; Invitrogen) was performed with 3 μ L of digested DNA (~30 ng) in a 30 μ L reaction using specific primers (S1

Table). The EpiTect Bisulfite Kit (Qiagen) was used for single cytosine methylation analysis according to the manufacturer's instructions. PCR products obtained with degenerated primers were cloned into pGEM-T Easy vectors (Promega) and sequenced. Data were analyzed with the online free software Kismeth (<http://katahdin.mssm.edu/kismeth>; [32]). Primers and PCR conditions are reported in S1 Table.

Results

Analysis of α -zein expression in reciprocal crosses of different *o2* lines

To simplify the analysis of the parent-of-origin effects of α -zeins, we used a set of previously characterized *o2* mutant lines that exhibit simplified expression patterns of the heavy-type zeins (zHs), which can further subdivided in zH1 and zH2 types (Fig 1A). The *Bianchio2It* (*Bo2It*), *NYRo2It* (*No2It*), *W22o2It*, and *3316o2It* inbred lines carry the *o2Italian* (*o2It*) allele (Table 1, Fig 1B and 1C) and retain the ability to express some members of zH1s. Conversely, the *Rossmanno2R* (*Ro2R*), *W23o2R*, and *Mo17o2R* (*Mo2R*) inbred lines carry different null-transcript *o2R* alleles and exhibit a more severe reduction of zHs (Table 1, Fig 1B and 1C) [29,33]. Accordingly, nine pairs of reciprocal crosses between the *o2It* and *o2R* lines were generated, and the expression of zH1s in hybrid endosperms was determined by SDS-PAGE. Because zein pattern accumulation is genotype specific, to facilitate the analysis, zeins were extracted mainly by omitting the 2-mercaptoethanol reducing agent. By using this approach, we improved the detection, comparison and quantification of α -zeins by reducing the presence of the other zein family members (see Materials and Methods and Fig 1B). The analysis allowed us to identify at least two pairs of reciprocal crosses in which the expression of zH1s occurred in a parent-of-origin manner (Table 1). For example, in reciprocal crosses between the *No2It* and *Ro2R* lines (NR and RN), the accumulation of zH1s preferentially occurred when the *No2It* line was used as the female parent in the cross (Fig 1B). The uniparental expression is, however, genetic background specific, as it only occurs when specific genetic backgrounds are combined together. For example, in reciprocal crosses between the *No2It* and *Mo2R* lines, a normal zH1s accumulation was observed in both the *No2It* x *Mo2R* (NM) and *Mo2R* x *No2It* (MN) crosses (Table 1 and Fig 1C). In this study, to characterize this phenomenon further, we focused on the *No2It* and *Ro2R* lines, as these materials have been extensively analyzed over the past 20 years.

The zH1 pattern displayed by the *No2It* and *No2It* x *Ro2R* (NR) endosperms was consistently weaker than those produced from other zein types, e.g., the light-type zeins (19-kDa). Because the endosperm is a triploid tissue that includes two maternal genomes and one paternal genome, it was important to confirm that the absence of α -zeins in the *Ro2R* x *No2It* (RN) endosperms was not the result of a dosage effect. For instance, the staining procedure had low sensitivity in detecting zH1s produced from one copy of the *No2It* genome in RN endosperms. Therefore, the transcript levels of α -zeins were determined by qRT-PCR by using RNA extracted from immature endosperms (20 DAP) with specific primers for each of the four α -zein subfamilies, *z1A*, *z1B*, *z1C* and *z1D* [34]. Although each subfamily produced a unique transcriptional profile among *No2It*, *Ro2R* and their reciprocal hybrids, only the *z1C* subfamily produced a clear parent-of-origin expression pattern (Fig 1D). In contrast, expression analysis of *z1C* in NM and MN crosses produced an additive transcriptional pattern, which was in line with the accumulation of zH1s in mature seeds. Of note, we observed a general reduction of transcripts encoding the 19-kDa α -zeins in the *Ro2R* compared to the *No2It*. Collectively, our results suggest that i) the parent-of-origin pattern of α -zeins observed in NR and RN mature endosperms reflected the reduced amounts of their transcripts observed in the RN immature endosperms and that ii) the phenomenon is genetic background specific.




Fig 1. Expression analysis of α -zein in *o2* hybrid endosperms. (A) Representative example of zein pattern of a wild type endosperm (NYR) as determined by SDS-PAGE analysis. Black arrows indicate the different members of the zein family: 27-kDa, 22-kDa, 19-kDa, 16-kDa, 15-kDa, 10-kDa. (B) SDS-PAGE analysis of zein extracts of *No2It* (zH1-plus), *Ro2R* (zH1-null), *No2It* x *Ro2R* (NR) and *Ro2R* x *No2It* (RN) mature endosperms. The first genotype in the cross is the seed parent. Representative examples of zein extraction performed with (+) or without (-) 2-mercaptoethanol are shown. As discussed in the Results section, this experimental strategy permitted us to simplify the analysis of α -zeins. (C) Example of additive accumulation pattern of α -zeins between the *No2It* and *Mo2R* lines. To confirm that detection of zH1s did not depend on protein loading issues, different amount of *No2It* x *Mo2R* (NM) and *Mo2R* x *No2It* (MN) extracts were loaded. (D) Quantitative RT-PCR analysis of the four α -zein subfamilies (*z1A*, *z1B*, *z1C* and *z1D*) in 20-DAP endosperm. Gene expression was normalized to *GRMZM2G023418*. Data are representative of three biological replicates. Error bars represent the SD of three technical replicates.

<https://doi.org/10.1371/journal.pone.0206993.g001>

Transcriptional analysis of α -zein regulators

The *o2It* allele encodes a protein that localizes to the nucleus and binds to the *O2* target sequence, although with a reduced efficiency compared to the wild-type protein [35]. To exclude the possibility that uniparental expression of α -zeins was due to imprinting regulation of their transcriptional activator, the *o2It* transcript level was analyzed by qRT-PCR in the same cDNA samples used to characterize the α -zein expression. In addition, the transcript levels of *OHP1/2* (*opaque-2 heterodimerizing protein 1 and 2*), *PBF* (*prolamin-box binding factor*), *GCN5* (*general control of amino-acid synthesis protein 5*), *ADA2* (*transcriptional adaptor 2*), *ZmMADS47*, and *ZmTaxilin*, *O11* (*opaque11*), *FL3* (*floury3*), *NKD1/2* (*naked endosperm 1 and 2*) were also analyzed, as these genes encode proteins that participate on α -zein regulation [7,36,37,38,39,40,41,42,43]. The results presented in Fig 2 show that none of the loci analyzed produced an expression pattern that could explain uniparental expression of α -zeins in NR and RN hybrids. It is important to note, however, that the rationale of this analysis was to detect gross transgressive transcriptional patterns between hybrids that could explain uniparental expression of α -zeins. Indeed, our analysis did not clarify the possible presence of regulator allelic variants between *No2It* and *Ro2R*, which may produce proteins with different properties, for example, weak or nonfunctional alleles that are expressed in a parent-of-origin manner.

Identification of α -zein alleles that are sensitive to parent-of-origin effects

To identify α -zeins that are subjected to parent-of-origin effects, we analyzed their expression in NR and RN endosperms using 2D-gel analysis. Based on the 2D profiles, three *No2It*-specific polypeptides were differentially detectable between the two reciprocals (Fig 3; black arrows), including one heavy-type (zH1b) and two light-type (zL1a and zL2b, belonging to the group of the 19kD polypeptides). An additional *No2It*-specific isoform (zH1a) was also differentially detectable between NR and RN endosperms. However, because of its low intensity, it remained unclear if zH1a reflected a true differentially expressed α -zein. Interestingly, 2D-gel analysis also revealed the presence of two additional isoforms, zH1c and zH1d, that were preferentially expressed in the NR endosperm (Fig 3; gray arrows). The data presented below will show that these latter isoforms belonged to the *Ro2R* genotype, and they may represent those very faint zH1 spots present in the 2D-gel of *Ro2R*. Although zH1d was detectable in both the reciprocals, its intensity was lower in RN, even if two maternal copies of this allele were present in the endosperm. Consequently, the expression of zH1c and zH1d in NR endosperm was favored upon paternal transmission. By determining the amino-terminal sequences of zH1b, zH1d and zL1a, we found that they all matched the consensus sequence of the *z1C* subfamily (S1 Fig) [28,44]; the fact that zL1a is the product of a zH gene type is not surprising, as zH sequences with internal deletions that modify their coding capacities have been described [28,45]. To clone the full-length coding sequences of α -zeins subjected to parent-of-origin expression, we prepared cDNA mini-libraries of *z1C* sequences from RNAs that were extracted




Fig 2. Transcriptional analysis of α -zein regulator genes. Quantitative RT-PCR analysis of α -zein regulator genes in 20-DAP endosperm. Gene expression was normalized to *GRMZM2G023418*. The first genotype in the cross is the seed parent. Data are representative of three biological replicates. Error bars represent the SD of three technical replicates.

<https://doi.org/10.1371/journal.pone.0206993.g002>




Fig 3. Identification of α -zein isoforms that are subject to parent-of-origin effects by 2D-gel analysis. Approximately 100 μ g of zein protein extract was loaded for each genotype, with the exception of Ro2R, for which 150 μ g of zein extract was used. The numbers above the No2It panel indicate the pH gradient that was used for the first dimension. The parentheses group the zH and zL isoforms. The black arrows indicate No2It-specific α -zeins that were differentially detected in the reciprocal NR and RN. Gray arrows indicate Ro2R-specific isoforms that were preferentially detected in NR endosperms. zL2c is an example of a No2-specific isoform that accumulated in an additive manner.

<https://doi.org/10.1371/journal.pone.0206993.g003>

from No2It and NR endosperms. We identified 4 clones that, based on their properties (length, nucleotide sequence, amino acid sequences, and *in silico*-predicted pI), were assigned to genes that encoded zL1a, zH1b, zH1c and zH1d, whose nucleotide sequences closely resembled the *azs22/D87*, *azs22/6*, *azs22/10* and *azs22/16* alleles [17,44,46], respectively (S2 Fig).

Accordingly, primers that were specific to the four sequences were designed by using the sequence information of α -zeins that were characterized in the B73 and BSS53 lines [44,46]. PCR analysis performed under stringent conditions confirmed that *zH1b* and *zL1a* belonged to the No2It genotype, whereas *zH1c* and *zH1b* belonged to the Ro2R genotype (Fig 4A, PCR panel). Following this, RT-PCR analysis was performed using endosperm RNAs that were extracted from the No2It and Ro2R lines and their reciprocals. As a control, a similar analysis was performed by using RNAs extracted from Mo2R, NM and MN endosperms. Overall, the results were in good agreement with data above. For instance, strongly maternally biased transcription was observed for *zH1b* and *zL1a*, whereas paternally biased transcription, though not exclusively, was observed for *zH1c* and *zH1b* (Fig 4A, RT-PCR panel). Finally, RT-PCR analysis confirmed that both *zL1a* and *zH1b* (No2It specific) were expressed in NM and MN endosperms (Fig 4B, RT-PCR panel), supporting both the no-suppressing characteristics of the Mo2R line on these No2It alleles and the genetic background-specific nature of the phenomenon under study.

Inheritance of α -zein uniparental expression in backcrosses and F2 progenies

Expression of 10- and 18-kDa δ zeins can occur in a parent-of-origin manner due to parental imprinting of the regulatory *dzt1* locus that controls their accumulation [5,47]. Therefore, we aimed to determine whether a similar mechanism might explain the uniparental expression patterns of α -zeins in hybrid seeds. For clarification purposes, SDS-PAGE analysis was used to analyze F2 seeds generated from NR and RN F1 plants and seeds generated from reciprocal backcrosses between the NR and RN F1 and Ro2R plants (Fig 5 and Table 2). If the Ro2R and No2It lines possessed a monogenic maternally expressed factor that negatively or positively regulated the accumulation of α -zeins, respectively, then the presence of zH1s should have occurred at a frequency of 50% in RN x Ro2R, NR x Ro2R, NR- and RN F2 seeds. In these crosses, in contrast to our expectations, we found a mixed situation. In some crosses, the expression of zH1s occurred at high frequency that ranged from 90% to 100%, whereas in other cases the expression dropped to 43% (Fig 5A and Table 2). On the other hand, Ro2R plants that were pollinated with either RN or NR plants produced zH-null seeds with frequencies that ranged from 88% to 100% (Fig 5A and Table 2), mimicking the expression pattern observed in the RN hybrid endosperm. These results suggest that α -zein suppression mediated by Ro2R can be unstable when this genotype is maternally inherited from either an RN or an NR hybrid plant. Interestingly, in some backcrosses and F2 seeds, we also observed reactivation of the zH2 zein type (Fig 5C).

3.4 DNA methylation analysis of α -zein alleles

Tissue-specific expression of α -zeins is epigenetically regulated [48,49,50,51,52]. Moreover, α -zeins show parental methylation differences within the endosperm [6]. Therefore, we aimed to




Fig 4. Allele specific analysis of α -zeins. (A) PCR and RT-PCR analysis of DNA and cDNA, respectively, of *No2It*, *NR*, *RN*, and *Ro2R* with primers specific to *zL1a*, *zH1b*, *zH1c*, *zH1d* alleles. (B) PCR and RT-PCR analysis of DNA and cDNA, respectively, of *No2It*, *NM*, *MN*, and *Mo2R* with primers specific to *zL1a*, *zH1b* alleles. For PCR analysis two independent genomic extractions were analyzed. Gene expression of *GRMZM2G023418* (*HK2*) was used as endogenous reference. RT-PCR (-RT) indicates that the reverse transcriptase enzyme was omitted in the cDNA reaction. Data are representative of two biological replicates. M1 and M2 indicate different molecular markers.

<https://doi.org/10.1371/journal.pone.0206993.g004>




Fig 5. Representative examples of SDS-PAGE of α -zein extracts from backcrosses and F2 endosperm progeny. (A) Differential accumulation of the zH1 type in the reciprocal backcrosses Ro2R x RN and RN x Ro2R. #1 and #2 indicate that these samples come from different ears. (B) Accumulation of zH1 type in RN F2 endosperms. (C) Reactivation of zH2 type in RN F2 endosperms. The Ro2R x RN zein extract was used as negative controls. Each lane represents a single seed zein extraction.

<https://doi.org/10.1371/journal.pone.0206993.g005>

determine whether allelic differences in DNA methylation might provide a possible explanation for the parent-of-origin effects that were observed at the studied loci. Accordingly, methylation-sensitive and methylation-dependent PCR analyses were performed using allele-specific primers to explore parental methylation differences of the *zL1a*, *zH1b*, *zH1c* and *zH1d* alleles in hybrid endosperms. Collectively, methylation-dependent PCR analysis confirmed that all of the α -zein alleles were subjected to maternal demethylation (Fig 6B). Conversely, methylation-sensitive PCR analysis, which detects methylation differences at specific cytosines, revealed that the extent of DNA demethylation was slightly different between alleles and hybrid genotypes (Fig 6B). The results also revealed a lack of a clear correlation between DNA

Table 2. zH1 profile of backcross and F2 seeds as determined by SDS-PAGE analysis.

Cross	Sample	% Observed zH1 presence vs. zH1 absence	% Expected zH1 presence vs. zH1 absence ^a
RN x No2It	20	100:0	50:50
RN x Ro2R #1	71	93:7	50:50
RN x Ro2R #2	14	43:57	50:50
RN x Ro2R #3	14	92:8	50:50
NR X Ro2R #1	22	100:0	50:50
NR X Ro2R #2	48	85:15	50:50
RN self	24	96:4	50:50
NR self	60	93:7	50:50
Ro2R x RN	22	0:100	0:100
Ro2R x NR #1	31	0:100	0:100
Ro2R x NR #2	48	12:88	0:100

a; percentages are those expected if α -zeins expression is determined by a maternally expressed monogenic factor. The first genotype in the cross is the seed parent. When present # indicates that for a specific cross different seed progeny were analyzed.

<https://doi.org/10.1371/journal.pone.0206993.t002>

methylation and zein expression. Specifically, the *zH1b* and *zL1a* alleles were paternally methylated in both the RN and MN crosses, whereas suppression was only observed in the former cross. Additionally, a single methylated paternal allele of *zH1d* produced more transcripts in NR than two demethylated maternal alleles in RN (Figs 4A and 6B). To further analyze whether a possible link existed between DNA methylation and α -zein expression, we analyzed the promoter region, which is more likely to affect gene expression than coding regions [53,54]. Because the α -zein promoters display high sequence identity and because the promoter sequences of *zH1b*, *zH1c*, *zH1d* and *zL1a* are unknown, we focused on *zL1a*. This allele closely resembles the *zp22/D87* allele (S2 Fig) and is characterized by an internal deletion that permits it to be discriminated from other paralogous sequences. Indeed, PCR analysis of the entire set of inbred lines that were used in this study revealed that only the BSSS53 (used as control) and *No2It* line produced identical PCR patterns, which were characterized by two different PCR products (S3 Fig). The smaller product was genotype specific and identified the *zL1a;zp22/D87* allele (promoter and part of the coding region), whereas the larger product was common to all of the genotypes that were analyzed and reflected a non-specific PCR product that arose due to the high identity of zein promoter sequences. Methylation-dependent PCR analysis that was conducted with the methylation-dependent enzymes McrBC and MspJI revealed that the *zL1a* promoter region was methylated to a similar extent in both of the pairs of reciprocal crosses that were analyzed (Fig 6B). A final attempt to identify methylation differences that could explain maternal expression vs. paternal suppression of α -zeins in hybrid endosperms was made by using bisulfite sequence analysis. Specifically, we focused on a portion of both the promoter and the coding region of *zL1a*. Upon maternal transmission, *zL1a* was unmethylated in all cytosine contexts in both NR and NM hybrids. In contrast, upon paternal transmission, *zL1a* retained similar levels of CG/CHG methylation in both RN (suppressive hybrid) and MN (non-suppressive hybrid; Fig 6C).

Discussion

The parent-of-origin effect that is exhibited by α -zeins in hybrid endosperms has been described as an example of genomic imprinting [26], an epigenetic mechanism that leads to




Fig 6. DNA methylation analysis of uniparental expressed α -zeins. (A) Schematic representation of the restriction maps of the *zH1b*, *zH1c*, *zH1d* and *zL1a* alleles. The positions of *HhaI* (H) restriction sites are shown. The solid bar indicates the nucleotides from the ATG to the stop codon. The dotted lines represent nucleotide

deletions between *zL1a* and the other three sequences. Arrowheads indicate the positions of the primers that were used in the PCR amplification of the coding (black) and promoter regions (gray). The positions of the O2 site and the prolamin-binding site (PBS) are reported. (B) Methylation-sensitive (*HhaI*) and methylation-dependent (*McrBC* and *MspPII*) PCR analyses. Undigested DNA was used as a control (c). White letters indicate the parental origin of the α -zein alleles in the reciprocals: maternal (M), paternal (P). (C) Bisulfite sequencing analysis of the promoter and coding region of *zL1a* in the NR and RN and the NM and MN hybrid endosperms. Each line represents a single sequence. The red, blue and green colors indicate CG, CHG and CHH sites, respectively. The portion of the promoter and coding region analyzed is indicated by the graph.

<https://doi.org/10.1371/journal.pone.0206993.g006>

monoallelic expression of genes in a parent-of-origin manner [3]. This hypothesis was supported by the observations that some α -zein isoforms are only maternally expressed in hybrid endosperms and that reductions in DNA methylation, an epigenetic mark that is often associated with imprinting expression, are primarily restricted to the maternal complement of α -zeins [26]. However, because α -zeins belong to a large gene family, and because of their high sequence identity, the specific sequence targets of this phenomenon are unknown. By using reciprocal crosses that were made between different $\alpha 2$ genotypes and that showed a simplified α -zein pattern, we identified four alleles sensitive to parent-of-origin effects; this permitted us to analyze for the first time both the expression and DNA methylation patterns of these genes in a parental allele-specific manner.

The first finding of our study is that uniparental expression of α -zeins is not restricted to the maternal complement. For instance, in the NR endosperm, we observed α -zeins that exhibited both maternal- and paternal-biased gene expression. The *No2It* α -zein alleles show an expression pattern that is consistent with maternally expressed genes, because their expression occurs in both the *No2It* and the NR endosperms [55]. In contrast, the expression of the *Ro2* α -zein alleles was paternally enhanced in one of the two hybrid endosperms (NR), a behavior that resembled the dominance-imprinting pattern described in mice, in which the phenotype of one cross differs from the phenotype of its reciprocal and parental genotypes [55]. The coexistence of maternally and paternally expressed, or suppressed, α -zein alleles within the same hybrid endosperm suggests that the expression of these loci does not simply depend on their parental origin but instead on a specific mechanism/factor that is maternally provided by one of the lines used in the cross.

Indeed, even parental DNA methylation differences that we detected at these alleles did not apparently correlate with gene expression. In this regard, the best example is provided by the analysis of the *No2It*-specific *zL1a* allele, which is maternally expressed in both NR and NM crosses. DNA methylation profiling of its promoter region, which is considered a good predictor of transcription activity [50, 51], revealed that this allele upon maternal transmission was totally depleted of cytosine methylation in both NR and NM endosperms. On the other hand, in both RN and MN crosses *zL1a* was paternally methylated at CG and CHG sites to a similar degree in both RN and MN crosses, despite it was only suppressed in the former hybrid.

A possible explanation for this result is that our analysis did not capture differences in DNA methylation that are critical to gene expression. These differences might be present within unknown regulatory regions located either upstream or downstream of the coding region. For example, the paternal expression and maternal silencing of the Arabidopsis gene *PHERES1* depends on parental methylation differences that target repeats located 1.5–2.6 kb downstream of the 3' end of the gene [56]. Alternatively, the silencing of α -zeins may require additional epigenetic marks that may or may not act in conjunction with DNA methylation. Indeed, demethylation of the 22-kDa α -zein promoters in maize callus does not promote α -zein expression even when their positive regulatory transcription factors O2 and PBF are co-expressed, suggesting that factors in addition to DNA methylation are involved in the regulation of α -zein expression [51].

The type of regulation observed for α -zeins is comparable to that already described for another zein subfamily: the δ -zeins [5,47]. The 10-kDa and 18-kDa δ -zeins are encoded by two genes that show parent-of-origin effects in a genetic background-specific manner as well as dominance-imprinting expression [5,47]. It has been proposed that the non-Mendelian regulation of δ -zeins is due to parental imprinting of the regulatory *dzt1* locus that post-transcriptionally controls their accumulation [5,47]. Thus, the simplest explanation is that expression of α -zeins may also depend on a regulatory factor that is itself the target of epigenetic regulation. Although the O2 protein is the main regulator of 22-kDa α -zeins, additional regulators have been identified [36,37,38,39,40,41,42,43], and others are likely to exist [57,58]. Interestingly, the *azs22/D87* and *azs22/6* alleles, namely, *zL1a* and *zH1b* in this study, respectively, are also expressed in the BSSS53 $\alpha 2$ line, supporting the idea that these alleles have become O2-independent [44]. Our transcriptional analysis of the α -zein regulators so far identified did not evidence expression patterns that may explain the uniparental expression of these genes. However, as mentioned in the Results section, our analysis only disclosed the lack of gross transgressive expression patterns at these regulatory loci in the NR and RN hybrids, and more detailed investigations will be required to know definitively whether these loci are involved in the phenomenon under study.

Supposing the existence of an endosperm factor, maternally expressed, that may either promote (*No2It*-specific) or suppress (*Ro2R*-specific) α -zein expression, one would expect to find the zH1-plus and zH1-null phenotype in 50% of seeds that have either the RN or the NR genotype as the maternal parent, respectively. In contrast, a penetrant zH1-null phenotype would be expected in endosperms that have the *Ro2R* genotype as the maternal parent. Our results reveal a complex situation that is not readily understood. The *Ro2R* x RN and *Ro2R* x NR endosperms displayed a penetrant zH1-null pattern, though not exclusive, which was comparable with that observed in the RN hybrid. In contrast, a more variable expression pattern was observed in RN x *Ro2R*, NR x *Ro2R*, RN x RN, and NR x NR endosperms. This behavior cannot be simply explained by the presence of a monogenic factor that is provided either by the *No2It* or *Ro2R* lines within the endosperm. Even considering the eventuality that a factor is provided by the seed parent either at cytoplasmic or sporophytic level, one would expect that all seeds of the above backcrosses and F2 would express α -zeins; this is clearly not the case. The complexity of these expression patterns is further evidenced by the findings that α -zeins of the zH2 types are reactivated in some backcrosses and F2 seeds. In this case, a simple mechanism of complementation between a regulatory factor and its target loci seems unlikely, because this condition would be already satisfied in the F1 hybrids. This latter phenomenon resembles the transgressive expression patterns that is observed in tomato, where novel expression states are observed in the F2 and in advanced generation [59].

Interestingly, there is again a commonality between the expression proprieties of α -zeins and δ -zeins. For instance, the ability of certain lines to suppress maternally the expression of the 18 kDa δ -zein is abolished after a round of heterozygosity with a non-suppressing line. To explain this finding, Wu and colleagues [47] suggested that *dzt1* might also be sensitive to paramutation: an epigenetic mechanism in which one allele causes a heritable change in the expression of a homologous allele [60]. These trans-allelic interactions are mediated by small interfering RNAs and lead to either losses or gains of DNA methylation at target loci [61,62,63]. Importantly, the same trans allelic interactions also generate transgressive expression patterns in hybrid plants and in advanced generations such as the F2 [59, 60]. We wonder if such transgressive behavior may be also depend on epigenome shock that is triggered by the epigenetic asymmetry of the endosperm parental genomes [17,64], likewise that one recently observed in the epihybrids of *Arabidopsis* [65].

Although at this time, our data cannot clarify the extent to which uniparental expression (F1) and transgressive expression (F2 and backcross) of α -zeins are linked, and if these phenomena might reflect, at least in part, paramutation-like phenomena, they show how gene expression strongly depends on genotype interactions that occur in hybrid/recombinant endosperms. Understanding the molecular basis of this phenomenon may help to clarify how hidden parental traits can be unlocked in advanced generations.

Supporting information

S1 Fig. Amino acid sequence comparison of α -zein identified in this study. The amino acid sequence of *zL1a*, *zH1b*, *zH1c* and *zH1d* was in silico deduced. Gaps (dotted lines) were introduced to maximize homology. At the amino-termini, the amino acids of each sequence that match those determined by the direct sequences of the two corresponding polypeptides are underlined.

(TIF)

S2 Fig. Neighbor-joining tree of α -zeins identified in this study with α -zeins identified in other maize lines. Zein sequences of the BSSS53 (BSSS), B73 and W22 inbred lines [44,46,66] were aligned with the *zL1a*, *zH1b*, *zH1c* and *zH1d* sequences (black arrows), and a phylogenetic Neighbor-joining tree was generated by using CLC Sequence Viewer. The light-blue boxes show the α -zein composition of the *z1C1* and *z1C2* loci among different inbred lines. Black numbers indicate the presence of a specific α -zein allele, whereas gray numbers indicate that in the *No2It* and *Ro2* lines the presence of a specific α -zein allele is unknown.

(TIF)

S3 Fig. PCR analysis of maize lines used in this study for the presence of the *zL1a* allele.

The BSSS53 and B73 inbred line are used as positive and negative control, respectively, for the presence of the *zL1a* allele.

(TIF)

S1 Table. List of primers used in the study and experimental conditions used.

(DOCX)

Acknowledgments

We thank Mario Motto and Serena Varotto for helpful comments on the manuscript, E. Quatrini for taking care of the plants, and anonymous reviewers for their helpful comments on an early version of this manuscript. Research was supported by a grant from MIUR, FIRB project No. RBNE01TYZF, and a special grant from MIUR and CNR for the Epigenomics Flagship Project (EPIGEN).

Author Contributions

Conceptualization: Angelo Viotti.

Funding acquisition: Angelo Viotti, Massimiliano Lauria.

Investigation: Silvana Castelli, Iride Mascheretti, Cristian Cosentino, Barbara Lazzari, Raul Pirona, Aldo Ceriotti, Massimiliano Lauria.

Project administration: Massimiliano Lauria.

Resources: Angelo Viotti, Massimiliano Lauria.

Supervision: Angelo Viotti, Massimiliano Lauria.

Validation: Silvana Castelli, Iride Mascheretti, Cristian Cosentino, Barbara Lazzari, Raul Pirona, Aldo Ceriotti, Massimiliano Lauria.

Visualization: Massimiliano Lauria.

Writing – original draft: Silvana Castelli, Angelo Viotti.

Writing – review & editing: Massimiliano Lauria.

References

1. Lisch D (2002) Mutator transposons. *Trends Plant Sci* 7: 498–504. PMID: [12417150](#)
2. Hollick JB (2017) Paramutation and related phenomena in diverse species. *Nature Reviews Genetics* 18: 5–23. <https://doi.org/10.1038/nrg.2016.115> PMID: [27748375](#)
3. Reik W, Walter J (2001) Genomic imprinting: parental influence on the genome. *Nat Rev Genet* 2: 21–32. <https://doi.org/10.1038/35047554> PMID: [11253064](#)
4. Kermicle JL (1970) Dependence of the R-mottled aleurone phenotype in maize on mode of sexual transmission. *Genetics* 66: 69–85. PMID: [17248508](#)
5. Chaudhuri S, Messing J (1994) Allele-specific parental imprinting of *dzt1*, a posttranscriptional regulator of zein accumulation. *Proc Natl Acad Sci U S A* 91: 4867–4871. PMID: [8197149](#)
6. Lund G, Messing J, Viotti A (1995) Endosperm-specific demethylation and activation of specific alleles of alpha-tubulin genes of *Zea mays* L. *Mol Gen Genet* 246: 716–722. PMID: [7898439](#)
7. Pirona R, Hartings H, Lauria M, Rossi V, Motto M (2005) Genetic control of endosperm development and of storage products accumulation in maize seeds. *Maydica* 50: 16.
8. Grossniklaus U, Vielle-Calzada JP, Hoepfner MA, Gagliano WB (1998) Maternal control of embryogenesis by MEDEA, a polycomb group gene in *Arabidopsis*. *Science* 280: 446–450. PMID: [9545225](#)
9. Chaudhury AM, Ming L, Miller C, Craig S, Dennis ES, Peacock WJ (1997) Fertilization-independent seed development in *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A* 94: 4223–4228. PMID: [9108133](#)
10. Ohad N, Margossian L, Hsu YC, Williams C, Repetti P, Fischer RL (1996) A mutation that allows endosperm development without fertilization. *Proc Natl Acad Sci U S A* 93: 5319–5324. PMID: [11607683](#)
11. Costa LM, Yuan J, Rouster J, Paul W, Dickinson H, Gutierrez-Marcos JF (2012) Maternal control of nutrient allocation in plant seeds by genomic imprinting. *Curr Biol* 22: 160–165. <https://doi.org/10.1016/j.cub.2011.11.059> PMID: [22245001](#)
12. Wolff P, Weinhofer I, Seguin J, Roszak P, Beisel C, Donoghue MT, et al. (2011) High-resolution analysis of parent-of-origin allelic expression in the *Arabidopsis* Endosperm. *PLoS Genet* 7: e1002126. <https://doi.org/10.1371/journal.pgen.1002126> PMID: [21698132](#)
13. Gehring M, Missirian V, Henikoff S (2011) Genomic analysis of parent-of-origin allelic expression in *Arabidopsis thaliana* seeds. *PLoS One* 6: e23687. <https://doi.org/10.1371/journal.pone.0023687> PMID: [21858209](#)
14. Luo M, Taylor JM, Spriggs A, Zhang H, Wu X, Russell S, et al. (2011) A genome-wide survey of imprinted genes in rice seeds reveals imprinting primarily occurs in the endosperm. *PLoS Genet* 7: e1002125. <https://doi.org/10.1371/journal.pgen.1002125> PMID: [21731498](#)
15. Xu W, Dai M, Li F, Liu A (2014) Genomic imprinting, methylation and parent-of-origin effects in reciprocal hybrid endosperm of castor bean. *Nucleic Acids Res* 42: 6987–6998. <https://doi.org/10.1093/nar/gku375> PMID: [24799438](#)
16. Xin M, Yang R, Li G, Chen H, Laurie J, Ma C, et al. (2013) Dynamic expression of imprinted genes associates with maternally controlled nutrient allocation during maize endosperm development. *Plant Cell* 25: 3212–3227. <https://doi.org/10.1105/tpc.113.115592> PMID: [24058158](#)
17. Waters AJ, Makarevitch I, Eichten SR, Swanson-Wagner RA, Yeh CT, Xu W, et al. (2011) Parent-of-origin effects on gene expression and DNA methylation in the maize endosperm. *Plant Cell* 23: 4221–4233. <https://doi.org/10.1105/tpc.111.092668> PMID: [22198147](#)
18. Zhang M, Zhao H, Xie S, Chen J, Xu Y, Wang K, et al. (2011) Extensive, clustered parental imprinting of protein-coding and noncoding RNAs in developing maize endosperm. *Proc Natl Acad Sci U S A* 108: 20042–20047. <https://doi.org/10.1073/pnas.1112186108> PMID: [22114195](#)
19. Xin M, Yang R, Yao Y, Ma C, Peng H, Sun Q, et al. (2014) Dynamic parent-of-origin effects on small interfering RNA expression in the developing maize endosperm. *BMC Plant Biol* 14: 192. <https://doi.org/10.1186/s12870-014-0192-8> PMID: [25055833](#)

20. Pignatta D, Erdmann RM, Scheer E, Picard CL, Bell GW, Gehring M. (2014) Natural epigenetic polymorphisms lead to intraspecific variation in Arabidopsis gene imprinting. *Elife* 3: e03198. <https://doi.org/10.7554/eLife.03198> PMID: 24994762
21. Waters AJ, Bilinski P, Eichten SR, Vaughn MW, Ross-Ibarra J, Springer NM (2013) Comprehensive analysis of imprinted genes in maize reveals allelic variation for imprinting and limited conservation with other species. *Proc Natl Acad Sci U S A* 110: 19639–19644. <https://doi.org/10.1073/pnas.1309182110> PMID: 24218619
22. Gehring M, Bubb KL, Henikoff S (2009) Extensive demethylation of repetitive elements during seed development underlies gene imprinting. *Science* 324: 1447–1451. <https://doi.org/10.1126/science.1171609> PMID: 19520961
23. Lauria M, Rupe M, Guo M, Kranz E, Pirona R, Viotti A, et al. (2004) Extensive maternal DNA hypomethylation in the endosperm of *Zea mays*. *Plant Cell* 16: 510–522. <https://doi.org/10.1105/tpc.017780> PMID: 14729913
24. Zemach A, Kim MY, Silva P, Rodrigues JA, Dotson B, Zilberman D. (2010) Local DNA hypomethylation activates genes in rice endosperm. *Proc Natl Acad Sci U S A* 107: 18729–18734. <https://doi.org/10.1073/pnas.1009695107> PMID: 20937895
25. Hsieh TF, Ibarra CA, Silva P, Zemach A, Eshed-Williams L, Fischer RL et al. (2009) Genome-wide demethylation of Arabidopsis endosperm. *Science* 324: 1451–1454. <https://doi.org/10.1126/science.1172417> PMID: 19520962
26. Lund G, Ciceri P, Viotti A (1995) Maternal-specific demethylation and expression of specific alleles of zein genes in the endosperm of *Zea mays* L. *Plant J* 8: 571–581. PMID: 7496402
27. Xu JH, Messing J (2008) Organization of the prolamin gene family provides insight into the evolution of the maize genome and gene duplications in grass species. *Proc Natl Acad Sci U S A* 105: 14330–14335. <https://doi.org/10.1073/pnas.0807026105> PMID: 18794528
28. Viotti A, Cairo G, Vitale A, Sala E (1985) Each zein gene class can produce polypeptides of different sizes. *Embo J* 4: 1103–1110. PMID: 16453609
29. Ciceri P, Castelli S, Lauria M, Lazzari B, Genga A, Bernard L, et al. (2000) Specific combinations of zein genes and genetic backgrounds influence the transcription of the heavy-chain zein genes in maize opaque-2 endosperms. *Plant Physiol* 124: 451–460. PMID: 10982458
30. Schmidt RJ, Burr FA, Aukerman MJ, Burr B (1990) Maize regulatory gene opaque-2 encodes a protein with a "leucine-zipper" motif that binds to zein DNA. *Proc Natl Acad Sci U S A* 87: 46–50. PMID: 2296602
31. Sekhon RS, Lin HN, Childs KL, Hansey CN, Buell CR, de Leon N. et al. (2011) Genome-wide atlas of transcription during maize development. *Plant Journal* 66: 553–563. <https://doi.org/10.1111/j.1365-3113X.2011.04527.x> PMID: 21299659
32. Gruntman E, Qi YJ, Slotkin RK, Roeder T, Martienssen RA, Sachidanandam R, et al. (2008) Kismeth: Analyzer of plant methylation states through bisulfite sequencing. *Bmc Bioinformatics* 9.
33. Bernard L, Ciceri P, Viotti A (1994) Molecular analysis of wild-type and mutant alleles at the Opaque-2 regulatory locus of maize reveals different mutations and types of O2 products. *Plant Mol Biol* 24: 949–959. PMID: 8204831
34. Feng L, Zhu J, Wang G, Tang Y, Chen H, Jin W et al. (2009) Expressional profiling study revealed unique expressional patterns and dramatic expressional divergence of maize alpha-zein super gene family. *Plant Mol Biol* 69: 649–659. <https://doi.org/10.1007/s11103-008-9444-z> PMID: 19112555
35. Gavazzi F, Lazzari B, Ciceri P, Gianazza E, Viotti A (2007) Wild-type opaque2 and defective opaque2 polypeptides form complexes in maize endosperm cells and bind the opaque2-zein target site. *Plant Physiol* 145: 933–945. <https://doi.org/10.1104/pp.107.103606> PMID: 17827273
36. Gontarek BC, Neelakandan AK, Wu H, Becraft PW (2016) NKD Transcription Factors Are Central Regulators of Maize Endosperm Development. *Plant Cell* 28: 2916–2936. <https://doi.org/10.1105/tpc.16.00609> PMID: 27895224
37. Li Q, Wang JC, Ye JW, Zheng XX, Xiang XL, Li C et al. (2017) The Maize Imprinted Gene Flouy3 Encodes a PLATZ Protein Required for tRNA and 5S rRNA Transcription through Interaction with RNA Polymerase III. *Plant Cell* 29: 2661–2675. <https://doi.org/10.1105/tpc.17.00576> PMID: 28874509
38. Zhang N, Qiao Z, Liang Z, Mei B, Xu Z, Song R (2012) *Zea mays* Taxilin protein negatively regulates opaque-2 transcriptional activity by causing a change in its sub-cellular distribution. *PLoS One* 7: e43822. <https://doi.org/10.1371/journal.pone.0043822> PMID: 22937104
39. Bhat RA, Borst JW, Riehl M, Thompson RD (2004) Interaction of maize Opaque-2 and the transcriptional co-activators GCN5 and ADA2, in the modulation of transcriptional activity. *Plant Mol Biol* 55: 239–252. <https://doi.org/10.1007/s11103-004-0553-z> PMID: 15604678

40. Vicente-Carbajosa J, Moose SP, Parsons RL, Schmidt RJ (1997) A maize zinc-finger protein binds the prolamins box in zein gene promoters and interacts with the basic leucine zipper transcriptional activator Opaque2. *Proc Natl Acad Sci U S A* 94: 7685–7690. PMID: [9207153](#)
41. Pysh LD, A M, Schmidt RJ (1993) OHP1: a maize basic domain/leucine zipper protein that interacts with opaque2. *Plant Cell* 5: 227. <https://doi.org/10.1105/tpc.5.2.227> PMID: [8453304](#)
42. Qiao Z, Qi W, Wang Q, Feng Y, Yang Q, Zhang N et al. (2016) ZmMADS47 Regulates Zein Gene Transcription through Interaction with Opaque2. *PLoS Genet* 12: e1005991. <https://doi.org/10.1371/journal.pgen.1005991> PMID: [27077660](#)
43. Feng F, Qi WW, Lv YD, Yan SM, Xu LM, Yang W et al. (2018) OPAQUE11 Is a Central Hub of the Regulatory Network for Maize Endosperm Development and Nutrient Metabolism. *Plant Cell* 30: 375–396. <https://doi.org/10.1105/tpc.17.00616> PMID: [29436476](#)
44. Song R, Llaca V, Linton E, Messing J (2001) Sequence, regulation, and evolution of the maize 22-kD alpha zein gene family. *Genome Res* 11: 1817–1825. <https://doi.org/10.1101/gr.197301> PMID: [11691845](#)
45. Messing J (1987) The genes encoding seed storage proteins in higher plants. In *Genetic Engineering*, Vol 6, (Rigby PWJ, ed) Harcourt Brace Jovanovich Publishers, London: 1–46.
46. Song RT, Messing J (2003) Gene expression of a gene family in maize based on noncollinear haplotypes. *Proceedings of the National Academy of Sciences of the United States of America* 100: 9055–9060. <https://doi.org/10.1073/pnas.1032999100> PMID: [12853580](#)
47. Wu Y, Goettel W, Messing J (2009) Non-Mendelian regulation and allelic variation of methionine-rich delta-zein genes in maize. *Theor Appl Genet* 119: 721–731. <https://doi.org/10.1007/s00122-009-1083-5> PMID: [19504256](#)
48. Bianchi MW, Viotti A (1988) DNA methylation and tissue-specific transcription of the storage protein genes of maize. *Plant Mol Biol* 11: 203–214. <https://doi.org/10.1007/BF00015672> PMID: [24272262](#)
49. Locatelli S, Piatti P, Motto M, Rossi V (2009) Chromatin and DNA modifications in the Opaque2-mediated regulation of gene transcription during maize endosperm development. *Plant Cell* 21: 1410–1427. <https://doi.org/10.1105/tpc.109.067256> PMID: [19482970](#)
50. Sturaro M, Viotti A (2001) Methylation of the Opaque2 box in zein genes is parent-dependent and affects O2 DNA binding activity in vitro. *Plant Mol Biol* 46: 549–560. PMID: [11516148](#)
51. Wu Y, Messing J (2012) Rapid divergence of prolamins gene promoters of maize after gene amplification and dispersal. *Genetics* 192: 507–519. <https://doi.org/10.1534/genetics.112.142372> PMID: [22798485](#)
52. Miclaus M, Xu JH, Messing J (2011) Differential gene expression and epiregulation of alpha zein gene copies in maize haplotypes. *PLoS Genet* 7: e1002131. <https://doi.org/10.1371/journal.pgen.1002131> PMID: [21731501](#)
53. Zhang X, Yazaki J, Sundaresan A, Cokus S, Chan SW, Chen H et al. (2006) Genome-wide high-resolution mapping and functional analysis of DNA methylation in Arabidopsis. *Cell* 126: 1189–1201. <https://doi.org/10.1016/j.cell.2006.08.003> PMID: [16949657](#)
54. Zilberman D, Gehring M, Tran RK, Ballinger T, Henikoff S (2007) Genome-wide analysis of Arabidopsis thaliana DNA methylation uncovers an interdependence between methylation and transcription. *Nat Genet* 39: 61–69. <https://doi.org/10.1038/ng1929> PMID: [17128275](#)
55. Wolf JB, Cheverud JM, Roseman C, Hager R (2008) Genome-wide analysis reveals a complex pattern of genomic imprinting in mice. *PLoS Genet* 4: e1000091. <https://doi.org/10.1371/journal.pgen.1000091> PMID: [18535661](#)
56. Makarevich G, Villar CB, Erilova A, Kohler C (2008) Mechanism of PHERES1 imprinting in Arabidopsis. *J Cell Sci* 121: 906–912. <https://doi.org/10.1242/jcs.023077> PMID: [18303047](#)
57. Fu J, Cheng Y, Linghu J, Yang X, Kang L, Zhang Z et al. (2013) RNA sequencing reveals the complex regulatory network in the maize kernel. *Nat Commun* 4: 2832. <https://doi.org/10.1038/ncomms3832> PMID: [24343161](#)
58. Li C, Qiao Z, Qi W, Wang Q, Yuan Y, Yang X et al. (2015) Genome-Wide Characterization of cis-Acting DNA Targets Reveals the Transcriptional Regulatory Framework of Opaque2 in Maize. *Plant Cell*.
59. Shivaprasad PV, Dunn RM, Santos BA, Bassett A, Baulcombe DC (2012) Extraordinary transgressive phenotypes of hybrid tomato are influenced by epigenetics and small silencing RNAs. *Embo J* 31: 257–266. <https://doi.org/10.1038/emboj.2011.458> PMID: [22179699](#)
60. Erhard KF Jr., Hollick JB (2011) Paramutation: a process for acquiring trans-generational regulatory states. *Curr Opin Plant Biol* 14: 210–216. <https://doi.org/10.1016/j.pbi.2011.02.005> PMID: [21420347](#)
61. Greaves IK, Groszmann M, Ying H, Taylor JM, Peacock WJ, Dennis ES (2012) Trans chromosomal methylation in Arabidopsis hybrids. *Proc Natl Acad Sci U S A* 109: 3570–3575. <https://doi.org/10.1073/pnas.1201043109> PMID: [22331882](#)

62. He G, Chen B, Wang X, Li X, Li J, He H et al. (2013) Conservation and divergence of transcriptomic and epigenomic variation in maize hybrids. *Genome Biol* 14: R57. <https://doi.org/10.1186/gb-2013-14-6-r57> PMID: 23758703
63. He G, Zhu X, Elling AA, Chen L, Wang X, Guo L et al. (2010) Global epigenetic and transcriptional trends among two rice subspecies and their reciprocal hybrids. *Plant Cell* 22: 17–33. <https://doi.org/10.1105/tpc.109.072041> PMID: 20086188
64. Zhang M, Xie S, Dong X, Zhao X, Zeng B, Chen J et al. (2013) Genome-wide high resolution parental-specific DNA and histone methylation maps uncover patterns of imprinting regulation in maize. *Genome Res* 24: 167–176. <https://doi.org/10.1101/gr.155879.113> PMID: 24131563
65. Rigal M, Becker C, Pelissier T, Pogorelcnik R, Devos J, Ikeda Yet al. (2016) Epigenome confrontation triggers immediate reprogramming of DNA methylation and transposon silencing in *Arabidopsis thaliana* F1 epihybrids. *Proceedings of the National Academy of Sciences of the United States of America* 113: E2083–E2092. <https://doi.org/10.1073/pnas.1600672113> PMID: 27001853
66. Dong JQ, Feng YP, Kumar D, Zhang W, Zhu TT, Luo MC et al. (2016) Analysis of tandem gene copies in maize chromosomal regions reconstructed from long sequence reads. *Proceedings of the National Academy of Sciences of the United States of America* 113: 7949–7956. <https://doi.org/10.1073/pnas.1608775113> PMID: 27354512