

Closed Limb Fractures With Compromised Vascularization: A Narrative Review

Christopher Vannabouathong¹ , Emil Schemitsch², Bradley Petrisor³ and Mohit Bhandari³

¹OrthoEvidence, Inc, Burlington, ON, Canada. ²Department of Surgery, Western University, London, ON, Canada. ³Department of Surgery, McMaster University, Hamilton, ON, Canada.

Clinical Medicine Insights: Arthritis and Musculoskeletal Disorders
Volume 12: 1–6
© The Author(s) 2019
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1179544119836742



ABSTRACT: Vascular compromised fractures typically result in a high rate of healing complications, such as avascular necrosis, nonunion, delayed union, and arthritis, which severely affect a patient's function and quality of life. The purpose of this review was to identify and describe the epidemiology and available treatment options for the most well-known vascular compromised closed fractures. The injuries discussed in detail in this review were scaphoid, lunate, femoral neck, and talar fractures. Current evidence suggests that optimal treatment for vascular compromised fractures is dependent on the degree of fracture displacement and comminution, and the patient's post-injury functional demands, age, and bone quality. Conservative efforts generally include casting or splinting with a period of immobilization. Surgery is indicated for substantially displaced fractures, patients who require higher functional demands and an earlier return to activity, or if complications occur following nonoperative treatment; however, operative intervention is typically performed for femoral neck fractures regardless of the amount of displacement. Various surgical techniques exist, though internal fixation with screws is a common procedure among these injuries and can be used in combination with other implants, such as plating or Kirschner wires (k-wires), when needed. Severe fracture comminution, poor bone quality, or arthritis can contraindicate the use of screws and more invasive intervention will be required. Bone grafting is done in some cases to enhance vascularity. Salvage procedures exist for patients who develop severe complications, but these will permanently alter the anatomy of the injured area and should be considered a last resort.

KEYWORDS: Vascular compromised fracture, scaphoid, lunate, femoral neck, talus

RECEIVED: January 30, 2019. **ACCEPTED:** February 12, 2019.

TYPE: Review

FUNDING: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The preparation of the manuscript was funded, in part, by a research grant from Acumed. Christopher Vannabouathong is an employee of OrthoEvidence, Inc.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Christopher Vannabouathong, OrthoEvidence, Inc, Burlington, ON L7N 3H8, Canada. Email: chris.vannabouathong@myorthovidence.com

Introduction

Vascular compromised fractures are injuries with a severely impaired healing process due to the limited blood flow to these areas.^{1–4} Because of this, fractures in these regions are associated with a high rate of necrosis and union complications, making the management of these injuries challenging.^{1,4–7} The purpose of this review was to identify and describe the epidemiology and treatment options for the most well-known vascular compromised closed fractures.

Background on scaphoid and lunate fractures

Fractures of the scaphoid represent about 50% to 90% of all carpal fractures in young, physically active individuals.^{1,8} Approximately 345 000 scaphoid injuries occur in the United States every year.³ With these injuries, the patient will present with wrist pain on the radial side after a traumatic event.⁹ There may also be tenderness in the anatomical snuffbox, pain on palpation of the scaphoid, pain with axial compression of the thumb, and pain in the thumb during range of motion (ROM).⁹ These clinical signs are important to investigate, as the injury may be missed on plain radiographs.⁹ Advanced imaging will help determine fracture stability and amount of displacement.⁹ Because of the vascular insufficiency present in the scaphoid bone, nonunion is a common complication following injury to this area, with rates ranging from 5% to 15%.^{1,3}

Other concerns include delayed union, arthritis, reduced wrist motion, and loss of strength.³ The probability of a scaphoid nonunion will be reduced with early recognition, appropriate treatment, and immobilization.¹⁰

Isolated lunate fractures are rare as they are typically associated with other wrist injuries; however, there are concerns that the injury is being missed on diagnosis and computerized tomography (CT) or magnetic resonance imaging (MRI) may be needed for proper identification.^{11–13} They account for approximately 0.5% to 6.5% of carpal fractures.^{11–13} The most common mechanism of action is falling on an outstretched hand, but the injury may occur following hyperflexion or a direct blow.¹³ Teisen and Hjarbaek categorized these fractures based on radiographs and vascularity, as described by Gelberman and Gross (Table 1).^{11,14,15}

Treatment of scaphoid fractures

If the injury is deemed minimally displaced, cast or splint immobilization may be employed; however, fracture consolidation via this method ranges from 8 to 12 weeks, which may be too long for a younger, more active patient.^{8,9,16} Once union is achieved, hand therapy may begin.⁹ Complications, including muscular atrophy, joint stiffness, and reduced grip strength, may occur and could delay the patient's return to work and activities.⁸



Table 1. The Teisen and Hjarbaek classification of lunate fractures.

Group I	Fracture of the volar pole, possibly affecting the volar nutrient artery
II	Chip fracture not affecting the main blood supply
III	Fracture of the dorsal pole, possibly affecting the dorsal nutrient artery
IV	Sagittal fracture through the body of the lunate
V	Transverse fracture through the body of the lunate

Different surgical fixation methods, screws, bone grafts, and approaches have been investigated in the literature.^{1,3,17} Bone grafting, typically from the radius, is done to enhance vascularity.^{1,10} Screw fixation provides structural support by compressing the 2 fragments and prevents bending at the fracture site; a single headless compression screw along the central longitudinal axis is the most common method of fixation.^{1,3,9,10} Designs have evolved since the original Herbert screw, claiming to improve load to failure, stiffness, and compression at the fracture site, though screw selection is currently based on the surgeon's personal preference, as a better understanding of the different compressive forces and clinical effects generated by these various screw designs is still required.^{9,18} Kirschner wires (k-wires) are helpful for stabilizing the ligamentous forces acting on the area and involve minor soft tissue damage, but can lead to pin tract infection, stiffness, and the need for re-intervention.^{1,8} Volar plating may be indicated when there is a large degree of comminution or bone loss.^{9,10} Generally, internal fixation is associated with good union rates, ranging from 85% to 90%, while also providing earlier return to mobilization.^{8,18} Not only is vascular compromise an issue, but instability can also be the cause of complications, such as nonunion, post-injury; therefore, a combination of these different treatments may be needed to address all underlying issues in some of these patients.¹ In a biomechanical study, Guss et al³ found that a 50% intact scaphoid with a headless compression screw provided similar load to failure and stiffness to a fully intact, uninjured scaphoid. There is inconsistent evidence on the use of k-wires alone, but in a case-control study of 25 patients with scaphoid nonunion, Allon et al¹ found that the addition of k-wires to headless screw fixation and bone grafting was safe and should be considered when suboptimal stability is present during surgery. When a nonunion occurs, salvage procedures include proximal row carpectomy, scaphoid excision with arthrodesis, or total wrist fusion; however, these operations can lead to permanent changes in wrist function.¹⁰ For such cases, fixation with a headless compression screw, with or without bone grafting, is currently the standard when attempting to stabilize and unite the scaphoid.^{10,19} Metaphyseal decompression of the distal radius, along with the insertion of compression screws, is a more recently developed method that

may increase vascularization of the radial column of the carpus for scaphoid nonunion.¹⁷ The addition of a volar plate to screw fixation can provide more buttress support, if needed (Figure 1).¹⁰

Treatment of lunate fractures

Conservative treatment includes prolonged immobilization with a plaster cast for at least 6 weeks, which may be reserved for undisplaced fractures.^{11,12,20}

Displaced lunate injuries are treated via internal fixation with screws, and possibly k-wires, as it ensures good reduction and compression; however, this is not possible if the fracture is comminuted (Figure 1).¹¹⁻¹³ This treatment method is also an alternative for patients who do not want to experience prolonged immobilization (ie, want a quicker return to work) when the injury is treated conservatively.^{11,12} If chronic instability or arthrosis develops, salvage procedures include proximal row carpectomy and wrist arthrodesis (Figure 1).¹²

Background on femoral neck fractures

The number of hip fractures is rising due to the aging population, though such injuries can still occur in young adults.²¹⁻²³ In a recent study conducted by Rosengren et al,²⁴ the authors reported that the number of people aged ≥ 50 years rose from 1.6 to 2 million from 1980 to 2010 in Denmark and from 2.8 to 3.5 million from 1987 to 2011 in Sweden. During these same years, the number of annual hip fractures (per 10000 person-years) generally increased in these countries, from ~ 7 in men and ~ 28 in women to ~ 14 and ~ 39 , respectively, in Denmark and from ~ 23 to ~ 29 in Swedish men (incidence rates in Swedish women were similar between 1987 and 2010; $\sim 75/10000$ person-years).²⁴

Femoral neck fractures, in particular, are challenging as they are at an increased risk of complications, such as fixation failure, malunion, nonunion, and avascular necrosis of the femoral head.^{23,25} They may occur due to either low- or high-energy trauma, or because of a sport injury, and can be classified using the Garden or Pauwels classification system (Tables 2 and 3).^{5,25-27} Femoral neck fractures are a leading cause of disability and morbidity in the elderly, and, in the United States, yearly incidence rates for this injury have ranged from 146 to 242 per 100000 adults.²⁸ Displaced fractures, relative to nondisplaced, are associated with higher complication rates.^{5,29} For example, Campenfeldt et al⁵ found rates of 23%, 15%, and 28% for nonunion, avascular necrosis, and reoperation, respectively, in patients with displaced fractures, whereas these rates for those with nondisplaced fractures were 0%, 12%, and 8%.

Treatment of femoral neck fractures

The aim of surgery is to reduce pain and allow for early mobilization.²¹ It is believed that early fracture reduction and stabilization via surgery may improve femoral head blood flow, resulting in a greater likelihood of union and less complications;




Figure 1. Overview and treatment of scaphoid and lunate fractures.

Table 2. The Garden classification of femoral neck fractures.

Type	I	Non-displaced, incomplete fracture
	II	Non-displaced, complete fracture
	III	Partially displaced, complete fracture
	IV	Fully displaced, complete fracture

Table 3. The Pauwels classification of femoral neck fractures.

Type	I	Up to 30° between the fracture line of the distal fragment and the horizontal line—compressive forces are dominant
	II	Within 30° to 50° between the fracture line of the distal fragment and the horizontal line—shearing force occurs and may have a negative effect on bone healing
	III	≥50° between the fracture line of the distal fragment and the horizontal line—shearing force is predominant and is associated with a significant amount of varus force which will more likely result in fracture displacement and varus collapse

delayed treatment may also increase the risk of mortality.^{23,25,30–32} Although there is no definitive criteria as to what is considered an “unacceptable delay” for surgical intervention, guidelines

suggest that surgery should be performed within 24 h of injury.³² Surgical options for a femoral neck fracture include internal fixation or an arthroplasty procedure.^{5,21,22,28} A patient less than 70 years old should undergo internal fixation, regardless of the degree of displacement, which is hypothesized to preserve the hip joint while allowing the patient to function.^{5,21,23,25} Relative to nonoperative treatment, internal fixation lessens the risk of displacement and allows earlier weightbearing and rehabilitation.³³ Another reason internal fixation is recommended for patients <60 years is that they are at a higher risk of experiencing a revision with a total hip replacement due to their greater life expectancy.⁵ The procedure can be done with screws, resulting in good functional outcomes for most patients.^{5,23,25} For example, 76% of patients in the study by Campenfeldt et al⁵ achieved a “good” or “excellent” result on the Harris Hip Score 24 months post-surgery. There is variation in the size and number of screws, and their angle of insertion and placement, with no clear consensus on the best approach.^{23,25} Bone grafting may be done along with compression screw insertion to help improve vascularity in the region.^{23,33} Another method of internal fixation includes a sliding hip screw.^{25,33} After screw insertion, the sliding and compressing of fragments during loading are hypothesized to induce fracture healing while controlling the shear forces, but femoral neck shortening may




Figure 2. Overview and treatment of femoral neck fractures.

occur.²⁵ The patient's bone quality and the amount of fracture comminution are other factors to consider as they may contraindicate internal fixation.^{22,25,29} Arthroplasty is typically done for patients who are older (ie, ≥ 60 years) and have displaced fractures, or when a patient experiences a failure with less invasive treatment.^{21,22,25,28} For older patients, arthroplasty promotes early mobilization and weightbearing, reducing the likelihood that the patient will remain bedridden and encounter subsequent comorbidities (Figure 2).²²

Background on talar fractures

Talar fractures account for about 0.1% to 0.3% of all fractures and 1% of foot and ankle fractures.^{4,7,34} The talus is the most superiorly located bone of the foot and is vital in maintaining ankle function and ROM.⁶ Injuries are typically due to high-energy trauma, such as a motor vehicle collision or fall.^{7,34} Plain radiographs will be required for diagnosis, but a CT scan or MRI may be used to further characterize the fracture.^{7,35} Clinically, the patient will have pain and swelling at the ankle with the inability to bear weight.⁷ Talar fractures may be missed on initial examination, being diagnosed as an ankle sprain.³⁶

Table 4. The Hawkins classification of talar neck fractures.

Type I	Non-displaced
II	Displacement with subluxation/dislocation of the subtalar joint
III	Displacement with subluxation/dislocation of the subtalar and ankle joints
IV	Displacement with subluxation/dislocation of the subtalar, ankle, and talonavicular joints

Changes in gait will be observed and post-traumatic arthritis is another concern.^{7,34}

The talus comprises a neck, head, body, lateral process, and posterior process; not all regions of the talus are vascularly compromised.^{6,7} Talar neck fractures are the most common type as this area is the weakest part of the bone, representing about half of these injuries.^{6,7} The severity of talar neck fractures can be classified using the methods described by Hawkins (Table 4).³⁷ Avascular necrosis, osteoarthritis, and worse clinical scores increase with fracture severity.^{6,7} Talar body fractures account for about 20% of all talar fractures.⁷ The articular surfaces of the ankle and, occasionally, subtalar joints are generally involved, increasing the likelihood of developing osteoarthritis.⁷

Treatment of talar fractures

A nondisplaced talar neck fracture with adequate alignment may be treated conservatively with immobilization in a lower leg cast for 6 weeks, allowing for partial weightbearing only.^{6,7,35,38} Rehabilitation may be started once the fracture is radiographically healed.³⁸

Type II to IV talar neck fractures require immediate internal fixation.^{6,7} Talar body fractures are almost always managed operatively; however, complications are still common as the blood supply becomes disrupted.⁷ Screws may provide optimal compression except when there is extensive comminution; in such cases, a plate may also be required or an excision may be performed, though the latter procedure can alter hindfoot stability and biomechanics.^{7,35} K-wires may help achieve closed reduction and temporary fixation prior to screw insertion.³⁹ If avascular necrosis develops, another procedure may be required, such as talectomy, pantalar arthrodesis, tibiotalar arthrodesis, or tibio calcaneal arthrodesis (Figure 3).⁷

Conclusions

Current evidence suggests that optimal treatment for vascular compromised fractures is dependent on the degree of fracture displacement and comminution, and the patient's post-injury functional demands, age, and bone quality. Conservative efforts generally include casting or splinting with a period of




Figure 3. Overview and treatment of talar neck and body fractures.

immobilization. Surgery is indicated for substantially displaced fractures, patients who require higher functional demands and an earlier return to activity, or if complications occur following nonoperative treatment; however, operative intervention is typically performed for femoral neck fractures regardless of the amount of displacement. Internal fixation with screws is a common procedure among these injuries and can be used in combination with other implants, such as plating or k-wires, when needed. Severe fracture comminution, poor bone quality, or arthritis can contraindicate the use of screws and more invasive intervention will be required. Bone grafting is done in some cases to enhance vascularity. Salvage procedures exist for patients who develop severe complications, but these will permanently alter the anatomy of the injured area and should be considered a last resort.

Author Contributions

CV and MB contributed to the design of the study, and acquisition and interpretation of data. CV drafted the manuscript. MB, ES and BP reviewed and revised the manuscript prior to submission. All authors approved the final version of the manuscript.

ORCID iD

Christopher Vannabouathong  <https://orcid.org/0000-0002-9694-6364>

REFERENCES

- Allon R, Kramer A, Wollstein R. Intramedullary screw and Kirschner wire fixation for unstable scaphoid nonunion. *J Hand Microsurg.* 2016;8:150–154.
- Beltran MJ, Mitchell PM, Collinge CA. Posterior to anteriorly directed screws for management of talar neck fractures. *Foot Ankle Int.* 2016;37:1130–1136.
- Guss MS, Mitgang JT, Sapienza A. Scaphoid healing necessary for unrestricted activity: a biomechanical cadaver model. *J Hand Surg Am.* 2017;43:134–138.
- Hernigou P, Dubory A, Flouzat Lachaniette CH, Khaled I, Chevallier N, Rouard H. Stem cell therapy in early post-traumatic talus osteonecrosis. *Int Orthop.* 2018;42:2949–2956.
- Campefeldt P, Hedstrom M, Ekstrom W, Al-Ani AN. Good functional outcome but not regained health related quality of life in the majority of 20–69 years old patients with femoral neck fracture treated with internal fixation: a prospective 2-year follow-up study of 182 patients. *Injury.* 2017;48:2744–2753.
- Jordan RK, Bafna KR, Liu J, Ebraheim NA. Complications of talar neck fractures by Hawkins classification: a systematic review. *J Foot Ankle Surg.* 2017;56:817–821.
- Barnett JR, Ahmad M, Khan W, O' Gorman A. The diagnosis, management and complications associated with fractures of the talus. *Open Orthop J.* 2017;11:460–466.
- Hohenberger GM, Berzins U, Bakota B, Holweg P, Clement B, Grechenig S. Scaphoid screw placement under minimal radiation exposure. *Injury.* 2017;48:S47–S50.
- Suh N, Grewal R. Controversies and best practices for acute scaphoid fracture management. *J Hand Surg Eur Vol.* 2018;43:4–12.
- Dodds SD, Williams JB, Seiter M, Chen C. Lessons learned from volar plate fixation of scaphoid fracture nonunions. *J Hand Surg Eur Vol.* 2018;43:57–65.
- Dana C, Doursounian L, Nourissat G. Arthroscopic treatment of a fresh lunate bone fracture detaching the scapholunate ligament. *Chir Main.* 2010;29:114–117.
- Hsu AR, Hsu PA. Unusual case of isolated lunate fracture without ligamentous injury. *Orthopedics.* 2011;34:e785–e789.
- Saberi S, Arabzadeh A, Farhoud AR. Lunate osteochondral fracture treated by excision: a case report and literature review. *Trauma Mon.* 2016;21:e22378.
- Gelberman RH, Gross MS. The vascularity of the wrist. Identification of arterial patterns at risk. *Clin Orthop Relat Res.* 1986;202:40–49.
- Teisen H, Hjarbaek J. Classification of fresh fractures of the lunate. *J Hand Surg Br.* 1988;13:458–462.
- Moatshe G, Godin JA, Chahla J, et al. Clinical and radiologic outcomes after scaphoid fracture: injury and treatment patterns in national football league combine athletes between 2009 and 2014. *Arthroscopy.* 2017;33:2154–2158.
- Dedeoglu SS, Imren Y, Cabuk H, Tekin AC, Ture YC, Gurbuz H. Results of percutaneous fixation and distal radius core decompression in scaphoid waist non-unions treated without grafting. *Hand Surg Rehabil.* 2017;37:43–47.
- Patel S, Tiedeken N, Qvick L, Debski RE, Kaufmann R, Fowler JR. Interfragmentary compression forces vary based on scaphoid bone screw type and fracture location [published online ahead of print December 1, 2017]. *Hand (N Y).* doi:10.1177/1558944717745663.
- Luchetti TJ, Rao AJ, Fernandez JJ, Cohen MS, Wysocki RW. Fixation of proximal pole scaphoid nonunion with non-vascularized cancellous autograft. *J Hand Surg Eur Vol.* 2018;43:66–72.
- Shunmugam M, Phadnis J, Watts A, Bain GI. Lunate fractures and associated radiocarpal and midcarpal instabilities: a systematic review. *J Hand Surg Eur Vol.* 2018;43:84–92.
- De Jong L, Klem TMAL, Kuijper TM, Roukema GR. The minimally invasive anterolateral approach versus the traditional anterolateral approach (Watson-Jones) for hip hemiarthroplasty after a femoral neck fracture: an analysis of clinical outcomes. *Int Orthop.* 2018;42:1943–1948.
- Sonaje JC, Meena PK, Bansawal RC, Bobade SS. Comparison of functional outcome of bipolar hip arthroplasty and total hip replacement in displaced femoral neck fractures in elderly in a developing country: a 2-year prospective study. *Eur J Orthop Surg Traumatol.* 2017;28:493–498.
- Wang XJ, Zhang ZH, Li L, Su YX, Wei L. Quadratus femoris muscle pedicle bone flap transplantation in the treatment of femoral neck fracture for Chinese young and middle-aged patients: a systematic review and meta-analysis. *Chin J Traumatol.* 2017;20:347–351.
- Rosengren BE, Björk J, Cooper C, Abrahamson B. Recent hip fracture trends in Sweden and Denmark with age-period-cohort effects. *Osteoporos Int.* 2017;28:139–149.
- Guimaraes JAM, Rocha LR, Noronha Rocha TH, et al. Vertical femoral neck fractures in young adults: a closed fixation strategy using a transverse cancellous lag screw. *Injury.* 2017;48:S10–S16.
- Garden RS. Low-angle fixation in fractures of the femoral neck. *J Bone Joint Surg Br.* 1961;43-B:647–663.
- Shen M, Wang C, Chen H, Rui Y-F, Zhao S. An update on the Pauwels classification. *J Orthop Surg Res.* 2016;11:161–161.

28. Ju DG, Rajae SS, Mirocha J, Lin CA, Moon CN. Nationwide analysis of femoral neck fractures in elderly patients: a receding tide. *J Bone Joint Surg Am.* 2017; 99:1932–1940.
29. Mingli F, Huiliang S, Guanglei C, et al. A clinical study on arthroplasty for failed internal fixation of hip fractures and review of literature. *Pak J Med Sci.* 2017;33:798–803.
30. Matassi F, Carulli C, Munz G, Lualdi C, Civinini R, Innocenti M. Preliminary results of an early vs delayed timing of surgery in the management of proximal femur fragility fractures. *Clin Cases Miner Bone Metab.* 2015;12:11–17.
31. Sebestyén A, Boncz I, Sándor J, Nyárády J. Effect of surgical delay on early mortality in patients with femoral neck fracture. *Int Orthop.* 2008;32:375–379.
32. Simunovic N, Devereaux PJ, Bhandari M. Surgery for hip fractures: does surgical delay affect outcomes? *Indian J Orthop.* 2011;45:27–32.
33. Zhang LL, Zhang Y, Ma X, Liu Y. Multiple cannulated screws vs. dynamic hip screws for femoral neck fractures: a meta-analysis. *Orthopade.* 2017;46: 954–962.
34. Braun BJ, Pelz P, Veith NT, et al. Long-term pathological gait pattern changes after talus fractures—dynamic measurements with a new insole. *Int Orthop.* 2018;42:1075–1082.
35. Shank JR, Benirschke SK, Swords MP. Treatment of peripheral talus fractures. *Foot Ankle Clin.* 2017;22:181–192.
36. Young KW, Park YU, Kim JS, Cho HK, Choo HS, Park JH. Misdiagnosis of talar body or neck fractures as ankle sprains in low energy traumas. *Clin Orthop Surg.* 2016;8:303–309.
37. Hawkins LG. Fractures of the neck of the talus. *J Bone Joint Surg Am.* 1970; 52:991–1002.
38. Valderrabano V, Perren T, Ryf C, Rillmann P, Hintermann B. Snowboarder's talus fracture: treatment outcome of 20 cases after 3.5 years. *Am J Sports Med.* 2005;33:871–880.
39. Vesely R, Kelbl M, Kacian J, Hulka V, Kocis J, Kunovsky R. Talar neck fractures treated using percutaneous screw fixation. *Acta Chir Orthop Traumatol Cech.* 2017;84:120–124.