

New structure–activity relationships of chalcone inhibitors of breast cancer resistance protein: polyspecificity toward inhibition and critical substitutions against cytotoxicity

Luciana Pereira Rangel^{1,2,*}
Evelyn Winter^{1,3,*}
Charlotte Gauthier¹
Raphaël Terreux⁴
Louise D Chiaradia-Delatorre⁵
Alessandra Mascarello⁵
Ricardo J Nunes⁵
Rosendo A Yunes⁵
Tania B Creczynski-Pasa³
Sira Macalou¹
Doriane Lorendeau¹
Hélène Baubichon-Cortay¹
Antonio Ferreira-Pereira²
Attilio Di Pietro¹

¹Equipe Labellisée Ligue 2013, BMSSI UMR 5086 CNRS/Université Lyon 1, Institut de Biologie et Chimie des Protéines, Lyon, France; ²Department of General Microbiology, Institute of Microbiology, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil; ³Department of Pharmaceutical Sciences, PPGFAR, Federal University of Santa Catarina, Florianópolis, Santa Catarina, Brazil; ⁴Equipe BISI, BMSSI UMR 5086 CNRS/Université Lyon 1, Institut de Biologie et Chimie des Protéines, Lyon, France; ⁵Department of Chemistry, Federal University of Santa Catarina, Florianópolis, Santa Catarina, Brazil

*These authors contributed equally to this work

Correspondence: Attilio Di Pietro
Equipe Labellisée Ligue 2013, BMSSI UMR 5086 CNRS/Université Lyon 1, Institut de Biologie et Chimie des Protéines, 7 Passage du Vercors, 69367 Lyon cedex 07, France
Tel +33 4 7272 2629
Fax +33 4 7272 2604
Email a.dipietro@ibcp.fr

Abstract: Adenosine triphosphate-binding cassette subfamily G member 2 (ABCG2) plays a major role in cancer cell multidrug resistance, which contributes to low efficacy of chemotherapy. Chalcones were recently found to be potent and specific inhibitors, but unfortunately display a significant cytotoxicity. A cellular screening against ABCG2-mediated mitoxantrone efflux was performed here by flow cytometry on 54 chalcone derivatives from three different series with a wide panel of substituents. The identified leads, with submicromolar IC₅₀ (half maximal inhibitory concentration) values, showed that the previously identified 2'-OH-4',6'-dimethoxyphenyl, as A-ring, could be efficiently replaced by a 2'-naphthyl group, or a 3',4'-methylenedioxyphenyl with lower affinity. Such a structural variability indicates polyspecificity of the multidrug transporter for inhibitors. At least two methoxyl groups were necessary on B-ring for optimal inhibition, but substitution at positions 3, 4, and 5 induced cytotoxicity. The presence of a large *O*-benzyl substituent at position 4 and a 2'-naphthyl as A-ring markedly decreased the cytotoxicity, giving a high therapeutic ratio, which constitutes a critical requirement for future in-vivo assays in animal models.

Keywords: ABC transporters, BCRP/ABCG2, cancer chemotherapy, drug transport, inhibitory chalcone derivatives, multidrug resistance transporters.

Introduction

Cancer cells display a strong ability to acquire resistance to antitumor drugs, termed multidrug resistance (MDR), which constitutes a critical hurdle to cancer therapy. While several mechanisms can induce cellular resistance to chemotherapeutics, the energy-dependent drug efflux, mediated by adenosine triphosphate-binding cassette (ABC) transporters, is recognized to play a major role. Overexpression of multidrug ABC transporters in tumors alters anticancer drug efficacy by significantly reducing their intracellular accumulation. ABC subfamily G member 2 (ABCG2), also called ABCP for its abundance in placenta,¹ BCRP (breast cancer resistance protein),² or MXR (mitoxantrone resistance protein),³ was the most recently discovered of the three main multidrug ABC transporters overexpressed in cancer cells, after ABC subfamily C member 1 (ABCC1)/MDR protein 1 (MRP1)⁴ and ABCB1/MDR1/P-glycoprotein.⁵ ABCG2 was shown to confer resistance to a wide variety of anticancer agents,³ revealing a characteristic polyspecificity toward transport drug substrates. Its clinical relevance was demonstrated in both adult and childhood acute myeloid leukemia,^{6,7} and it was clearly involved in drug bioavailability at various protection barriers.^{8,9} Overcoming MDR

against anticancer agents, which is of importance for future clinical treatments, might be achieved through effective inhibitors of multidrug ABC transporters. ABCG2 inhibitors were therefore investigated in a number of studies, leading to identification of a few potent compounds.^{10,11} Different types of inhibitors were characterized: firstly, nonselective inhibitors already known as ABCB1 inhibitors and then considered as dual inhibitors, such as GF120918/elacridar¹² and XR9576/tariquidar.¹³ A few inhibitors appeared to be selective: fumitremorgin C, from *Aspergillus fumigatus*,¹⁴ which was however highly neurotoxic and of which synthetic derivatives, such as Ko143, were prepared.¹⁵ Derivatives of XR9576/tariquidar¹⁶ and GF120918/elacridar¹⁷ were also ABCG2-specific, but displayed cytotoxicity and a limited in-vivo bioavailability.¹⁸ A flavonoid-binding site was identified for chrysin and other flavones,^{19–21} while methoxy-*trans*-stilbenes appeared less toxic.²² Lower-affinity rotenoids,²³ acridone derivatives,¹⁷ and chromones²⁴ appeared to bind to the same site, as for a number of flavones, and benzo-pyran/furane derivatives, of which three-dimensional quantitative structure-activity relationship analyses allowed to establish a molecular model.^{25,26} Recent studies with chalcones and derivatives (Figure 1) reported a role of substituents, essentially OH, methoxyl (OMe), and Cl, on both A- and B-rings,^{27,28} but the results were not always easily interpretable since both rings were simultaneously substituted. In addition, we showed that OMe groups on phenyl B-ring, especially at position 4, were quite detrimental towards cytotoxicity.²⁹ Our present aim was therefore to use unsubstituted A-ring for studying the role of a large panel of B-ring substitutions, on both inhibition potency and cytotoxicity. We show here, with a number of chalcone derivatives, that large unsubstituted groups such as 2'-naphthyl or 3',4'-methylenedioxy-phenyl efficiently replaced the phenyl A-ring, supporting polyspecificity of ABCG2 for inhibitors. At least two OMe were required on the phenyl B-ring for optimal inhibition, although inducing a significant cytotoxicity, whereas most other substituents were not, or only weakly, efficient toward inhibition. Using both a large *O*-benzyl substituent at critical B-ring position 4 and a 2'-naphthyl group as A-ring additively decreased cytotoxicity and gave a high therapeutic ratio (TR), making chalcone 45 a good candidate for future in-vivo trials in mice models.

Materials and methods

Synthesis, purification, and physicochemical analyses of chalcones

All reagents used were of analytical grade, and purchased from Merck KGaA, Darmstadt, Germany or Sigma-Aldrich (St Louis, MO, USA), except




Figure 1 Core structure of generic chalcones (**A**) and substituted derivatives studied in our previous work (**B–K**).²⁹

Abbreviations: OMe, methoxyl; Me, methyl.

for 2,4,6-trimethoxy-acetophenone, xanthoxylin [2-hydroxy-4,6-dimethoxyaceto-phenone], 2-hydroxy-3-bromo-4,6-dimethoxyacetophenone, and 3-methoxy-4-(phenylmethoxy)-benzaldehyde, which were prepared according to Figure 2, as previously described.^{30–34} The chalcones were prepared by magnetic stirring of the acetophenone derivative (1.2 mmol), methanol (20 mL), KOH 50% w/v (5 mL), and the corresponding aldehyde (1.2 mmol), at room temperature for 24 hours. Distilled water and hydrochloric acid (10%) were added to the




Figure 2 Synthesis of chalcones.

Notes: a = KOH 50% w/v, CH₃OH, room temperature, 24 hours. In classical chalcones, both A and B are phenyl groups.

reaction for total precipitation of the products. The compounds were then obtained by vacuum filtration and later recrystallized from dichloromethane and hexane. All chalcones were previously described in the literature.^{30,32,35–50} The purified compounds were obtained in yields ranging from 40% to 99%. Melting points were determined with a Microquímica MGAPF-301 apparatus (Monte Mor, São Paulo, Brazil). Infrared spectra were recorded with an Abb Bomen FTLA 2000 spectrometer (Zurich, Switzerland) on KBr disks. Nuclear magnetic resonance (¹H and ¹³C) measurements were recorded on a Varian Inc. Oxford AS-400 (Palo Alto, CA, United States) (400 MHz) spectrometer, using tetramethylsilane as an internal standard. Elemental analysis was carried out with a CE Instruments CHNS EA 1110 (Hindley, United Kingdom). Percentages of C and H were in agreement with the product formula (within ± 0.4% of theoretical values to C).

Compounds for biological assays

Mitoxantrone, calcein-AM, and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide (MTT) were purchased from Sigma-Aldrich. All other reagents were commercial products with the highest available purity grade. All chalcone derivatives were dissolved in DMSO (dimethylsulfoxide) and then diluted in Dulbecco's Modified Eagle's Medium (DMEM) high glucose medium. The stock solution was stored at –20°C and warmed to 25°C just before use.

Cell cultures

The human fibroblast HEK293 cell lines transfected with either *ABCG2* (HEK293-*ABCG2*) or the empty vector (HEK293-*pcDNA3.1* cells) were obtained as previously described.²¹ Flp-In-293, an isogenic HEK293 cell line, was co-transfected using Lipofectamine™ (Invitrogen, Carlsbad, CA, USA) with either the empty vector *pcDNA/FRT* or the *pcDNA/FRT-ABCC1* vector, in combination with the Flp recombinase vector *pOG44*, resulting in targeted integration of the expression vector to the same locus in each cell. HEK293 cell lines transfected with *ABCB1* (P-glycoprotein) were kindly provided by Dr SE Bates (National Cancer Institute [NCI] at the National Institutes of Health [NIH], Bethesda, MD, USA). All cells were maintained in DMEM high glucose, supplemented with 10% fetal bovine serum, 1% penicillin/streptomycin, and supplemented in some cases with either 0.75 mg/mL G418 (for HEK293-*pcDNA3.1* and HEK293-*ABCG2*), or 2 mg/mL G418 (for HEK293-*ABCB1*), or 200 µg/mL hygromycin B (for Flp-In-293-*pcDNA5/FRT* and Flp-In-293-*ABCC1*).

ABCG2- and *ABCB1*-mediated mitoxantrone transport

HEK293 cells were seeded at a density of 1×10^5 cells/well into 24-well culture plates. After 48 hours incubation, the cells were exposed to 5 µM mitoxantrone (HEK293-*ABCG2* and HEK293-*MDR1* cells) for 30 minutes at 37°C, in the presence or absence of compounds at various concentrations. After cell washing with phosphate buffer saline, the cells were trypsinized. The intracellular drug fluorescence was monitored by flow cytometry with a FACS Calibur cytometer (BD Biosciences, San Jose, CA, USA). At least 10,000 events were collected, for which the maximal fluorescence (100%) was the difference between geometric mean fluorescence of cells incubated with 5 µM GF120918 and without inhibitor.²³ For *ABCB1*-mediated mitoxantrone transport, the cells transfected with the empty vector were used as a control.

MRP1-mediated calcein transport

HEK293 cells transfected with either *ABCC1* or the empty vector were exposed to 0.2 µM calcein-AM and analyzed by flow cytometry as described above. The maximal fluorescence (100%) was the difference between geometric mean fluorescence of control cells (HEK293-*pcDNA3.1*) and *MRP1*-transfected cells, incubated with substrate but without inhibitor.

Cytotoxicity assays

HEK293 cells transfected by either *pcDNA3.1-ABCG2* (resistant cells) or the empty vector (control sensitive cells) were seeded into 96-well culture plates at a 1×10^4 cells/well density. After overnight incubation, the cells were treated with various concentrations of compounds for 72 hours at 37°C under 5% CO₂. Cell viability was evaluated with an MTT colorimetric assay.⁵¹ Control experiments were performed with DMEM high glucose containing 0.1% of DMSO (v/v). The results from at least three replicates were expressed as percentage of viable cells versus control cells, taken as 100%. The curves were fitted with the Sigma Plot™ (Systat Software Inc, San Jose, CA, USA) software.
















Statistical analysis

Each experiment was performed at least in triplicate. The data are presented as mean ± standard deviation.

Results and discussion New structure–activity relationships (SARs) among inhibitory chalcones









A total of 54 chalcone derivatives were investigated here, belonging to three different series listed in Tables 1–3:

Table 1 Inhibition of ABCG2-mediated mitoxantrone efflux by chalcones 1–23

First series		
Compound	Ring B	% inhibition at 5 μ M
1		15.3 \pm 6.7
2		26.5 \pm 7.1
3		40.9 \pm 1.0
4		12.8 \pm 7.5
5		13.2 \pm 5.2
6		28.8 \pm 11.2
7		13.7 \pm 7.5
8		24.7 \pm 3.9
9		73.0 \pm 10.1
10		15.1 \pm 4.6
11		31.8 \pm 3.8
12		8.3 \pm 8.1
13		57.2 \pm 8.3
14		33.3 \pm 4.0
15		18.7 \pm 2.7

(Continued)

Table 1 (Continued)

Compound	Ring B	% inhibition at 5 μ M
16		25.4 \pm 2.8
17		59.1 \pm 11.2
18		29.8 \pm 3.0
19		30.2 \pm 6.4
20		18.2 \pm 5.2
21		85.7 \pm 6.7
22		79.5 \pm 5.7
23		73.1 \pm 4.5

Notes: The preparation of all compounds is described in the Materials and methods section. Each chalcone was assayed at 5 μ M for its ability to inhibit mitoxantrone efflux from ABCG2-transfected HEK293 cells by flow cytometry as described. The relative inhibition was calculated by reference to 5 μ M GF120918 producing 100% inhibition.
















Abbreviations: OBut, buthoxy; Me, methyl; OMe, methoxy.

(1) the first series (Table 1), with 23 derivatives, containing a 3',4'-methylenedioxy-phenyl unit as A-ring; (2) the second series (Table 2), with 22 derivatives, containing a 2'-naphthyl group as A-ring; and (3) the third series (Table 3), with nine derivatives, containing a 1-naphthyl group or other substituents as B-ring.

New structure–activity relationships were identified by comparing the inhibitory activity of each compound at 5 μ M against ABCG2-mediated mitoxantrone efflux, measured by flow cytometry and using 5 μ M GF120918 as a reference for complete inhibition (Tables 1–3). The following substitutions were identified to positively contribute to inhibition.








1. A 3',4'-methylenedioxy-phenyl unit, replacing A-ring in compounds of the first series (Table 1), was as efficient in 9 as the active 2'-OH-4',6'-diOMe-phenyl (A-ring) substituent previously identified in chalcone i (Figure 1).²⁹

Table 2 Inhibition by chalcones 24–45

Second series		
Compound	Ring B	% inhibition at 5 μ M
24		39.7 \pm 6.4
25		16.7 \pm 6.8
26		9.2 \pm 1.7
27		2.0 \pm 1.4
28		42.9 \pm 1.2
29		22.5 \pm 8.8
30		23.8 \pm 13.1
31		11.0 \pm 2.8
32		49.7 \pm 1.5
33		96.9 \pm 7.4
34		17.4 \pm 8.1
35		33.7 \pm 8.7
36		14.6 \pm 6.9
37		29.0 \pm 11.0
38		6.7 \pm 1.3


(Continued)

Table 2 (Continued)

Compound	Ring B	% inhibition at 5 μ M
39		24.1 \pm 6.9
40		1.2 \pm 2.9
41		31.8 \pm 5.3
42		7.5 \pm 1.9
43		23.2 \pm 7.7
44		42.1 \pm 8.8
45		81.5 \pm 4.4

Note: The conditions were the same as in Table 1.**Abbreviations:** OBut, butoxy; Me, methyl; OMe, methoxy.

- A 2'-naphthyl group replacing A-ring in compounds of the second series (Table 2) was also quite efficient in 33, as indolyl in chalcones j and k (Figure 1).²⁹ Such a polyspecificity toward A-ring chalcones is consistent with that observed for different classes of flavonoidic and flavonoid-like inhibitors, assumed to bind to the same site or to overlapping sites, such as hydrophobic flavones,^{19–21} rotenoids,²³ acridones,¹⁷ tariquidar derivatives,¹⁶ methoxy *trans*-stilbenes,²² and other chalcones.^{27–29} Polyspecificity of inhibitory sites is also consistent with the well known polyspecificity of ABCG2⁵² and other multidrug ABC transporters toward large panels of transport substrates. It is worthwhile mentioning that the gain-of-function R482T/G mutation in ABCG2, which was found to extend the transport substrate panel to rhodamine 123 and anthracyclins,⁵³ also allowed inhibitors such as GF120918 and imatinib to be transported.⁵²
- In both series, at least two OMe (in 9, 22, 23, and 33) were required for optimal inhibition when comparing with compounds with only one OMe (8, 19, 32) or without any OMe (1–7, 10–18, 20, 24–31, 34–43). The only exception was 44, which suggested a negative contribution of OMe at position 6 versus position 5 in 33. One of the two OMe could be efficiently replaced by *O*-benzyl at position 4 (in

Table 3 Inhibition by chalcones 46–54**Third series**

Compound	Ring A	Ring B	% inhibition at 5 μ M
46			88.2 \pm 5.6
47			73.0 \pm 2.6
48			49.2 \pm 7.2
49			57.0 \pm 6.7
50			82.3 \pm 11.1
51			21.3 \pm 9.0
52			60.9 \pm 6.7
53			63.7 \pm 12.7
54			55.3 \pm 8.9

Note: The conditions were the same as in Table 1.

Abbreviation: OMe, methoxyl.

21 and 45). Interestingly, an *O*-benzyl substituent was also shown to positively contribute to ABCG2 inhibition in chromones.²⁴

- In contrast, all other tested substituents on B-ring were poorly efficient, such as Cl (in 1–4, 17, and 26–29), Br (in 7 and 38), F (in 15, 16, 36, and 37), NO₂ (in 5, 6, 24, and 25), methyl (Me) (in 10, 18, 31, and 43), N(Me)₂ (in 11 and 34) and buthoxyl (in 14 and 35), similarly as using a naphthyl group to replace B-ring (in 12, 13, 40, and 41).
- Comparable positive contributions were observed from the third series of chalcones (Table 3): the requirement

for at least two OMe (in 46, 50, 52, 53, and 54), or one OMe and one *O*-benzyl (in 47), while the low efficiency of 51 was most likely related to the charged carboxyl preventing diffusion through the plasma membrane and therefore the compound uptake.

Selectivity for ABCG2 and cytotoxicity

The 54 chalcones described in this work were then investigated for their ability to inhibit mitoxantrone efflux by ABCB1/MDR1/P-glycoprotein, and calcein efflux by ABCC1/MRP1. As shown in Figure 3, all chalcones were




Figure 3 Inhibition by chalcones of P-glycoprotein (ABCB1)- and MRPI (ABCC1)-drug efflux activity. All chalcones from the three series were checked individually at 10 μM by flow cytometry for their ability to inhibit either ABCB1-mediated mitoxantrone efflux (grey bars) or ABCC1-mediated calcein efflux (black bars). The control of complete inhibition was performed with either 5 μM GF120919 or 30 μM MK571, respectively.

Abbreviations: ABC, adenosine triphosphate-binding cassette; CTRL, control.

essentially specific for ABCG2, since no significant inhibition was produced in most cases. A low inhibition of ABCB1 activity was however observed at 10 μM with two strong ABCG2 inhibitors, 9 and 33, both containing three OMe groups on the B-ring. Three other compounds slightly inhibited ABCC1-mediated drug efflux, by reference to MK571: 34 and 11 containing a 4-N(Me)₂ group on B-ring, and 48 containing a 2'-OH on A-ring. The N(Me)₂ substituent present in 11 and 34 might allow these chalcones to be transported, as observed for ABCB1 transport substrates,⁵⁴ which could explain lower specificity. In contrast, the three potent ABCG2 inhibitors containing a large 4-O-benzyl (ie,

21, 45, and 47) were highly selective and did not inhibit ABCB1 nor ABCC1.

The most potent compounds, inhibiting by more than 70% at 5 μM , were further analyzed for both their IC₅₀ (half maximal inhibitory concentration) values by flow cytometry (Table 4), and cytotoxicity by an MTT cell survival test (Figure 4A and B). The different compounds greatly varied in cytotoxicity, with 9, containing a 3',4'-methylenedioxy-phenyl as A-ring and a 3,4,5-triOMe substituted B-ring, being the most cytotoxic with an IG₅₀ (concentration producing 50% growth inhibition) value of 4.4 μM , giving a low TR (TR = IG₅₀/IC₅₀) of 7, similarly to 22 with a 2,5-diOMe substitution (TR = 8).

Table 4 TR values of potent chalcone inhibitors

Compound	A-ring substituents	B-ring substituents	IG ₅₀ (μM)	IC ₅₀ (μM)	TR
9	methylenedioxy-phenyl	3,4,5-triOMe-phenyl	4.4 \pm 0.5	0.64 \pm 0.16	7
22	methylenedioxy-phenyl	2,5-diOMe-phenyl	8.1 \pm 1.3	1.07 \pm 0.20	8
46	2',4'-diOMe-phenyl	methylenedioxy-phenyl	11.7 \pm 1.8	0.64 \pm 0.12	18
23	methylenedioxy-phenyl	3,4-diOMe-phenyl	12.9 \pm 1.6	0.54 \pm 0.06	24
21	methylenedioxy-phenyl	3-OMe,4-OBz-phenyl	17.5 \pm 1.2	0.25 \pm 0.12	70
47	4'-Br-phenyl	3-OMe,4-OBz-phenyl	28.7 \pm 5.4	0.29 \pm 0.04	99
50	2'-OH, 4',6'-diOMe-phenyl	1-naphthyl	20.6 \pm 0.8	0.22 \pm 0.07	94
33	2-naphthyl	2,4,5-triOMe-phenyl	34.9 \pm 2.5	0.23 \pm 0.06	152
45	2-naphthyl	3-OMe,4-OBz-phenyl	>100	0.23 \pm 0.07	>435

Notes: The cytotoxicity IG₅₀ values were evaluated from Figure 4A and B. The IC₅₀ values were determined with increasing inhibitor concentrations up to 10 μM by flow cytometry as in Tables 1–3.

Abbreviations: IC₅₀, half maximal inhibitory concentration; IG₅₀, concentration producing 50% growth inhibition; OBz, O-benzyl; OMe, methoxyl; TR, therapeutic ratio.




Figure 4 Intrinsic cytotoxicity on HEK293 cells and sensitization of BCRP (ABCG2)-transfected cells to mitoxantrone. The survival upon 72-hour culture of control sensitive cells in the presence of increasing concentrations of the indicated chalcones was quantified by MTT assays, as described in the Materials and methods section. Compared intrinsic cytotoxicities are shown in (A) for 21 (black triangles), 22 (black circles), 23 (white circles), and 47 (white triangles), and in (B) for 9 (black circles), 33 (white triangles), 45 (black squares), 46 (white circles), and 50 (black triangles). The sensitization to mitoxantrone, in (C) of the growth of ABCG2-transfected resistant cells (black circles) by 1 μ M 50 (black triangles) reached a similar level as control sensitive cells (white circles).

Abbreviations: ABC, adenosine triphosphate-binding cassette; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide; HEK293, human embryonic kidney HEK293.

A rather high cytotoxicity was also observed for 23, with 3,4-diOMe substitution ($IG_{50} = 12.9 \mu\text{M}$, TR = 24), and 46, with a 3,4-methylenedioxy-phenyl instead of B-ring, which confirmed the cytotoxic role of OMe at positions 3, 4, and 5 of phenyl B-ring previously observed when A-ring was an OH- and OMe-substituted phenyl.²⁹ Therefore, the 3',4'-methylenedioxy-phenyl as A-ring behaved similarly to the above substituted phenyl, but with a 3–4-fold lower affinity. By contrast, the addition of *O*-benzyl at position 4 was positive in 21, versus 23, by increasing threefold the TR (70 versus 24), as due to both increased affinity for inhibition and decreased cytotoxicity (Figure 4A). The effect was even more marked in 47 with a 4'-Br-phenyl as A-ring (TR = 99). Quite interestingly, the presence of a naphthyl group not only increased inhibition affinity but also lowered the cytotoxic effect of OMe substituents, especially when present as A-ring in 33 ($IG_{50} = 34.9 \mu\text{M}$ and TR = 152) versus B-ring in 50 ($IG_{50} = 20.6 \mu\text{M}$ and TR = 94) (Figure 4B and Table 4). This naphthyl positive effect on cytotoxicity appeared additive to that induced by 4-*O*-benzyl in 45 ($IG_{50} > 100 \mu\text{M}$, TR > 435). Such decreases in cytotoxicity might be possibly due to the prevention of interactions with unknown cellular target(s) inducing cell death.

The potent inhibition of lead compounds against ABCG2-mediated mitoxantrone efflux in flow cytometry experiments was confirmed by their ability to chemosensitize cell growth of resistant cells ($IG_{50} = 82.8 \pm 21.7 \text{ nM}$) to the cytotoxic drug, as illustrated in Figure 4C for 50: a 1 μM noncytotoxic concentration of the chalcone efficiently chemosensitized cell growth to mitoxantrone, giving an IG_{50} value of $17.5 \pm 2.2 \text{ nM}$, similar to that of sensitive control cells ($14.5 \pm 3.6 \text{ nM}$). A comparable efficiency was observed with 45 ($IG_{50} = 16.4 \pm 3.1 \text{ nM}$), whereas 22 only produced

a partial sensitization ($26.7 \pm 11.4 \text{ nM}$) (data not shown) in agreement to its 4–5-fold lower affinity for inhibition (see Table 4). The fact that the same low concentrations of 50 and 45 chemosensitized cell growth in MTT assays as efficiently as they inhibited drug efflux in flow cytometry suggests that these chalcones were not mainly metabolized nor transported during the 72-hour timescale of the experiments. Chalcone 45, which is a potent and selective inhibitor with low toxicity, therefore constitutes a good candidate for future in-vivo assays in animal models.

Conclusion

Original data about ABCG2 inhibitory sites and molecular mechanism have been obtained in the present work, thanks to a wide variety of chalcones and derivatives. Firstly, the flavonoid-specific inhibitory site appeared to be polyspecific since it could accommodate diverse large substituents instead of the original phenyl A-ring, with 2'-naphthyl and 3',4'-methylenedioxy-phenyl behaving quite efficiently toward inhibition. Secondly, substitution at position 4 of the phenyl B-ring was quite critical toward both inhibition and cytotoxicity, with a major role played by an *O*-benzyl group. The present data of cytotoxicity obtained from in-vitro screening on culture cells constitute an interesting first step that requires to be further developed for the identified lead compounds, such as chalcone 45, toward in-vivo pharmacokinetics and pharmacodynamics in mice.

Acknowledgments

Drs RW Robey and SE Bates, NCI-NIH, Bethesda, MD, USA, are acknowledged for providing HEK-293 cells transfected by ABCB1 (MDR-19), and GlaxoSmithKline for GF120918. LPR and EW were recipient of mobility fellowships from the

Brazilian Federal Agency for the Support and Evaluation of Graduate Education (CAPES) (Process numbers 1842/08-0 and 8792127, respectively), and CG was recipient of a doctoral fellowship from the Ligue Nationale Contre le Cancer. Financial support was provided by the CNRS and Université Lyon 1 (UMR 5086), and grants from the Région Rhône-Alpes (CIBLE 2010), the Ligue Nationale Contre le Cancer (Equipe labellisée Ligue 2013) and International ANR-NIH (2010-INTB-1101) to ADP.

Disclosure

The authors declare no conflicts of interest in this work.

References

- Allikmets R, Schriml LM, Hutchinson A, Romano-Spica V, Dean M. A human placenta-specific ATP-binding cassette gene (ABCP) on chromosome 4q22 that is involved in multidrug resistance. *Cancer Res.* 1998;58(23):5337–5339.
- Doyle LA, Yang W, Abruzzo LV, et al. A multidrug resistance transporter from human MCF-7 breast cancer cells. *Proc Natl Acad Sci U S A.* 1998;95(26):15665–15670.
- Miyake K, Mickley L, Litman T, et al. Molecular cloning of cDNAs which are highly overexpressed in mitoxantrone-resistant cells: demonstration of homology to ABC transport genes. *Cancer Res.* 1999;59(1):8–13.
- Endicott JA, Ling V. The biochemistry of P-glycoprotein-mediated multidrug resistance. *Annu Rev Biochem.* 1989;58:137–171.
- Cole SPC, Bhardwaj G, Gerlach JH, et al. Overexpression of a transporter gene in a multidrug-resistant human lung cancer cell line. *Science.* 1992;258(5088):1650–1654.
- Steinbach D, Sell W, Voigt A, Hermann J, Zintl F, Sauerbray A. BCRP gene expression is associated with a poor response to remission induction therapy in childhood acute myeloid leukemia. *Leukemia.* 2002;16(8):1443–1447.
- Ho MM, Hogge DE, Ling V. MDR1 and BCRP1 expression in leukemic progenitors correlates with chemotherapy response in acute myeloid leukemia. *Exp Hematol.* 2008;36(4):433–442.
- Seamon JA, Rugg CA, Emanuel S, et al. Role of the ABCG2 drug transporter in the resistance and oral bioavailability of a potent cyclin-dependent kinase/Aurora kinase inhibitor. *Mol Cancer Ther.* 2006;5(10):2459–2467.
- Breedveld P, Beijnen JH, Schellens JHM. Use of P-glycoprotein and BCRP inhibitors to improve oral bioavailability and CNS penetration of anticancer drugs. *Trends Pharmacol Sci.* 2006;27(1):17–24.
- Ahmed-Belkacem A, Pozza A, Macalou S, Perez-Victoria JM, Boumendjel A, Di Pietro A. Inhibitors of cancer cell multidrug resistance mediated by breast cancer resistance protein (BCRP/ABCG2). *Anticancer Drugs.* 2006;17(3):239–244.
- Boumendjel A, Macalou S, Valdameri G, et al. Targeting the multidrug ABCG2 transporter with flavonoid inhibitors: in vitro optimization and in vivo validation. *Curr Med Chem.* 2011;18(22):3387–3401.
- de Bruin M, Miyake K, Litman T, Robey RW, Bates SE. Reversal of resistance by GF120918 in cell lines expressing the ABC half-transporter, MXR. *Cancer Lett.* 1999;146(2):117–126.
- Robey RW, Steadman K, Polgar O, et al. Pheophorbide is a specific probe for ABCG2 function and inhibition. *Cancer Res.* 2004;64(4):1242–1246.
- Rabindran SK, He H, Singh M, et al. Reversal of a novel multidrug resistance mechanism in human colon carcinoma cells by fumitremorgin C. *Cancer Res.* 1998;58(24):5850–5858.
- Allen JD, van Loevezijn A, Lakhai JM, et al. Potent and specific inhibition of the breast cancer resistance protein multidrug transporter in vitro and in mouse intestine by a novel analogue of fumitremorgin C. *Mol Cancer Ther.* 2002;1(6):417–425.
- Kühnle M, Egger M, Muller C, et al. Potent and selective inhibitors of breast cancer resistance protein (ABCG2) derived from the P-glycoprotein (ABCB1) modulator tariquidar. *J Med Chem.* 2009;52(4):1190–1197.
- Boumendjel A, Macalou S, Ahmed-Belkacem A, Blanc M, Di Pietro A. Acridone derivatives: design, synthesis, and inhibition of breast cancer resistance protein ABCG2. *Bioorg Med Chem.* 2007;15(8):2892–2897.
- Arnaud O, Boumendjel A, Gèze A, et al. The acridone derivative MBLI-87 sensitizes breast cancer resistance protein-expressing xenografts to irinotecan. *Eur J Cancer.* 2011;47(4):640–648.
- Zhang S, Yang X, Morris ME. Flavonoids are inhibitors of breast cancer resistance protein (ABCG2)-mediated transport. *Mol Pharmacol.* 2004;65(5):1208–1216.
- Zhang S, Yang X, Coburn RA, Morris ME. Structure activity relationships and quantitative structure activity relationships for the flavonoid-mediated inhibition of breast cancer resistance protein. *Biochem Pharmacol.* 2005;70(4):627–639.
- Ahmed-Belkacem A, Pozza A, Munoz-Martinez F, et al. Flavonoid structure-activity studies identify 6-prenylchrysin and tectochrysin as potent and specific inhibitors of breast cancer resistance protein ABCG2. *Cancer Res.* 2005;65(11):4852–4860.
- Valdameri G, Pereira Rangel L, Sapatafora C, et al. Methoxy stilbenes as potent, specific, untransported and noncytotoxic inhibitors of breast cancer resistance protein. *ACS Chem Biol.* 2012;7(2):322–330.
- Macalou S, Ahmed-Belkacem A, Borelli F, et al. Non-prenylated rotenoids, a new class of potent breast cancer resistance protein inhibitors. *J Med Chem.* 2007;50(8):1933–1938.
- Valdameri G, Genoux-Bastide E, Peres B, et al. Substituted chromones as highly-potent, nontoxic inhibitors, specific for the breast cancer resistance protein ABCG2. *J Med Chem.* 2012;55(2):966–970.
- Nicolle E, Boccard J, Guilet D, et al. Breast cancer resistance protein (BCRP/ABCG2): New inhibitors and QSAR studies by a 3D linear solvation-energy approach. *Eur J Pharm Sci.* 2009;38(1):39–46.
- Nicolle E, Boumendjel A, Macalou S, et al. QSAR analysis and molecular modeling of ABCG2-specific inhibitors. *Adv Drug Deliver Rev.* 2009;61(1):34–46.
- Han Y, Riwanto M, Go ML, Ee PLR. Modulation of breast cancer resistance protein (BCRP/ABCG2) by non-basic chalcone analogues. *Eur J Pharm Sci.* 2008;35(1–2):30–41.
- Juvalé K, Pape VFS, Wiese M. Investigation of chalcones and benzochalcones as inhibitors of breast cancer resistance protein. *Bioorg Med Chem.* 2012;20(1):346–355.
- Valdameri G, Gauthier C, Terreux R, et al. Investigation of chalcones as selective inhibitors of the breast cancer resistance protein ABCG2: critical role of methoxy substituents in both inhibition and cytotoxicity. *J Med Chem.* 2012;55(7):3193–3200.
- Chiaradia LD, dos Santos R, Vitor CE, et al. Synthesis and pharmacological activity of chalcones derived from 2,4,6-trimethoxyacetophenone in RAW 264.7 cells stimulated by LPS: quantitative structure-activity relationships. *Bioorg Med Chem.* 2008;16(2):658–667.
- Gulati KC, Seth SR, Venkataraman K. Phloracetophenone. *Organic Synth.* 1943;2:522.
- Boeck P, Bandeira Falcão CAB, Leal PC, et al. Synthesis of chalcone analogues with increased antileishmanial activity. *Bioorg Med Chem.* 2006;14(5):1538–1545.
- Cechinel-Filho V, Vaz ZR, Zunino L, Calixto JB, Yunes RA. Synthesis of xanthoxylone derivatives with antinociceptive and antioedematogenic activities. *Eur J Med Chem.* 1996;31(10):833–839.
- Tsai S, Klinman JP. De Novo design and utilization of photolabile caged substrates as probes of hydrogen tunneling with horse liver alcohol dehydrogenase at sub-zero temperatures: a cautionary note. *Bioorg Chem.* 2003;31(2):172–190.

35. Insuasty B, Orozco F, Quiroga J, Abonia R, Noguera M, Cobo J. Microwave induced synthesis of novel 8,9-dihydro-7H-pyrimido[4,5-b][1,4]diazepines as potential antitumor agents. *Eur J Med Chem.* 2008;43(9):1955–1962.
36. Borchhardt DM, Mascarello A, Chiaradia LD, et al. Biochemical evaluation of a series of synthetic chalcone and hydrazone derivatives as novel inhibitors of cruzain from *Trypanosoma cruzi*. *J Braz Chem Soc.* 2010;21(1):142–150.
37. Damazio RG, Zanatta AP, Cazarolli LH, et al. Nitrochalcones: Potential in vivo insulin secretagogues. *Biochimie.* 2009;91(11–12):1493–1598.
38. Sadashivamurthy B, Basavaraju YB. New tetralone esters as intermediates for the synthesis of podophyllotoxin analogs. *Indian J Heterocycl Chem.* 2006;15(3):259–262.
39. Sathisha AD, Hemakumar KH, Basavaraju YB. Synthesis of new tetralone ester intermediates for podophyllotoxin analogues. *Indian J Heterocycl Chem.* 2007;17(1):15–18.
40. Parmar VS, Sharma S, Rathore JS, et al. Carbon-13 nuclear magnetic resonance studies on 1,3-diphenylprop-2-enones. *Magn Reson Chem.* 1990;28(5):470–474.
41. Wattanasin S, Murthy WS. An improved procedure for the preparation of chalcones and related enones. *Synthesis.* 1980;8:647–650.
42. Damazio RG, Zanatta AP, Cazarolli LH, et al. Antihyperglycemic activity of naphthylchalcones. *Eur J Med Chem.* 2010;45(4):1332–1337.
43. Thirunaryanan G. Effects of substituent on ¹H and ¹³C NMR chemical shifts in substituted styryl 2-naphthyl ketones. *Acta Cienc Indica Chem.* 2003;29(3):147–150.
44. Winter E, Chiaradia LD, de Cordova CAS, Nunes RJ, Yunes RA, Crezynski-Pasa TB. Naphthylchalcones induce apoptosis and caspase activation in a leukemia cell line: The relationship between mitochondrial damage, oxidative stress, and cell death. *Bioorg Med Chem.* 2010;18(22):8026–8034.
45. Robinson TP, Hubbard RB, Ehlers TJ, Arbiser JL, Goldsmith DJ, Bowen JP. Synthesis and biological evaluation of aromatic enones related to curcumin. *Bioorg Med Chem Lett.* 2005;13(12):4007–4013.
46. Patel HS, Patel VK, Dixit BC. Synthesis and antimicrobial activity of novel 2-amino-4-naphthyl-6-substituted phenylpyrimidine. *Orient J Chem.* 2001;17(3):142–150.
47. Chiaradia LD, Martins PGA, Cordeiro MNS, et al. Synthesis, biological evaluation, and molecular modeling of chalcone derivatives as potent inhibitors of *Mycobacterium tuberculosis* protein tyrosine phosphatases (PtpA and PtpB). *J Med Chem.* 2012;55(1):390–402.
48. Formento Navarini AL, Chiaradia LD, Mascarello A, et al. Hydroxychalcones induce apoptosis in B16-F10 melanoma cells via GSH and ATP depletion. *Eur J Med Chem.* 2009;44(4):1630–1637.
49. Chiaradia LD, Mascarello A, Purificação M, et al. Synthetic chalcones as efficient inhibitors of *Mycobacterium tuberculosis* protein tyrosine phosphatase PtpA. *Bioorg Med Chem Lett.* 2008;18(23):6227–6230.
50. Campos-Buzzi F, Pereira de Campos J, Tonini PP, et al. Antinociceptive effects of synthetic chalcones obtained from xanthoxyline. *Arch Pharm.* 2006;339(7):361–365.
51. Mosmann T. Rapid colorimetric assay for cellular growth and survival. Application to proliferation and cyto-toxicity assays. *J Immunol Methods.* 1983;65(1–2):55–63.
52. Telbisz A, Hegedus C, Ozvegy-Laczka C, et al. Antibody binding shift assay for rapid screening of drug interactions with the human ABCG2 multidrug transporter. *Eur J Pharm Sci.* 2012;45(1–2):101–109.
53. Sarkadi B, Ozvegy-Laczka C, Nemet K, Varadi A. ABCG2 – a transporter for all seasons. *FEBS Lett.* 2004;567(1):116–120.
54. Ford JM, Prozialeck WC, Hait WN. Structural features determining activity of phenothiazines and related drugs for inhibition of cell growth and reversal of multidrug resistance. *Mol Pharmacol.* 1989;35(1):105–115.

Drug Design, Development and Therapy

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which

Submit your manuscript here: <http://www.dovepress.com/drug-design-development-and-therapy-journal>

Dovepress

has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.