

Targeted SARS-CoV-2 treatment is associated with decreased mortality in immunocompromised patients with COVID-19

Emmanuel Lafont^{1,2}, H  l  ne Pere^{2,3}, David Lebeaux^{2,4}, Geoffrey Cheminet^{1,2}, Eric Thervet^{2,5},
Romain Guillemain⁶ and Adrien Flahault^{2,5*}

¹Service de M  decine Interne, H  pital Europ  en Georges Pompidou, AP-HP Centre, Universit   Paris Cit  , Paris, France; ²Universit   Paris Cit  , Paris, France; ³Service de Microbiologie, Unit   de Virologie, H  pital Europ  en Georges Pompidou, AP-HP Centre, Universit   Paris Cit  , Paris, France; ⁴Service de Microbiologie, Unit   Mobile d'Infectiologie, H  pital Europ  en Georges Pompidou, AP-HP Centre, Universit   Paris Cit  , Paris, France; ⁵Service de N  phrologie, H  pital Europ  en Georges Pompidou, AP-HP Centre, Universit   Paris Cit  , Paris, France; ⁶Service de Chirurgie Cardiaque, H  pital Europ  en Georges Pompidou, AP-HP Centre, Universit   Paris Cit  , Paris, France

*Corresponding author. E-mail: adrien.flahault@aphp.fr

Received 31 May 2022; accepted 6 July 2022

Background: Little is known about targeted (antiviral or monoclonal antibody) anti-SARS-CoV-2 treatment in immunocompromised patients with COVID-19.

Objectives: To assess the real-life efficacy and tolerance of targeted treatment of COVID-19 in immunocompromised patients.

Patients and methods: Single-centre retrospective case series of immunocompromised patients with COVID-19 between December 2021 and March 2022. We recorded all cases of COVID-19 among immunocompromised patients treatment between 20 December 2021 and 15 March 2022. Choice of treatment was left to the physician's decision, according to internal treatment protocol, treatment availability and circulating variants. Main outcome was death from COVID-19 after no treatment or targeted treatment.

Results: Sixty-seven immunocompromised patients [38 male; median (IQR) age, 53 (43–63) years], with a median (IQR) follow-up of 60 (47–80) days. Ten patients did not receive any targeted treatment. Targeted treatment consisted of IV curative remdesivir ($n=22$), sotrovimab ($n=16$), tixagevimab/cilgavimab ($n=13$) and casirivimab/imdevimab ($n=1$). Ten patients (15%) presented severe COVID-19 and 2 (3%) died from Omicron COVID-19. Comparing patients who received targeted anti-SARS-CoV-2 treatment and no prophylaxis, ($n=42$; 81%) with those who did not ($n=10$; 19%), death rate was significantly lower in treated patients [$n=0$ (0%) versus $n=2$ (20%); $P=0.034$]. No severe adverse events were reported among treated patients. Among 15 patients who received tixagevimab/cilgavimab as pre-exposure prophylaxis, 6 received an additional curative treatment and none died from COVID-19.

Conclusions: Our results suggest that targeted COVID-19 treatment, including direct antivirals or monoclonal antibodies, is safe and efficient and could be proposed in high-risk immunocompromised patients.

Introduction

Immunocompromised patients are at very high risk of developing severe coronavirus disease 2019 (COVID-19), with high morbidity and mortality rates.^{1–3} Moreover, low post-vaccinal immune responses have been reported among them, especially in solid organ transplant recipients.^{4–6} A number of targeted treatments have been developed to prevent and to cure COVID-19 in at-risk patients, and were recently made available in France. Monoclonal antibody combinations casirivimab/imdevimab and tixagevimab/cilgavimab have been used both as pre-exposure

prophylaxis (PREP) and as curative treatment, while sotrovimab has been used for early curative treatment only. The direct antiviral remdesivir has also shown its efficacy when prescribed early to prevent severe forms of COVID-19.^{7–9} However, clinical trials evaluating the efficacy of these preventive and curative treatments have been conducted among non-vaccinated, mostly non-immunocompromised, patients prior to the occurrence of the Omicron variant. While the occurrence of the Omicron variant has been associated with reduced severity of COVID-19,¹⁰ recent data show that immunocompromised patients remain at a high risk of morbidity and mortality from COVID-19, possibly due to

the lack of vaccine efficacy in this population^{11,12} and Omicron escape from casirivimab/imdevimab prophylaxis.¹³ Whether anti-SARS-CoV-2 preventative and curative targeted treatments are well tolerated and are efficient to decrease morbidity and mortality in immunocompromised patients is unknown. Thus, we aimed to evaluate the outcome of immunocompromised patients with COVID-19, who received or did not receive targeted treatments in a real-life setting within our institution.

Patients and methods

Study population

We performed a monocentric observational retrospective study in one university hospital (Hôpital Européen Georges Pompidou, Paris, France). We included all immunocompromised patients aged 18 years or over with laboratory-confirmed (PCR or antigenic test) COVID-19, treated and/or followed in three departments (Nephrology, Internal Medicine and Cardiac Surgery departments) from 20 December 2021 to 15 March 2022. Our aim was to propose targeted curative treatment to all severely immunocompromised patients with COVID-19. Early treatment was defined by treatment less than 10 days after symptom onset, while late treatment was given 10 or more days after symptom onset. Contraindications, availability of the different treatments and efficacy of monoclonal antibodies towards the circulating SARS-CoV-2 variants were also considered in the decision. Choice of treatment was left to the physician's decision, according to internal treatment protocol, patient characteristics, treatment availability and circulating variants. Specific curative treatment was provided as soon as possible in immunocompromised patients, when a positive diagnosis was confirmed, independently of the presence of symptoms or the reason for testing (symptoms, COVID-19 contact or systematic testing). The study was approved by the hospital's Internal Review Board (CERAPHP Centre, ref 2022-03-08).

Data collection and definitions

Baseline and follow-up information was collected from the medical file in patients regularly seen at our institution for causes other than COVID-19.

Patients for whom follow-up data could not be obtained were contacted by e-mail or by phone.

Data on PREP (history of previous COVID-19, vaccination, monoclonal antibodies), cause of immunosuppression, medical history, details on COVID-19 disease presentation, management and outcomes were collected. Low post-vaccinal immune response was defined as IgG anti-spike (anti-S) ≤ 264 binding antibody units (BAU)/mL at least 15 days after at least three doses of anti-SARS-CoV-2 vaccination.¹⁴ Severe COVID-19 was defined as pneumonia with oxygen requirement. Targeted anti-SARS-CoV-2 treatment included curative treatment by monoclonal antibodies and/or antivirals, and convalescent plasma, and did not include non-specific treatment including tocilizumab and dexamethasone.

Statistical analysis

We conducted descriptive analyses using median (IQR) for quantitative variables and count (percent) for qualitative variables. Comparisons were made using the Mann-Whitney test for quantitative variables and the Fisher exact test for qualitative variables.

Results

From 20 December 2021 to 15 March 2022, 67 immunocompromised patients were treated in our institution (hospital-based or outpatient care) for COVID-19, 70% of whom were solid organ transplant recipients. Variant determination was available for 49 (73%) patients. Among these, 46 (94%) were infected with the Omicron variant, including 34 (69%) BA.1, 3 (6%) BA.2, 9 (18%) undetermined Omicron variant and 3 (6%) Delta variant.

Clinical characteristics and patient management are described in Table S1 (available as [Supplementary data](#) at JAC Online) and in Figure 1, and clinical outcomes are provided in Table S2. Eighteen patients (27%) were hospitalized, 10 (15%) presented severe COVID-19 disease and 2 (3%) died. Death from COVID-19 was observed for two solid organ transplant recipients infected with the Omicron BA.1 variant. Both had a low




Figure 1. Study flowchart. Tixagevimab/cilgavimab prophylaxis was considered effective if administered at least 5 days before symptom onset. Early treatment was defined by targeted treatment administered less than 10 days after symptom onset while late treatment was given at least 10 days after symptom onset. One patient treated with casirivimab/imdevimab, infected by a subsequently identified Omicron variant, was considered non-treated, while one patient in the treated group was treated with casirivimab/imdevimab after identification of a Delta variant. This figure appears in colour in the online version of *JAC* and in black and white in the print version of *JAC*.

Table 1. Comparison of patients treated or not treated with targeted curative COVID-19 treatment

Characteristic	Targeted curative treatment (n=42; 81%)	No treatment (n=10; 19%)	P value
Age, years, median (IQR)	52 (45–63)	55 (36–60)	0.71
Male sex, n (%)	23 (55)	7 (70)	0.49
Cause of immunosuppression, n (%)			
Solid organ transplantation	29 (69)	6 (60)	0.71
Calcineurin inhibitor	23 (55)	6 (60)	1
Mycophenolate mofetil	21 (50)	5 (50)	1
Everolimus	11 (26)	1 (10)	0.42
Steroids	36 (86)	7 (70)	0.35
Belatacept	4 (10)	0 (0)	0.58
Rituximab	6 (14)	3 (30)	0.35
Other immunosuppressive agent	2 (5)	0 (0)	1
Cyclophosphamide	5 (12)	0 (0)	0.57
Active cancer with chemotherapy	3 (7)	0 (0)	1
Comorbidities			
Diabetes mellitus, n (%)	19 (45)	3 (30)	0.49
Chronic heart failure, n (%)	4 (10)	0 (0)	0.58
Chronic respiratory failure, n (%)	1 (2)	0 (0)	1
Obesity, n (%)	8 (19)	3 (30)	0.42
eGFR, mL/min/1.73 m ² , median (IQR)	55 (32–67)	55 (24–93)	0.69
End-stage renal disease with dialysis, n (%)	3 (7)	1 (10)	1
COVID-19 prophylaxis, n (%)			
History of COVID-19	1 (2)	0 (0)	1
At least 3 doses of vaccine	35 (83)	7 (70)	0.38
Anti-S >264 BAU/mL	1 (3)	2 (22)	0.086
Casirivimab/imdevimab PREP	14 (33)	4 (40)	0.72
Tixagevimab/cilgavimab PREP	0 (0)	0 (0)	—
COVID-19 disease			
Asymptomatic, n (%)	2 (5)	2 (20)	0.16
Fever, n (%)	26 (62)	5 (50)	0.5
Cough, n (%)	28 (67)	6 (60)	0.72
Dyspnoea, n (%)	15 (36)	3 (30)	1
Oxygen requirement, n (%)	5 (12)	3 (30)	0.17
Hospitalization, n (%)	12 (29)	3 (30)	1
Intensive care, n (%)	2 (5)	2 (20)	0.17
Death from COVID-19, n (%)	0 (0)	2 (20)	0.034
Follow-up duration, days, median (IQR)	63 (55–85)	59 (46–90)	0.62

Patients who received tocilizumab and other non-SARS-CoV-2-targeted treatments only are listed in the no-treatment column. Patients receiving PREP at least 5 days before symptom onset were excluded from this analysis.

post-vaccinal immune response (anti-S < 30 BAU/mL after three and four doses) and received no specific anti-SARS-CoV-2 treatment.

A total of 10 patients did not receive tixagevimab/cilgavimab prophylaxis and no targeted treatment either, and a total of 57 patients did receive some treatment: 15 of them received tixagevimab/cilgavimab prophylaxis, 6 of whom also received targeted treatment, and 42 with no prophylaxis received targeted treatment. Causes for the non-administration of specific anti-SARS-CoV-2 treatment are provided in Table S3. Figure 1 provides details on type and time of treatment.

None of the patients who received tixagevimab/cilgavimab prophylaxis or specific anti-SARS-CoV-2 treatment died from COVID-19. The death rate was significantly lower in patients

who received specific anti-SARS-CoV-2 treatment compared with those who did not [$n=0$ (0%) versus $n=2$ (20%); $P=0.034$], while patient characteristics were similar between the two groups (Table 1). No severe adverse events were noted among treated patients.

Discussion

Our study reports the COVID-19 outcomes in immunocompromised patients, most of them with low post-vaccinal immune responses, infected by Omicron variant and who received, or did not receive, a targeted anti-SARS-CoV-2 treatment. We show that treatment with specific anti-SARS-CoV-2 treatment was associated with a lower death rate and no reported adverse events.

None of the patients who received curative targeted treatment for COVID-19 died in our cohort. Patients were treated with the antiviral remdesivir as well as monoclonal antibodies, and choice of treatment was constrained by drug availability and kidney function (remdesivir being contraindicated in patients with estimated glomerular filtration rate (eGFR) below 30 mL/min/1.73 m²). Choice of monoclonal antibody was also determined by knowledge of *in vitro* resistance of the SARS-CoV-2 variant to monoclonal antibodies, requiring regular changes in COVID-19 treatment protocols. For example, Chavarot et al.¹¹ reported clinical efficacy of sotrovimab in kidney transplant recipients with Omicron COVID-19; however, the occurrence of the BA.2 variant, which escapes sotrovimab binding, currently limits the use of sotrovimab in clinical practice.¹⁵ In contrast, antivirals have shown persistent *in vitro* efficacy against COVID-19, independently of the SARS-CoV-2 variant,¹⁶ making these drugs a promising alternative for the treatment of COVID-19.

While a small number of patients with COVID-19 had received COVID-19 prophylaxis with tixagevimab/cilgavimab in our study, we found that none of them died from COVID-19. These results are in line with those reported by Bertrand et al.,¹⁷ who reported no deaths among 28 patients with Omicron COVID-19 who had received tixagevimab/cilgavimab prophylaxis.

In our study, 20% of patients without targeted treatment died. These results are similar to those reported by Chavarot et al.,¹¹ who found a high mortality rate (approximately 20% at 30 days) from Omicron COVID-19 among untreated kidney transplant recipients including 50% with low post-vaccinal immune response. Bertrand et al.¹⁷ also reported a high mortality rate from Omicron COVID-19 in kidney transplant recipients with no or low response to vaccination and no tixagevimab/cilgavimab prophylaxis (5/56; 9%, with no available information on whether or not some of these patients were treated with targeted treatment after infection). In contrast, overall mortality from COVID-19 in our severely immunocompromised cohort was low ($n=2$; 3%), with most patients receiving targeted treatment either prophylactically, curatively or both.

A limitation to our retrospective study is the lack of randomization, the relatively small sample size with a low number of events and the fact that choice of treatment was left to the physician's decision and treatment availability. However, to our knowledge, it is the largest published cohort of immunocompromised patients infected with SARS-CoV-2 during the Omicron era and treated with targeted treatments, using a real-life approach since all treatments were not available at a given time and treatment strategies needed to be adapted to circulating variants. The duration of follow-up is relatively prolonged, allowing for a reliable estimation of morbidity and mortality related to COVID-19.

In conclusion, we found that SARS-CoV-2 infection in immunocompromised patients treated with targeted anti-SARS-CoV-2 treatment was associated with no deaths and a good safety profile, despite low post-vaccinal immune responses. In contrast, patients who did not receive treatment were at high risk of death, similarly to previous reports.^{11,17} Hence, we propose that early, targeted COVID-19 treatment should be discussed in high-risk immunocompromised patients, and that a treatment strategy can include both antivirals and monoclonal antibodies according to

the patient profile, the current SARS-CoV-2 epidemiology and drug availability at a given time and place.

Funding

This study was carried out as part of our routine work.

Transparency declarations

None of the authors declare any competing interests linked with the present study.

Author contributions

All authors contributed to the manuscript. A.F. and E.L. were responsible for conception and design. A.F. and E.L. were responsible for data collection. A.F. was responsible for analysis. All authors were responsible for the interpretation of data. A.F. and E.L. wrote the first version of the manuscript. All authors critically revised and approved the final version of the manuscript.

Data availability

Data are available upon reasonable request.

Supplementary data

Tables S1 to S3 are available as [Supplementary data](#) at JAC Online.

References

- Gatti M, Rinaldi M, Bussini L et al. Clinical outcome in solid organ transplant recipients affected by COVID-19 compared to general population: a systematic review and meta-analysis. *Clin Microbiol Infect* 2022; S1198–743X(22)00116–1.
- Avouac J, Drumez E, Hachulla E et al. COVID-19 outcomes in patients with inflammatory rheumatic and musculoskeletal diseases treated with rituximab: a cohort study. *Lancet Rheumatol* 2021; **3**: e419–26.
- Belsky JA, Tullius BP, Lamb MG et al. COVID-19 in immunocompromised patients: a systematic review of cancer, hematopoietic cell and solid organ transplant patients. *J Infect* 2021; **82**: 329–38.
- Hadjadj J, Planas D, Ouedrani A et al. Immunogenicity of BNT162b2 vaccine against the Alpha and Delta variants in immunocompromised patients with systemic inflammatory diseases. *Ann Rheum Dis* 2022; **81**: 720–8.
- Furer V, Eviatar T, Zisman D et al. Immunogenicity and safety of the BNT162b2 mRNA COVID-19 vaccine in adult patients with autoimmune inflammatory rheumatic diseases and in the general population: a multi-centre study. *Ann Rheum Dis* 2021; **80**: 1330–8.
- Asderakis A, Khalid U, Koimtzis G et al. An analysis of serological response and infection outcomes following Oxford-AstraZeneca (AZD1222) and Pfizer-BioNTech (mRNA BNT162b2) SARS-CoV-2 vaccines in kidney and kidney pancreas transplants. *Transplantation* 2022; **106**: 1421–9.
- Gupta A, Gonzalez-Rojas Y, Juarez E et al. Early treatment for COVID-19 with SARS-CoV-2 neutralizing antibody sotrovimab. *N Engl J Med* 2021; **385**: 1941–50.
- Gottlieb RL, Vaca CE, Paredes R et al. Early remdesivir to prevent progression to severe COVID-19 in outpatients. *N Engl J Med* 2022; **386**: 305–15.
- Levin MJ, Ustianowski A, De Wit S et al. Intramuscular AZD7442 (tixagevimab-cilgavimab) for prevention of COVID-19. *N Engl J Med* 2022; **386**: 2188–200.

- 10** Ulloa AC, Buchan SA, Daneman N *et al.* Estimates of SARS-CoV-2 omicron variant severity in Ontario, Canada. *JAMA* 2022; **327**: 1286–8.
- 11** Chavarot N, Melenotte C, Amrouche L *et al.* Early treatment with sotrovimab monoclonal antibody in kidney transplant recipients with omicron infection. *Kidney Int* 2022; S0085-2538(22)00272-1.
- 12** Al Jurdi A, Gassen RB, Borges TJ *et al.* Suboptimal antibody response against SARS-CoV-2 omicron variant after third dose of mRNA vaccine in kidney transplant recipients. *Kidney Int* 2022; **101**: 1282–86.
- 13** Flahault A, Touchard J, Péré H *et al.* Breakthrough omicron COVID-19 infections in patients receiving the REGEN-Cov antibody combination. *Kidney Int* 2022; **101**:824–5.
- 14** Feng S, Phillips DJ, White T *et al.* Correlates of protection against symptomatic and asymptomatic SARS-CoV-2 infection. *Nat Med* 2021; **27**: 2032–40.
- 15** Bruel T, Hadjadj J, Maes P *et al.* Serum neutralization of SARS-CoV-2 omicron sublineages BA.1 and BA.2 in patients receiving monoclonal antibodies. *Nat Med* 2022; **28**: 1297–302.
- 16** Vangeel L, Chiu W, De Jonghe S *et al.* Remdesivir, Molnupiravir and Nirmatrelvir remain active against SARS-CoV-2 Omicron and other variants of concern. *Antiviral Res* 2022; **198**: 105252.
- 17** Bertrand D, Laurent C, Lemée V *et al.* Efficacy of anti SARS-CoV-2 monoclonal antibodies prophylaxis and vaccination on omicron COVID-19 in kidney transplant recipients. *Kidney Int* 2022; S0085-2538(22)00382-9.