

Original Research



OPEN ACCESS

Received: Oct 29, 2021

Revised: Dec 9, 2021

Accepted: Dec 13, 2021

Published online: Dec 23, 2021

Corresponding Author:

Jee-Seon Shim

Department of Preventive Medicine, Yonsei University College of Medicine, 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea.

Tel. +82-2-2228-0364

Fax. +82-2-392-8133

Email. shimjs@yuhs.ac

©2022 The Korean Nutrition Society and the Korean Society of Community Nutrition

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Jee-Seon Shim 

<https://orcid.org/0000-0002-8671-3153>

Funding

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2020R11A1A01064904).

Conflict of Interest

The author declare no potential conflicts of interests.

Ultra-processed foods and total sugars intake in Korea: evidence from the Korea National Health and Nutrition Examination Survey 2016–2018

Jee-Seon Shim [§]

Department of Preventive Medicine, Yonsei University College of Medicine, Seoul 03722, Korea

ABSTRACT

BACKGROUND/OBJECTIVES: Dietary sugars intake worldwide is stable or decreasing, but overall sugars intake remains above the recommended level. Some studies suggest that ultra-processed foods (UPFs) drive excessive sugars intake. However, UPF consumption in Korea and its association with sugars intake have not yet been studied. This study aimed to estimate the contribution of UPF consumption to total sugars intake and to investigate the association between UPF consumption and total sugars intake in Koreans.

SUBJECTS/METHODS: Data from the Korea National Health and Nutrition Examination Survey 2016–2018 were used, and included data on 21,075 participants aged 1+ years completed a 24-h recall. Food items reported in the 24-h recall were classified according to the NOVA system as UPFs, processed foods, processed culinary ingredients, or unprocessed or minimally processed foods.

RESULTS: The average daily energy intake was 1,996 kcal, and UPFs accounted for 26.2% of the total energy intake (% TE). The average total sugars intake was 63.1 g (13.0% TE), and 44.9% of total sugars intake came from UPFs. Among the entire population, 15.8% exceeded the recommended limit for total sugars within 20% TE, and excessive total sugars intake was more prevalent among females (19.5%) and children (21.1%). The prevalence of excessive total sugars intake showed a significantly increasing tendency across the quartiles of UPF energy contribution, ranging from 11.9% in the lowest quartile to 23.2% in the highest quartile. Even after adjustment for sociodemographic variables, UPF consumption was positively associated with the prevalence of excessive total sugars intake (P for trend < 0.001). This association was similar in subgroup analyses by sex and age.

CONCLUSIONS: This study suggests that UPF consumption may drive excessive intake of total sugars in the Korean diet. Our findings conclusively establish that restricting UPF consumption may be an efficient way to reduce sugars intake in Korean population.

Keywords: Food processing; dietary sugars; nutrition surveys; diet surveys

INTRODUCTION

Excessive sugars intake has been shown to be detrimental to human health [1-5]. Dietary Sugars, which refer to monosaccharides and disaccharides, are consumed in the form of

sugars naturally present in foods and also those added to foods during food processing by the manufacturer, food preparation by the cook, or at the table by the consumer [6]. There is no evidence that the consumption of natural sugars or foods high in natural sugars (e.g., milk and fruits) has adverse effects on health, but excessive consumption of added sugars has been reported to increase the risk of obesity, diabetes, cardiovascular diseases, and cancer [2-5]. Therefore, the World Health Organization strongly recommends reducing the intake of free sugars, including all sugars added to foods plus those naturally present in honey, syrup, and fruit juices, to less than 10% of total energy intake (% TE) [3]. The US recommends that added sugars intake should be reduced to less than 10% TE [7]. In contrast, the Dietary Reference Intakes for Koreans suggests that total sugars, including both natural and added sugars, should be within 10–20% TE [8]. Although the terminology used in guidelines varies and the cutoffs are slightly inconsistent, health authorities and governments are currently making efforts to reduce dietary sugars intake. A recent study that reviewed multiple national nutrition surveys showed that, although worldwide dietary sugars intake is decreasing or remains stable, it still remains above recommendation [9]. Although for Koreans the mean intake of total sugars is within the recommended range, the intake exceeds the cutoff in some subgroups (e.g., children and young adults), highlighting the need for intervention [10].

Ultra-processed foods have been demonstrated to drive excessive sugars intake [11-15]. These foods are industrial products manufactured from substances extracted from foods or derived from food constituents and synthesized from organic sources, with little or even no whole food [16]. They include sugar-sweetened drinks, snacks and cookies, chocolate, candies, breads and cakes, breakfast cereals, and diverse types of instant foods and ready-to-eat products. These foods are typically high in sugars, fat, and salt, energy density, and low in fiber, protein, minerals, and vitamins [16,17]. There is growing evidence of a significant positive, linear association between ultra-processed food consumption and added or free sugars intake [11-15], thereby suggesting that reducing ultra-processed food consumption could be an effective way of limiting excessive sugars intake. However, there is no evidence on the ultra-processed food consumption of Koreans and its association with dietary sugars intake. Therefore, this study aimed to estimate the contribution of ultra-processed food consumption to total sugars intake, and to investigate the association between ultra-processed food consumption and total sugars intake in Koreans.

SUBJECTS AND METHODS

Study population

This study used data from the Korea National Health and Nutrition Examination Survey (KNHANES), 2016–2018. KNHANES is a continuous, nationwide, cross-sectional survey that provides diverse information about the health and nutrition status of Koreans aged 1 year or older. KNHANES consists of health interviews, health examinations, and nutrition surveys. The survey protocols and procedures were approved by the Institutional Review Board (2018-01-03-P-A) of the Korea Disease Control and Prevention Agency (KDCA). Written informed consent was obtained from all participants.

In the KNHANES 2016–2018, a total of 31,639 individuals aged 1 year or older were sampled. Among them, 24,269 (76.6%) participated in at least one of the three component surveys. Of these, 21,271 completed a 24-h dietary recall of the nutrition survey. This study included 21,075 participants for which 24-h dietary recall data was available and were not pregnant or breastfed at the time of the survey.

Dietary assessment

Dietary intake was assessed via a 24-h recall. A trained dietitian surveyed details of foods and beverages consumed on the recall day using the multi-pass method, similar to the US Department of Agriculture (USDA) Automated Multi-Pass Method [18]. Individuals provided information on food description, quantity, and preparation method of each food item they consumed both within and outside the home, as well as additional details including time, place, accompanying meals, and so on. All food items were coded and edited. For a dish (i.e., multi-ingredient food such as Bibimbap), dietary information was disaggregated into individual ingredients by applying the participant's home recipe for the dish or the standard recipe developed for the KNHANES [19-21]. Dietary energy and nutrient intake were calculated by applying the food composition database to food intake data [19-21]. The food composition database used until 2015 did not include the total sugars content in food, but recently KDCA has established a database for total sugars content, permitting to evaluate dietary total sugars intake of Koreans since 2016 [22]. More details on dietary assessment methods and databases applied in nutrient intake calculation can be found elsewhere [17,19-21].

Food classification according to processing

For this study, we used the food information derived from the raw dataset released by the KDCA. In the 24-h dietary recall of KNHANES 2016–2018, a total of 3,894 food items were reported to be consumed either on their own or as ingredients in dishes. All reported items were classified into one of the four NOVA food groups and subgroups within each NOVA group. The NOVA food classification system classifies foods based on the nature, extent, and purpose of food processing [16,23], and comprises the following groups: group 1, unprocessed or minimally processed foods; group 2, processed culinary ingredients; group 3, processed foods; and group 4, ultra-processed foods. More details on the NOVA system and food classification are described elsewhere [16,17,23]. Briefly, unprocessed or minimally processed foods include raw foods directly obtained from natural or minimally altered foods without any additional ingredients (e.g., natural raw food, dried, ground, crushed, frozen, or vacuumed packed foods). Processed culinary ingredients are foods derived from group 1 foods or from nature via processing such as pressing, extracting, and refining (e.g., plant oil, sugar, animal fat, salt). These foods are usually used when cooking and seasoning group 1 foods. Processed foods are food products made by adding group 2 foods to group 1 foods (e.g., canned fish, fruit jam, and fermented food). The main purpose of this processing is to store group 1 foods for a long time and to improve their sensory qualities. These foods are edible by themselves and in combination with other foods. Finally, ultra-processed foods are products formulated mostly or entirely with industrial substances derived from foods and additives. These foods contain little or no group 1 foods and are generally created for the pursuit of convenience (i.e., easy consumption anytime and anywhere), high palatability, and profit. Ultra-processed foods include sugar-sweetened beverages (SSBs), soft drinks, packaged snacks, confectionaries, and diverse types of instant foods.

Sociodemographic variables

This study included sex, age, residential area, and income as sociodemographic variables. Sociodemographic variables were categorized as follows: sex (male and female), age (1–18 years, 19–49 years, 50–64 years, and 65 years or older), residential area (urban and rural), and income (lowest: quartile 1–highest: quartile 4).

Statistical analysis

We estimated dietary energy and total sugars intake for the entire population according to the NOVA food groups and subgroups. Energy intake from each food group was expressed as absolute (kcal/day) and relative intake (% TE). Total sugars intake was expressed as absolute intake (g/day), relative intake (% TE and % of total sugars intake), and the proportion of energy from total sugars among energy intake within each food group (%). We also compared the average intake of energy and total sugars according to the four NOVA food groups by sex and age groups and tested whether the intake differed by sex and age groups.

Next, we estimated total sugars intake (% TE) and the prevalence of excessive total sugars intake ($\geq 20\%$ TE from total sugars) across quartiles of dietary energy contribution of ultra-processed foods. This cutoff ($\geq 20\%$ TE from total sugars) is the upper limit recommended by the Korean Nutrition Society (KNS) [8]. Crude and adjusted logistic regression analyses were performed to estimate the prevalence ratio (PR) and 95% confidence interval (CI) for the association between quartiles of ultra-processed foods (% TE) and the prevalence of excessive total sugars intake. In the adjusted models, sex, age group, residence area, and income were included as confounding factors. Linear trends were tested using the median value of ultra-processed food consumption (% TE) for each quartile. We additionally assessed whether the total sugars intake (% TE) increased across quartiles of ultra-processed foods using linear regression analyses. All analyses were stratified by sex and age. For subgroup analyses, sex- and age-specific quartiles of ultra-processed food consumption were used.

According to the comprehensive action plan for reducing sugars consumption launched by the Korea Ministry of Food and Drug Safety (KMFDA) in 2016, sugars intake should be less than 10% TE from processed foods. Thus, we additionally analyzed the association between ultra-processed food consumption and total sugars intake based on the standards recommended by the KMFDA. We calculated the total sugars intake from both processed and ultra-processed foods (% TE) of NOVA food groups and estimated the prevalence of excessive total sugar intake, defined as $\geq 10\%$ TE from total sugars from both processed and ultra-processed foods. Total sugars intake and the prevalence of excessive total sugars intake from processed and ultra-processed foods were presented according to quartiles of energy contribution of ultra-processed foods. Crude and adjusted logistic regression analyses were performed to assess whether the prevalence differed across quartiles of ultra-processed foods. Adjustments were made for sex, age group, residence area, and income. Linear trends were tested using the median of ultra-processed food consumption (% TE) of each quartile. Linear regression analyses were used to assess whether the total sugars intake from processed and ultra-processed foods (% TE) increased across quartiles of ultra-processed foods. Subgroup analyses were performed in a similar manner.

Sample weights and survey designs were considered in all analyses. PROC SURVEYMEANS, PROC SURVEYLOGISTIC, and PROC SURVEYPREG procedures were used. Data analyses were performed using SAS software (version 9.4; SAS Institute, Cary, NC, USA), and results were regarded as significant at a P -value < 0.05 .

RESULTS

Survey-weighted proportion of the study population are depicted in **Table 1**.

Table 1. Distribution of study population

| Variable | No. | Weighted % |
|----------------------------|--------|------------|
| Sex | | |
| Male | 9,440 | 50.6 |
| Female | 11,635 | 49.4 |
| Age group (yrs) | | |
| Children (0–18) | 4,417 | 17.4 |
| Young adults (19–49) | 7,524 | 45.7 |
| Middle-aged adults (50–64) | 4,663 | 22.9 |
| Elderly (≥ 65) | 4,471 | 14.0 |
| Residence | | |
| Urban | 17,203 | 85.1 |
| Rural | 3,872 | 14.9 |
| Household income | | |
| Low (Q1) | 3,720 | 14.8 |
| Low-middle (Q2) | 5,298 | 24.5 |
| High-middle (Q3) | 5,905 | 29.6 |
| High (Q4) | 6,099 | 31.1 |

This analysis was performed using 1-day 24-h dietary recall data from the KNHANES 2016–2018 ($n = 21,075$). The complex KNHANES sampling design and the sample weights were accounted for in this analysis. KNHANES, Korea National Health and Nutrition Examination Survey.

Dietary intake of total energy and total sugars by food groups

The average daily energy intake among Koreans aged 1 year or older was 1,996 kcal, with 26.2% TE arising from ultra-processed foods (**Table 2**). The average total sugars intake was 63.1 g and total sugars accounted for 13.0% TE. Nearly half of the daily total sugars intake (44.9%) came from ultra-processed foods. The contribution of total sugars to energy intake in each food group was 9.4% TE in unprocessed or minimally processed foods, but that within ultra-processed foods was up to 25.0% TE.

Table 3 shows the dietary intake of total energy and total sugars according to the four NOVA food groups by sex and age groups. The consumption of ultra-processed foods (% TE) was higher in males and in the younger population (both $P < 0.001$). The absolute intake of total sugars was slightly higher in males, but the relative intake to total energy (% TE) was higher in females (both $P < 0.001$). Males consumed nearly half (47.9%) of daily total sugars from ultra-processed foods, whereas females presented a total sugars intake from ultra-processed foods lower than that of males (41.9%) and instead consumed more total sugars from unprocessed or minimally processed foods (all $P < 0.001$). By age group, both absolute and relative intake of total sugars were higher in the younger population (all $P < 0.001$).

Association between ultra-processed food consumption and total sugars intake

Among the entire population, 15.8% exceeded the upper limit for total sugars (20% TE) recommended by the KNS (**Table 4**). Excessive total sugars intake was more prevalent among females (19.5%) and children (21.1%). The prevalence of excessive total sugars intake showed a significant increasing tendency across the quartiles of ultra-processed food energy contribution, ranging from 11.9% in the lowest quartile to 23.2% in the highest quartile. After adjusting for sociodemographic variables, ultra-processed food consumption was positively associated with the prevalence of excessive total sugars intake (P for trend < 0.001). The adjusted PR of excessive total sugars intake was 2.40 (95% CI, 2.10–2.75) in the highest quartile of ultra-processed food energy contribution compared with the lowest. Similar associations were observed in all subgroup analyses stratified by sex and age.

Table 2. Dietary intake of total energy and total sugars according to NOVA food groups and subgroups

| Food groups | Energy intake | | Total sugars intake | | | Content of total sugars |
|---|---------------------|-----------------|---------------------|-----------------|-----------------|-------------------------|
| | Absolute (kcal/day) | Relative (% TE) | Absolute (g/day) | Relative (% TE) | Relative (% TS) | % E ¹⁾ |
| Unprocessed or minimally processed foods | 1,135.9 ± 6.9 | 59.2 ± 0.2 | 25.5 ± 0.3 | 5.5 ± 0.1 | 39.9 ± 0.3 | 9.4 ± 0.1 |
| Fruits | 82.1 ± 1.4 | 4.5 ± 0.1 | 15.4 ± 0.3 | 3.4 ± 0.1 | 20.9 ± 0.3 | 48.0 ± 0.4 |
| Vegetables ²⁾ | 57.6 ± 0.5 | 3.0 ± 0.0 | 4.9 ± 0.1 | 1.0 ± 0.0 | 10.3 ± 0.1 | 33.4 ± 0.2 |
| Milk and plain yogurts | 42.0 ± 0.9 | 2.4 ± 0.0 | 2.9 ± 0.1 | 0.7 ± 0.0 | 4.9 ± 0.1 | 7.8 ± 0.1 |
| Potatoes | 30.8 ± 1.2 | 1.6 ± 0.1 | 1.6 ± 0.1 | 0.3 ± 0.0 | 1.9 ± 0.1 | 2.7 ± 0.1 |
| Legumes, nuts and seeds | 34.3 ± 0.7 | 1.8 ± 0.0 | 0.4 ± 0.0 | 0.1 ± 0.0 | 0.8 ± 0.0 | 2.2 ± - |
| Grains | 613.2 ± 4.5 | 33.3 ± 0.2 | 0.3 ± 0.0 | 0.1 ± 0.0 | 0.8 ± 0.0 | 0.2 ± - |
| Eggs | 45.3 ± 0.8 | 2.3 ± 0.0 | 0.1 ± 0.0 | 0.0 ± 0.0 | 0.2 ± 0.0 | 0.3 ± - |
| Fish and sea foods | 39.9 ± 0.9 | 1.9 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.1 ± 0.0 | 0.3 ± - |
| Coffee and tea without sugar | 3.9 ± 0.1 | 0.2 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.2 ± - |
| Meats | 186.9 ± 3.7 | 8.2 ± 0.1 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Processed culinary ingredients | 85.2 ± 1.2 | 4.0 ± 0.0 | 3.7 ± 0.1 | 0.7 ± 0.0 | 6.3 ± 0.1 | 18.3 ± 0.2 |
| Sugars ³⁾ | 18.6 ± 0.3 | 0.9 ± 0.0 | 3.6 ± 0.1 | 0.7 ± 0.0 | 6.3 ± 0.1 | 59.5 ± 0.4 |
| Animal fats | 5.4 ± 0.4 | 0.2 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.1 ± 0.0 |
| Salt, vinegars, and others | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.7 ± 0.0 |
| Starch | 3.5 ± 0.2 | 0.1 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Plant oils | 57.6 ± 0.8 | 2.7 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Processed foods | 225.1 ± 3.5 | 10.6 ± 0.1 | 4.1 ± 0.1 | 0.8 ± 0.0 | 8.8 ± 0.1 | 15.6 ± 0.2 |
| Salted or pickled vegetables ⁴⁾ | 31.0 ± 0.3 | 1.7 ± 0.0 | 2.5 ± 0.0 | 0.5 ± 0.0 | 6.1 ± 0.1 | 28.6 ± 0.1 |
| Fruit jams, canned fruits | 4.8 ± 0.2 | 0.2 ± 0.0 | 0.9 ± 0.0 | 0.2 ± 0.0 | 1.3 ± 0.1 | 25.6 ± 0.4 |
| Noodles, starchy jello | 122.1 ± 2.8 | 5.6 ± 0.1 | 0.3 ± 0.0 | 0.1 ± 0.0 | 0.6 ± 0.0 | 0.6 ± 0.0 |
| Fermented alcoholic beverages | 36.2 ± 1.3 | 1.4 ± 0.0 | 0.2 ± 0.0 | 0.0 ± 0.0 | 0.5 ± 0.0 | 0.3 ± 0.0 |
| Canned or bottled fish and sea foods | 10.8 ± 0.3 | 0.6 ± 0.0 | 0.1 ± 0.0 | 0.0 ± 0.0 | 0.2 ± 0.0 | 1.2 ± 0.1 |
| Others (seasoned nuts, seeds, etc.) | 1.8 ± 0.2 | 0.1 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.4 ± 0.0 |
| Soybean curd | 18.5 ± 0.5 | 0.9 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.1 ± 0.0 | 0.2 ± 0.0 |
| Ultra-processed foods | 549.7 ± 5.7 | 26.2 ± 0.2 | 29.8 ± 0.4 | 6.0 ± 0.1 | 44.9 ± 0.3 | 25.0 ± 0.2 |
| Sugar-sweetened beverages ⁵⁾ | 46.3 ± 0.8 | 2.5 ± 0.0 | 5.5 ± 0.1 | 1.5 ± 0.0 | 10.0 ± 0.2 | 19.8 ± 0.3 |
| Soft drinks, fruit and vegetable drinks | 33.8 ± 0.9 | 1.6 ± 0.0 | 7.7 ± 0.2 | 1.2 ± 0.0 | 8.9 ± 0.2 | 26.1 ± 0.5 |
| Sweetened milk and its products | 42.8 ± 1.0 | 2.3 ± 0.1 | 4.7 ± 0.1 | 1.0 ± 0.0 | 6.7 ± 0.1 | 12.1 ± 0.2 |
| Cereals, breads, cakes, sandwiches, etc. | 140.7 ± 2.7 | 6.8 ± 0.1 | 4.7 ± 0.1 | 0.9 ± 0.0 | 6.7 ± 0.1 | 5.9 ± 0.1 |
| Traditional sauce | 31.1 ± 0.4 | 1.5 ± 0.0 | 2.2 ± 0.0 | 0.4 ± 0.0 | 4.4 ± 0.1 | 18.5 ± 0.2 |
| Confectionary | 12.9 ± 0.6 | 0.7 ± 0.0 | 1.6 ± 0.1 | 0.3 ± 0.0 | 2.1 ± 0.1 | 6.3 ± 0.2 |
| Others (instant sauce, condiments, etc.) | 31.7 ± 0.8 | 1.5 ± 0.0 | 1.4 ± 0.0 | 0.3 ± 0.0 | 2.5 ± 0.1 | 23.4 ± 0.3 |
| Cookies, chips, and snacks | 37.5 ± 1.2 | 1.9 ± 0.1 | 1.2 ± 0.1 | 0.2 ± 0.0 | 1.9 ± 0.1 | 2.5 ± 0.1 |
| Fish and meat processed foods | 52.6 ± 1.5 | 2.5 ± 0.1 | 0.4 ± 0.0 | 0.1 ± 0.0 | 0.9 ± 0.0 | 2.7 ± 0.1 |
| Instant noodles and dumplings | 37.0 ± 1.4 | 1.9 ± 0.1 | 0.3 ± 0.0 | 0.1 ± 0.0 | 0.7 ± 0.0 | 0.3 ± 0.0 |
| Instant cooked rice, soup, and other dishes | 13.1 ± 0.6 | 0.7 ± 0.0 | 0.1 ± 0.0 | 0.0 ± 0.0 | 0.1 ± 0.0 | 0.4 ± 0.0 |
| Distilled alcoholic beverages | 70.1 ± 2.5 | 2.4 ± 0.1 | 0.1 ± 0.0 | 0.0 ± 0.0 | 0.1 ± 0.0 | 0.6 ± 0.0 |
| Total | 1,996 ± 1 | 100 | 63.1 | 13.0 | 100 | 13.0 |

Data are presented as mean ± SE.

This analysis was performed using 1-day 24-h dietary recall data from the KNHANES 2016–2018 (n = 21,075). The complex KNHANES sampling design and the sample weights were accounted for in this analysis.

TE, total energy intake; TS, total sugars intake; KNHANES, Korea National Health and Nutrition Examination Survey.

¹⁾% of energy from total sugars intake within each food group.

²⁾Vegetables include mushrooms and seaweeds.

³⁾Sugars include sugar, honey, and molasses.

⁴⁾Salted or pickled vegetables include diverse types of kimchi.

⁵⁾Sugar-sweetened beverages include coffee or tea products with added sugar or milk, cocoa, or other sugar-sweetened beverages.

Association between ultra-processed food consumption and total sugars intake from both processed and ultra-processed foods

Table 5 shows the association between ultra-processed food consumption and excessive total sugars intake based on the recommendation of the KMFDA. Among the entire population, 22.1% consumed more than 10% TE from total sugars from both processed and ultra-processed foods. Excessive total sugars intake from both processed and ultra-processed foods was more prevalent in females (21.3%) and children (37.3%).

Table 3. Dietary intake of total energy and total sugars according to NOVA food groups by sex and age groups

| Food groups | Energy intake | | Total sugars intake | | | Content of total sugars |
|--|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| | Absolute (kcal/day) | Relative (% TE) | Absolute (g/day) | Relative (% TE) | Relative (% TS) | % E ¹⁾ |
| Sex | | | | | | |
| Male (n = 9,440) | 2,329.2 ± 15.8 | | 67.7 ± 0.7 | 11.9 ± 0.1 | | |
| Unprocessed or minimally processed foods | 1,291.2 ± 10.4 | 57.7 ± 0.3 | 24.6 ± 0.4 | 4.5 ± 0.1 | 36.1 ± 0.4 | 7.9 ± 0.1 |
| Processed culinary ingredients | 97.9 ± 1.7 | 4.0 ± 0.1 | 3.9 ± 0.1 | 0.7 ± 0.02 | 6.4 ± 0.1 | 17.6 ± 0.3 |
| Processed foods | 276.9 ± 5.2 | 11.2 ± 0.2 | 4.6 ± 0.1 | 0.8 ± 0.01 | 9.6 ± 0.2 | 15.0 ± 0.2 |
| Ultra-processed foods | 663.2 ± 8.7 | 27.1 ± 0.3 | 34.6 ± 0.5 | 6.0 ± 0.1 | 47.9 ± 0.4 | 24.8 ± 0.2 |
| Female (n = 11,635) | 1,654.7 ²⁾ ± 9.1 | | 58.3 ²⁾ ± 0.5 | 14.1 ²⁾ ± 0.1 | | |
| Unprocessed or minimally processed foods | 977.0 ²⁾ ± 6.3 | 60.8 ²⁾ ± 0.3 | 26.5 ²⁾ ± 0.4 | 6.5 ²⁾ ± 0.1 | 43.8 ²⁾ ± 0.4 | 10.9 ²⁾ ± 0.1 |
| Processed culinary ingredients | 72.1 ²⁾ ± 1.2 | 4.1 ²⁾ ± 0.1 | 3.4 ²⁾ ± 0.1 | 0.8 ²⁾ ± 0.02 | 6.3 ²⁾ ± 0.1 | 19.1 ²⁾ ± 0.3 |
| Processed foods | 172.0 ²⁾ ± 3.1 | 9.9 ²⁾ ± 0.2 | 3.5 ²⁾ ± 0.1 | 0.9 ± 0.02 | 8.1 ²⁾ ± 0.1 | 16.3 ²⁾ ± 0.2 |
| Ultra-processed foods | 433.6 ²⁾ ± 5.4 | 25.3 ²⁾ ± 0.3 | 24.9 ²⁾ ± 0.4 | 6.0 ± 0.1 | 41.9 ²⁾ ± 0.4 | 25.1 ± 0.2 |
| Age group (yrs) | | | | | | |
| Children (aged 1–18; n = 4,417) | 1,828.5 ± 18.7 | | 66.6 ± 0.9 | 14.9 ± 0.1 | | |
| Unprocessed or minimally processed foods | 1,007.2 ± 11.2 | 56.7 ± 0.4 | 22.0 ± 0.5 | 5.2 ± 0.1 | 35.5 ± 0.5 | 9.3 ± 0.2 |
| Processed culinary ingredients | 82.3 ± 2.2 | 4.2 ± 0.1 | 2.6 ± 0.1 | 0.5 ± 0.02 | 4.4 ± 0.1 | 14.7 ± 0.4 |
| Processed foods | 148.5 ± 5.1 | 7.6 ± 0.2 | 2.5 ± 0.1 | 0.5 ± 0.02 | 4.6 ± 0.2 | 14.5 ± 0.4 |
| Ultra-processed foods | 590.6 ± 10.3 | 31.5 ± 0.4 | 39.5 ± 0.8 | 8.7 ± 0.1 | 55.5 ± 0.6 | 28.2 ± 0.3 |
| Young adults (aged 19–49; n = 7,524) | 2,174.4 ± 17.3 | | 65.8 ± 0.7 | 12.6 ± 0.1 | | |
| Unprocessed or minimally processed foods | 1,121.4 ± 11.0 | 53.3 ± 0.3 | 22.1 ± 0.4 | 4.4 ± 0.1 | 34.0 ± 0.4 | 8.7 ± 0.2 |
| Processed culinary ingredients | 105.7 ± 1.9 | 4.7 ± 0.1 | 4.4 ± 0.1 | 0.8 ± 0.02 | 7.5 ± 0.2 | 18.7 ± 0.3 |
| Processed foods | 273.6 ± 5.6 | 12.1 ± 0.2 | 4.4 ± 0.1 | 0.9 ± 0.02 | 9.3 ± 0.2 | 14.5 ± 0.2 |
| Ultra-processed foods | 673.7 ± 8.7 | 29.9 ± 0.3 | 34.8 ± 0.6 | 6.5 ± 0.1 | 49.2 ± 0.4 | 23.8 ± 0.2 |
| Middle-aged adults (aged 50–64; n = 4,663) | 1,989.2 ± 16.4 | | 63.5 ± 0.8 | 13.2 [*] ± 0.1 | | |
| Unprocessed or minimally processed foods | 1,244.7 ± 10.9 | 64.8 ± 0.4 | 33.7 ± 0.6 | 7.2 ± 0.1 | 48.9 ± 0.5 | 11.0 ± 0.2 |
| Processed culinary ingredients | 69.6 ± 1.5 | 3.3 ± 0.1 | 3.3 ± 0.1 | 0.7 ± 0.02 | 5.7 ± 0.2 | 19.3 ± 0.4 |
| Processed foods | 230.3 ± 5.9 | 10.8 ± 0.2 | 4.9 ± 0.1 | 1.0 ± 0.02 | 10.3 ± 0.2 | 16.8 ± 0.3 |
| Ultra-processed foods | 444.6 ± 8.3 | 21.0 ± 0.3 | 21.5 ± 0.5 | 4.4 ± 0.1 | 35.1 ± 0.5 | 24.2 ± 0.3 |
| Elderly (aged 65+; n = 4,471) | 1,630.4 ²⁾ ± 14.5 | | 49.0 ²⁾ ± 0.8 | 11.9 ²⁾ ± 0.2 | | |
| Unprocessed or minimally processed foods | 1,165.5 ²⁾ ± 11.3 | 72.8 ²⁾ ± 0.3 | 27.5 ²⁾ ± 0.7 | 6.6 ²⁾ ± 0.1 | 50.2 ²⁾ ± 0.7 | 9.3 ²⁾ ± 0.2 |
| Processed culinary ingredients | 47.0 ²⁾ ± 1.4 | 2.7 ²⁾ ± 0.1 | 2.9 ²⁾ ± 0.2 | 0.7 ²⁾ ± 0.04 | 5.9 ²⁾ ± 0.2 | 20.1 ²⁾ ± 0.6 |
| Processed foods | 153.0 ²⁾ ± 4.5 | 8.7 ²⁾ ± 0.2 | 3.6 ²⁾ ± 0.1 | 0.9 ²⁾ ± 0.03 | 10.3 ²⁾ ± 0.3 | 18.7 ²⁾ ± 0.3 |
| Ultra-processed foods | 264.8 ²⁾ ± 5.7 | 15.8 ²⁾ ± 0.3 | 15.0 ²⁾ ± 0.3 | 3.7 ²⁾ ± 0.1 | 33.6 ²⁾ ± 0.6 | 26.2 ²⁾ ± 0.3 |

Data are presented as mean ± SE.

This analysis was performed using 1-day 24-h dietary recall data from the KNHANES 2016–2018 (n = 21,075). The complex KNHANES sampling design and the sample weights were accounted for in this analysis.

TE, total energy intake; TS, total sugars intake; KNHANES, Korea National Health and Nutrition Examination Survey.

¹⁾% of energy from total sugars intake within each food group.

²⁾P-value < 0.05 of differences in mean dietary intakes between sex and age groups.

The prevalence of excessive total sugar intake from both processed and ultra-processed foods showed a significantly increasing tendency across the quartiles of ultra-processed food energy contribution, ranging from 0.9% in the lowest quartile to 47.2% in the highest quartile. After adjusting for sociodemographic variables, ultra-processed food consumption still had a strong positive association with excessive total sugars intake from both processed and ultra-processed foods (*P* for trend < 0.001).

DISCUSSION

This study estimated the contribution of ultra-processed food consumption to total sugars intake and assessed whether ultra-processed food consumption is associated with total sugars intake using data from a recent national nutrition survey. In the Korean diet, the mean consumption of total sugars was 13.0% TE, and nearly half (44.9%) of total sugars intake came from ultra-processed foods. Among the entire population, 15.8% consumed more than

Ultra-processed foods and total sugars intake

Table 4. Associations between dietary contribution of UPF and total sugars intake

| Dietary contribution of UPF (% TE) | Total sugars intake (% TE) | | Excessive sugars intake ($\geq 20\%$ TE from total sugars) | | | | |
|--|----------------------------|------------------|---|------|-----|-----------------------|-----------------------|
| | Quartile | Mean (range) | Mean \pm SE | % | SE | PR (95% CI) | adjPR (95% CI) |
| Total (n = 21,075) | | 26.2 | 13.0 | 15.8 | | | |
| 1st | | 5.4 (0–10.6) | 10.9 \pm 0.1 | 11.9 | 0.5 | 1.00 | 1.00 |
| 2nd | | 15.8 (10.6–21.4) | 12.4 \pm 0.1 | 11.4 | 0.6 | 0.95 (0.82–1.09) | 0.97 (0.84–1.12) |
| 3rd | | 28.2 (21.4–35.9) | 13.5 \pm 0.1 | 15.5 | 0.6 | 1.36 (1.19–1.55) | 1.42 (1.24–1.62) |
| 4th | | 50.8 (35.9–100) | 14.8 \pm 0.1 | 23.2 | 0.7 | 2.23 (1.96–2.54) | 2.40 (2.10–2.75) |
| P-value | | | < 0.001 ¹⁾ | | | < 0.001 ²⁾ | < 0.001 ³⁾ |
| Sex | | | | | | | |
| Male (n = 9,440) | | 27.1 | 11.9 | 12.1 | | | |
| 1st | | 6.1 (0–11.7) | 9.5 \pm 0.2 | 6.4 | 0.6 | 1.00 | 1.00 |
| 2nd | | 17.1 (11.7–22.8) | 11.8 \pm 0.2 | 9.5 | 0.8 | 1.52 (1.20–1.94) | 1.50 (1.17–1.92) |
| 3rd | | 29.5 (22.8–36.9) | 12.7 \pm 0.2 | 13.1 | 0.8 | 2.20 (1.72–2.81) | 2.12 (1.65–2.72) |
| 4th | | 51.3 (36.9–100) | 13.4 \pm 0.2 | 18.3 | 0.9 | 3.27 (2.58–4.15) | 3.21 (2.52–4.07) |
| P-value | | | < 0.001 | | | < 0.001 | < 0.001 |
| Female (n = 11,635) | | 25.3 | 14.1 | 19.5 | | | |
| 1st | | 4.8 (0–9.8) | 12.1 \pm 0.2 | 16.7 | 0.8 | 1.00 | 1.00 |
| 2nd | | 14.8 (9.8–20.3) | 13.2 \pm 0.2 | 13.9 | 0.8 | 0.80 (0.67–0.96) | 0.81 (0.68–0.97) |
| 3rd | | 27.1 (20.3–34.8) | 14.4 \pm 0.1 | 17.7 | 0.8 | 1.07 (0.91–1.26) | 1.09 (0.93–1.29) |
| 4th | | 50.4 (34.8–100) | 16.5 \pm 0.2 | 28.8 | 1.0 | 2.02 (1.74–2.35) | 2.16 (1.84–2.53) |
| P-value | | | < 0.001 | | | < 0.001 | < 0.001 |
| Age group (yrs) | | | | | | | |
| Children (aged 1–18; n = 4,417) | | 31.5 | 14.9 | 21.1 | | | |
| 1st | | 10.9 (0–18.0) | 11.8 \pm 0.2 | 9.2 | 0.9 | 1.00 | 1.00 |
| 2nd | | 23.6 (18.0–28.9) | 13.5 \pm 0.2 | 12.6 | 1.2 | 1.42 (1.07–1.88) | 1.42 (1.07–1.89) |
| 3rd | | 34.9 (28.9–42.0) | 15.7 \pm 0.2 | 22.8 | 1.5 | 2.92 (2.23–3.83) | 2.92 (2.23–3.82) |
| 4th | | 54.4 (42.0–100) | 18.4 \pm 0.3 | 38.4 | 1.8 | 6.16 (4.80–7.90) | 6.24 (4.86–8.00) |
| P-value | | | < 0.001 | | | < 0.001 | < 0.001 |
| Young adults (aged 19–49; n = 7,524) | | 29.9 | 12.6 | 13.9 | | | |
| 1st | | 8.0 (0–14.6) | 10.2 \pm 0.2 | 9.1 | 0.8 | 1.00 | 1.00 |
| 2nd | | 20.2 (14.6–26.3) | 12.3 \pm 0.2 | 10.5 | 0.8 | 1.17 (0.92–1.78) | 1.19 (0.94–1.50) |
| 3rd | | 33.2 (26.3–41.2) | 13.1 \pm 0.2 | 14.6 | 0.9 | 1.70 (1.35–2.14) | 1.80 (1.43–2.67) |
| 4th | | 56.0 (41.2–100) | 14.4 \pm 0.2 | 20.8 | 1.1 | 2.61 (2.07–3.28) | 2.70 (2.15–3.40) |
| P-value | | | < 0.001 | | | < 0.001 | < 0.001 |
| Middle-aged adults (aged 50–64; n = 4,663) | | 21.0 | 13.2 | 17.1 | | | |
| 1st | | 4.1 (0–8.2) | 12.4 \pm 0.3 | 16.4 | 1.3 | 1.00 | 1.00 |
| 2nd | | 12.5 (8.2–17.2) | 13.2 \pm 0.2 | 15.1 | 1.3 | 0.91 (0.70–1.17) | 0.92 (0.71–1.20) |
| 3rd | | 22.9 (17.2–29.4) | 14.0 \pm 0.2 | 19.1 | 1.3 | 1.21 (0.95–1.53) | 1.28 (1.01–1.63) |
| 4th | | 43.0 (29.4–94.9) | 13.1 \pm 0.3 | 17.7 | 1.3 | 1.10 (0.85–1.41) | 1.20 (0.93–1.54) |
| P-value | | | < 0.001 | | | 0.208 | 0.043 |
| Elderly (aged 65+; n = 4,471) | | 15.8 | 11.9 | 13.1 | | | |
| 1st | | 2.5 (0–5.3) | 10.1 \pm 0.3 | 12.6 | 1.3 | 1.00 | 1.00 |
| 2nd | | 8.4 (5.3–11.7) | 11.0 \pm 0.2 | 8.5 | 1.0 | 0.65 (0.45–0.92) | 0.64 (0.44–0.91) |
| 3rd | | 16.2 (11.7–22.3) | 13.1 \pm 0.3 | 14.4 | 1.3 | 1.16 (0.85–1.60) | 1.24 (0.90–1.71) |
| 4th | | 36.3 (22.3–100) | 13.6 \pm 0.3 | 16.9 | 1.3 | 1.41 (1.05–1.19) | 1.43 (1.07–1.93) |
| P-value | | | < 0.001 | | | < 0.001 | < 0.001 |

This analysis was performed using 1-day 24-h dietary recall data from the KNHANES 2016–2018 (n = 21,075). The complex KNHANES sampling design and the sample weights were accounted for in this analysis.

UPF, ultra-processed foods; TE, total energy intake; PR, prevalence ratio; CI, confidence interval; adjPR, adjusted prevalence ratio; KNHANES, Korea National Health and Nutrition Examination Survey.

¹⁾P-values of differences in total sugars intake (% TE) across quartiles of UPF consumption were derived from linear regression analyses after adjustment for sex, age group, residence area, and income.

²⁾P for trend of crude PR of excessive total sugars intake across quartiles of UPF consumption.

³⁾P for trend of adjPR of excessive total sugars intake across quartiles of UPF consumption. Sex, age group, residence area, and income were mutually adjusted in the analyses of the entire population and each subgroup.

the recommended limit of 10% TE from total sugars. Ultra-processed food consumption had a significantly positive linear association with total sugars intake as well as excessive intake of total sugars. Such associations were similarly found regardless of sex and age groups.

Table 5. Associations between dietary contribution of UPF and total sugars intake from PF and UPF

| Dietary contribution of UPF (% TE) | Total sugars intake (% TE) | | Excessive sugars intake (> 10% TE from total sugars of PF and UPF) | | | |
|---|----------------------------|-----------------------|--|-----|-----------------------|-----------------------|
| | Mean (range) | Mean ± SE | % | SE | PR (95% CI) | adjPR (95% CI) |
| Total (n = 21,075) | 26.2 | 6.8 | 22.1 | | | |
| 1st | 5.4 (0–10.6) | 2.7 ± 0.04 | 0.9 | 0.2 | 1.0 | 1.0 |
| 2nd | 15.8 (10.6–21.4) | 5.5 ± 0.05 | 8.0 | 0.4 | 9.5 (6.6–13.7) | 8.7 (6.0–12.5) |
| 3rd | 28.2 (21.4–35.9) | 7.7 ± 0.08 | 27.4 | 0.8 | 41.3 (28.9–59.0) | 35.7 (24.9–51.1) |
| 4th | 50.8 (35.9–100) | 10.7 ± 0.1 | 47.2 | 0.9 | 97.6 (68.7–138.5) | 81.4 (57.2–115.9) |
| <i>P</i> -value | | < 0.001 ¹⁾ | | | < 0.001 ²⁾ | < 0.001 ³⁾ |
| Sex | | | | | | |
| Male (n = 9,440) | 27.1 | 6.8 | 21.9 | | | |
| 1st | 6.1 (0–11.7) | 2.9 ± 0.04 | 0.9 | 0.2 | 1.0 | 1.0 |
| 2nd | 17.1 (11.7–22.8) | 5.8 ± 0.08 | 9.9 | 0.7 | 12.5 (7.0–22.5) | 11.2 (6.3–20.2) |
| 3rd | 29.5 (22.8–36.9) | 7.8 ± 0.1 | 29.1 | 1.1 | 47.2 (26.6–83.6) | 39.8 (22.4–70.8) |
| 4th | 51.3 (36.9–100) | 10.0 ± 0.2 | 43.3 | 1.3 | 87.5 (50.2–152.5) | 71.1 (40.7–124.1) |
| <i>P</i> -value | | < 0.001 | | | < 0.001 | < 0.001 |
| Female (n = 11,635) | 25.3 | 6.8 | 22.3 | | | |
| 1st | 4.8 (0–9.8) | 2.4 ± 0.05 | 1.0 | 0.2 | 1.0 | 1.0 |
| 2nd | 14.8 (9.8–20.3) | 5.2 ± 0.07 | 7.0 | 0.5 | 7.7 (4.9–12.3) | 7.2 (4.5–11.5) |
| 3rd | 27.1 (20.3–34.8) | 7.5 ± 0.1 | 25.3 | 1.0 | 34.7 (22.1–54.5) | 30.5 (19.4–47.9) |
| 4th | 50.4 (34.8–100) | 11.4 ± 0.2 | 51.6 | 1.1 | 109.2 (69.5–171.7) | 93.0 (58.9–146.7) |
| <i>P</i> -value | | < 0.001 | | | < 0.001 | < 0.001 |
| Age group (yrs) | | | | | | |
| Children (aged 1–18; n = 4,417) | 31.5 | 9.2 | 37.3 | | | |
| 1st | 10.9 (0–18.0) | 4.0 ± 0.1 | 3.6 | 0.6 | 1.0 | 1.0 |
| 2nd | 23.6 (18.0–28.9) | 7.4 ± 0.1 | 25.5 | 1.6 | 9.2 (6.4–13.5) | 9.3 (6.4–13.4) |
| 3rd | 34.9 (28.9–42.0) | 10.3 ± 0.2 | 48.2 | 1.8 | 25.2 (17.3–36.8) | 25.3 (17.3–37.0) |
| 4th | 54.4 (42.0–100) | 14.5 ± 0.3 | 68.5 | 1.6 | 59.0 (40.7–85.6) | 59.4 (40.9–86.2) |
| <i>P</i> -value | | < 0.001 | | | < 0.001 | < 0.001 |
| Young adults (aged 19–49; n = 7,524) | 29.9 | 7.3 | 25.0 | | | |
| 1st | 8.0 (0–14.6) | 3.4 ± 0.1 | 1.9 | 0.4 | 1.0 | 1.0 |
| 2nd | 20.2 (14.6–26.3) | 6.5 ± 0.1 | 16.5 | 1.0 | 10.4 (6.5–16.5) | 10.6 (6.7–16.8) |
| 3rd | 33.2 (26.3–41.2) | 8.3 ± 0.1 | 32.5 | 1.2 | 25.2 (16.1–39.4) | 25.8 (16.5–40.3) |
| 4th | 56.0 (41.2–100) | 10.7 ± 0.2 | 47.2 | 1.5 | 46.8 (29.8–73.5) | 47.7 (30.3–75.1) |
| <i>P</i> -value | | < 0.001 | | | < 0.001 | < 0.001 |
| Middle-aged adults (aged 50–64; n = 4,663) | 21.0 | 5.4 | 12.5 | | | |
| 1st | 4.1 (0–8.2) | 2.4 ± 0.1 | 0.5 | 0.2 | 1.0 | 1.0 |
| 2nd | 12.5 (8.2–17.2) | 4.8 ± 0.1 | 3.3 | 0.6 | 6.44 (2.8–14.9) | 6.7 (2.9–15.5) |
| 3rd | 22.9 (17.2–29.4) | 6.4 ± 0.1 | 17.7 | 1.3 | 41.0 (18.6–90.6) | 44.2 (20.1–97.3) |
| 4th | 43.0 (29.4–94.9) | 7.7 ± 0.2 | 27.6 | 1.6 | 72.9 (33.2–160.0) | 80.8 (36.9–176.7) |
| <i>P</i> -value | | < 0.001 | | | < 0.001 | < 0.001 |
| Elderly (aged 65+; n = 4,471) | 15.8 | 4.6 | 9.0 | | | |
| 1st | 2.5 (0–5.3) | 1.7 ± 0.1 | 0.9 | 0.3 | 1.0 | 1.0 |
| 2nd | 8.4 (5.3–11.7) | 3.8 ± 0.1 | 1.2 | 0.4 | 1.3 (0.5–3.4) | 1.4 (0.6–3.4) |
| 3rd | 16.2 (11.7–22.3) | 5.6 ± 0.1 | 7.2 | 0.8 | 8.4 (4.0–17.4) | 8.7 (4.2–18.1) |
| 4th | 36.3 (22.3–100) | 7.5 ± 0.2 | 27.1 | 1.6 | 39.9 (9.9–79.9) | 42.3 (21.2–84.5) |
| <i>P</i> -value | | < 0.001 | | | < 0.001 | < 0.001 |

This analysis was performed using 1-day 24-h dietary recall data from the KNHANES 2016–2018 (n = 21,075). The complex KNHANES sampling design and the sample weights were accounted for in this analysis.

TE, total energy intake; PF, processed foods; UPF, ultra-processed foods; PR, prevalence ratio; CI, confidence interval; adjPR, adjusted prevalence ratio; KNHANES, Korea National Health and Nutrition Examination Survey.

¹⁾*P*-values of differences in total sugars intake (% TE) across quartiles of UPF consumption were derived from linear regression analyses after adjustment for sex, age group, residence area, and income.

²⁾*P* for trend of crude PR of excessive total sugars intake across quartiles of UPF consumption.

³⁾*P* for trend of adjPR of excessive total sugars intake across quartiles of UPF consumption. Sex, age group, residence area, and income were mutually adjusted in the analyses of the entire population and each subgroup.

Excessive dietary sugars intake is a worldwide problem, although sugars intake seems to be slightly decreasing or stable in some countries and regions [9]. A previous study using data from the National Health and Nutrition Examination Survey 2011–2018 reported the

average added sugars intake of the US population to be 67.8 g/day, corresponding to 12.7% TE [24]. The UK National Diet and Nutrition Survey 2008–2014 found that 12.4% of total energy was obtained by consuming free sugars [11]. In recent epidemiologic studies of eight Latin American countries, the mean total sugars intake was 99.4 g/day, which accounted for 20.1% TE, and 65.9% (65.5 g/day) came from added sugars, which accounted for 13.2% TE [25]. Although the total dietary sugar intake of Koreans estimated in our study was not as high as that of other countries, it was not lower than that of the KNHANES 2008–2011 (12.8% TE) [10]. In addition, both higher intake of dietary sugars and foods high in sugars among children and adolescents were observed [10,22]; thus highlighting the need for tightly monitoring sugars intake in these risk populations.

To date, the relationship between dietary sugars intake and human health has mainly been investigated at the level of SSBs, which are the main source of dietary sugars [26–30]. SSBs have poor nutritional profiles, less satiation, and excessive caloric value; thus, they have been targeted for public health intervention [31]. However, recently, various types of highly processed foods have infiltrated into the human diet and are widely consumed [32], which has aroused attention to the overall consumption of ultra-processed foods other than SSBs. Several studies have evaluated the levels of food processing and the excessive consumption of dietary sugars. In the US diet, ultra-processed foods contributed 57.9% of the total energy and comprised 89.7% of added sugars. Additionally, consumption of added sugars increased linearly across quintiles of ultra-processed food consumption from 7.5% TE in the lowest quintile to 19.5% TE in the highest [12]. Similar associations have been observed in other countries. In the UK, ultra-processed foods accounted for 56.8% of total energy and 64.7% of total free sugars, and Australians consumed 42.0% of total energy and 73.8% of total free sugars from ultra-processed foods [11,14]. Statistically significant positive associations were found between quintiles of ultra-processed food consumption and both the mean free sugars intake and the prevalence of excessive free sugars intake among almost all age groups [11,14]. These tendency was similar in countries with relatively low levels of ultra-processed food consumptions. In Spain and Chile, ultra-processed foods accounted for one-third of TE, but individuals in the highest quintile of ultra-processed food consumption were three to four times more likely to exceed the 10% TE upper limit for added sugars [13,15]. Our results are consistent with these results found in previous studies.

To our knowledge, this is the first study to assess the consumption of ultra-processed foods and examine its relationship with total sugars intake in Korea. Although both the levels of ultra-processed food consumption and dietary sugars intake were not as high as those in other countries [9,11–13,15,24], it seems reasonable that the consumption of these foods drives excessive sugars intake, suggesting that limiting the consumption of ultra-processed foods could be an effective way of reducing dietary sugars intake.

Our study had several strengths. We used data from nationwide representative samples of Koreans aged 1 year or older, which increases the generalizability of our findings. We analyzed dietary intake data obtained by the 24-h recall method, which assesses individuals' actual food consumption over the previous day. Thus, the actual dietary intake of food and nutrients could be assessed, unlike other studies using household expenditure surveys [15,33]. In our study, all food items reported in the 24-h recall were classified according to the NOVA system. This food classification system has been widely used to categorize foods according to the nature, extent, and purpose of food processing [16,23]. Moreover, three researchers independently categorized foods according to the concept and guidelines of

the NOVA classification system, and discrepancies in classification were resolved through discussion, as described in our previous studies [17].

However, our study also presents several limitations. First, the dietary data we used were data on single-day intake assessed by self-report. Thus, the ultra-processed food consumption and total sugars intake estimated in our study may not reflect individuals' usual intake [34]. In addition, ultra-processed food consumption and total sugars intake may be underestimated because of the tendency of under-reporting of unhealthy foods [35], although it is unlikely that this may affect the association between ultra-processed food consumption and total sugars intake. Finally, we estimated the dietary intake of total sugars, including both natural and added sugars, rather than added or free sugars, which is commonly considered more important to human health. Information on the content of added sugars in food is not yet available in Korea, thereby hindering the calculation of their contribution to total intake. Thus, we additionally estimated the intake of total sugars from specific food groups, such as processed and ultra-processed foods, and investigated the association between ultra-processed foods and dietary sugars intake from specific food groups and found strong relationships between them.

In conclusion, our findings indicate that the mean total sugars intake of Koreans accounted for 13.0% of TE, and ultra-processed foods contributed to one-fourth of the total energy and nearly half of the total sugars intake. The mean total sugars intake was below the limited level among all sex and age groups, but higher consumption of ultra-processed food was significantly associated with increased intake of total sugars and increased prevalence of excessive total sugars intake in all groups. This study demonstrates that restricting ultra-processed food consumption may be an efficient way to reduce sugars intake in the Korean diet.

REFERENCES

1. Prinz P. The role of dietary sugars in health: molecular composition or just calories? *Eur J Clin Nutr* 2019;73:1216-23.
[PUBMED](#) | [CROSSREF](#)
2. Debras C, Chazelas E, Srour B, Kesse-Guyot E, Julia C, Zelek L, Agaësse C, Druesne-Pecollo N, Galan P, Hercberg S, et al. Total and added sugar intakes, sugar types, and cancer risk: results from the prospective NutriNet-Santé cohort. *Am J Clin Nutr* 2020;112:1267-79.
[PUBMED](#) | [CROSSREF](#)
3. World Health Organization. *Guideline: Sugars Intake for Adults and Children*. Geneva: World Health Organization; 2015.
4. Singh GM, Micha R, Khatibzadeh S, Lim S, Ezzati M, Mozaffarian D. Global Burden of Diseases Nutrition and Chronic Diseases Expert Group (NutriCoDE). Estimated global, regional, and national disease burdens related to sugar-sweetened beverage consumption in 2010. *Circulation* 2015;132:639-66.
[PUBMED](#) | [CROSSREF](#)
5. Bray GA, Popkin BM. Dietary sugar and body weight: have we reached a crisis in the epidemic of obesity and diabetes?: health be damned! Pour on the sugar. *Diabetes Care* 2014;37:950-6.
[PUBMED](#) | [CROSSREF](#)
6. Sigman-Grant M, Morita J. Defining and interpreting intakes of sugars. *Am J Clin Nutr* 2003;78:815S-826S.
[PUBMED](#) | [CROSSREF](#)
7. US Government Printing Office. *Dietary Guidelines for Americans, 2020*. 8th ed. Washington, D.C.: US Government Printing Office; 2020.
8. Korean Nutrition Society, Ministry of Health and Welfare. *Dietary Reference Intakes for Koreans 2020*. Sejong: Ministry of Health and Welfare; 2020.

9. Wittekind A, Walton J. Worldwide trends in dietary sugars intake. *Nutr Res Rev* 2014;27:330-45.
[PUBMED](#) | [CROSSREF](#)
10. Lee HS, Kwon So, Yon M, Kim D, Lee JY, Nam J, Park SJ, Yeon JY, Lee SK, Lee HY, Kwon OS, Kim CI. Dietary total sugar intake of Koreans: based on the Korea National Health and Nutrition Examination Survey (KNHANES), 2008–2011. *J Nutr Health* 2014;47:268-76.
[CROSSREF](#)
11. Rauber F, Louzada MLDC, Martinez Steele E, Rezende LFM, Millett C, Monteiro CA, Levy RB. Ultra-processed foods and excessive free sugar intake in the UK: a nationally representative cross-sectional study. *BMJ Open* 2019;9:e027546.
[PUBMED](#) | [CROSSREF](#)
12. Martinez Steele E, Baraldi LG, Louzada ML, Moubarac JC, Mozaffarian D, Monteiro CA. Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study. *BMJ Open* 2016;6:e009892.
[PUBMED](#) | [CROSSREF](#)
13. Cediel G, Reyes M, da Costa Louzada ML, Martinez Steele E, Monteiro CA, Corvalán C, Uauy R. Ultra-processed foods and added sugars in the Chilean diet (2010). *Public Health Nutr* 2018;21:125-33.
[PUBMED](#) | [CROSSREF](#)
14. Machado PP, Steele EM, Louzada MLDC, Levy RB, Rangan A, Woods J, Gill T, Scrinis G, Monteiro CA. Ultra-processed food consumption drives excessive free sugar intake among all age groups in Australia. *Eur J Nutr* 2020;59:2783-92.
[PUBMED](#) | [CROSSREF](#)
15. Latasa P, Louzada MLDC, Martinez Steele E, Monteiro CA. Added sugars and ultra-processed foods in Spanish households (1990–2010). *Eur J Clin Nutr* 2018;72:1404-12.
[PUBMED](#) | [CROSSREF](#)
16. Monteiro CA, Cannon G, Levy RB, Moubarac JC, Louzada ML, Rauber F, Khandpur N, Cediel G, Neri D, Martinez-Steele E, et al. Ultra-processed foods: what they are and how to identify them. *Public Health Nutr* 2019;22:936-41.
[PUBMED](#) | [CROSSREF](#)
17. Shim JS, Shim SY, Cha HJ, Kim J, Kim HC. Association between ultra-processed food consumption and dietary intake and diet quality in Korean adults. *J Acad Nutr Diet*. Forthcoming 2021.
[PUBMED](#) | [CROSSREF](#)
18. Blanton CA, Moshfegh AJ, Baer DJ, Kretsch MJ. The USDA automated multiple-pass method accurately estimates group total energy and nutrient intake. *J Nutr* 2006;136:2594-9.
[PUBMED](#) | [CROSSREF](#)
19. Kweon S, Kim Y, Jang MJ, Kim Y, Kim K, Choi S, Chun C, Khang YH, Oh K. Data resource profile: the Korea National Health and Nutrition Examination Survey (KNHANES). *Int J Epidemiol* 2014;43:69-77.
[PUBMED](#) | [CROSSREF](#)
20. Ministry of Health and Welfare, Korea Centers for Disease Control and Prevention. Guidelines for Nutrition Survey: Korea National Health and Nutrition Examination Survey 6th (2013–2015). Cheongju: Korea Centers for Disease Control and Prevention; 2015.
21. Korea Centers for Disease Control and Prevention. 2016–2018 Guidebook for Using the Data from Korea National Health and Nutrition Examination Survey. Cheongju: Korea Centers for Disease Control and Prevention; 2020.
22. Yeon SY, Kweon SH, Oh KW. The daily dietary sugar intake in Korea, 2018. *Public Health Wkly Rep* 2020;13:359-63.
23. Monteiro CA, Cannon G, Moubarac JC, Levy RB, Louzada ML, Jaime PC. The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutr* 2018;21:5-17.
[PUBMED](#) | [CROSSREF](#)
24. Ricciuto L, Fulgoni VL 3rd, Gaine PC, Scott MO, DiFrancesco L. Sources of added sugars intake among the U.S. population: analysis by selected sociodemographic factors using the National Health and Nutrition Examination Survey 2011–18. *Front Nutr* 2021;8:687643.
[PUBMED](#) | [CROSSREF](#)
25. Fisberg M, Kovalskys I, Gómez G, Rigotti A, Sanabria LYC, García MCY, Torres RGP, Herrera-Cuenca M, Zimberg IZ, Koletzko B, et al. Total and added sugar intake: assessment in eight Latin American Countries. *Nutrients* 2018;10:389.
[PUBMED](#) | [CROSSREF](#)
26. Shim JS, Kang NH, Lee JS, Kim KN, Chung HK, Chung HR, Kim HJ, Ahn YS, Chang MJ. Socioeconomic burden of sugar-sweetened beverages consumption in Korea. *Nutr Res Pract* 2019;13:134-40.
[PUBMED](#) | [CROSSREF](#)

27. Xi B, Huang Y, Reilly KH, Li S, Zheng R, Barrio-Lopez MT, Martinez-Gonzalez MA, Zhou D. Sugar-sweetened beverages and risk of hypertension and CVD: a dose-response meta-analysis. *Br J Nutr* 2015;113:709-17.
[PUBMED](#) | [CROSSREF](#)
28. Wiener RC, Shen C, Findley PA, Sambamoorthi U, Tan X. The association between diabetes mellitus, sugar-sweetened beverages, and tooth loss in adults: evidence from 18 states. *J Am Dent Assoc* 2017;148:500-509.e4.
[PUBMED](#) | [CROSSREF](#)
29. Bernabé E, Vehkalahti MM, Sheiham A, Aromaa A, Suominen AL. Sugar-sweetened beverages and dental caries in adults: a 4-year prospective study. *J Dent* 2014;42:952-8.
[PUBMED](#) | [CROSSREF](#)
30. Anari R, Amani R, Veissi M. Sugar-sweetened beverages consumption is associated with abdominal obesity risk in diabetic patients. *Diabetes Metab Syndr* 2017;11 Suppl 2:S675-8.
[PUBMED](#) | [CROSSREF](#)
31. Malik VS, Hu FB. Fructose and cardiometabolic health: what the evidence from sugar-sweetened beverages tells us. *J Am Coll Cardiol* 2015;66:1615-24.
[PUBMED](#) | [CROSSREF](#)
32. Vandevijvere S, Jaacks LM, Monteiro CA, Moubarac JC, Girling-Butcher M, Lee AC, Pan A, Bentham J, Swinburn B. Global trends in ultraprocessed food and drink product sales and their association with adult body mass index trajectories. *Obes Rev* 2019;20 Suppl 2:10-9.
[PUBMED](#) | [CROSSREF](#)
33. Moubarac JC, Batal M, Martins AP, Claro R, Levy RB, Cannon G, Monteiro C. Processed and ultra-processed food products: consumption trends in Canada from 1938 to 2011. *Can J Diet Pract Res* 2014;75:15-21.
[PUBMED](#) | [CROSSREF](#)
34. Shim JS, Oh K, Kim HC. Dietary assessment methods in epidemiologic studies. *Epidemiol Health* 2014;36:e2014009.
[PUBMED](#) | [CROSSREF](#)
35. Willett WC. *Nutritional Epidemiology*. Oxford: Oxford University Press; 2012.