

Research Article



Regulatory Effects of Apatinib in Combination with Piperine on MDM-2 Gene Expression, Glutathione Peroxidase Activity and Nitric Oxide Level as Mechanisms of Cytotoxicity in Colorectal Cancer Cells

Mahshid Mohammadian^{1*}, Zakieh Rostamzadeh Khameneh², Soraya Emamgholizadeh Minaei³, Meysam Ebrahimifar⁴, Kosar Esgandari⁵

¹Department of Medical Laboratory Sciences, Urmia University of Medical Sciences, Urmia, Iran.

²Solid Tumor Research Center, Urmia University of Medical Sciences, Urmia, Iran.

³Department of Medical Physics and Imaging, Urmia University of Medical Sciences, Urmia, Iran.

⁴Department of Toxicology, Shahreza Azad University, Shahreza, Iran.

⁵Department of Medical Physics, Urmia University of Medical Sciences, Urmia, Iran.

Article info

Article History:

Received: 27 Oct. 2020

Revised: 5 Feb. 2021

Accepted: 31 Mar. 2021

published: 3 Apr. 2021

Keywords:

- Colorectal cancer
- Apoptosis
- HCT-116 cells
- Apatinib
- Piperine

Abstract

Purpose: Apatinib has been utilized in colon cancer therapies but its efficiency and molecular mechanism are not fully understood. Chemotherapy in combination with non-toxic compounds can be an effective treatment strategy for cancer. Consequently, this study was carried out to evaluate the effects of apatinib and piperine on colorectal cancer (CRC) cell line and their potential anti-cancerous mechanisms in vitro.

Methods: The effects of apatinib and piperine on HCT-116 CRC cells were detected by assessing cell viability using MTT assay. The potential cytotoxic mechanisms of apatinib and piperine were investigated by evaluating MDM-2 gene expression ratio using real-time PCR assay. Moreover, the glutathione peroxidase (GPX) activity and nitric oxide (NO) levels were assessed by colorimetric assays.

Results: The proliferation rate of CRC cells decreased by increasing the concentrations of piperine or apatinib. When HCT-116 cells were treated with different concentrations of apatinib in combination with piperine, the synergistic effects were observed (combination index < 1). In HCT-116 cells treated with apatinib and piperine at the concentrations of 0.5×IC₅₀ and 0.2×IC₅₀, the MDM-2 gene expression was downregulated and NO levels increased compared to the untreated control cells and related single treatments. In addition, GPX activity significantly decreased in combination treatment at 0.5×IC₅₀ concentration of both agents versus single treatments.

Conclusion: Apatinib in combination with piperine could significantly inhibit the growth of CRC cells. These cytotoxic effects were induced by regulation of MDM-2 gene expression and inhibition of antioxidant marker.

Introduction

Colorectal cancer (CRC) is the third-largest cancer world wide^{1,2} and is related to a high rate of mortality.¹⁻⁵ Recently, clinical studies have been interested in detecting novel integrating targeted treatments and combination chemotherapy regimens.² Combination chemotherapy comes into importance in CRC treatment but drug resistance is inevitable.³ In this regard, evaluating new drug combinations can improve the treatment outcomes.^{3,5} Numerous efforts have been conducted for increase understanding of the CRC heterogeneity and propose the most effective-tailored treatment to any affected patient.⁶ Oncogene mutations, the existence of multi-drug resistance, and the intolerable side effects limit the efficiency of treatment. So, new treatment protocols

for CRC treatment are instantly required.⁷ Angiogenesis is an important feature of cancer growth and metastasis. Vascular endothelial growth factor (VEGF) binding to vascular endothelial growth factor receptor (VEGFR) induces the vascular endothelial cells proliferation and angiogenesis. Anti-angiogenic drugs show anti-cancer efficiency through blocking the binding of VEGF and VEGFR. Apatinib mesylate as a micromolecular VEGFR-2 inhibitor that binds to VEGFR-2 and strongly inhibits it, exerts anti-cancer efficacy.⁸ Also, piperine that is a Piperidine alkaloid in black pepper, can inhibit cancer cell growth, but the mechanism of action of piperine is not fully understood.⁹ Piperine has been extensively described to inhibit colon cancer growth by G₁ arrest in the cell cycle, apoptosis induction, and antitumor activities.^{9,10} In

*Corresponding Author: Mahshid Mohammadian, Email; mohammadian.m@umsu.ac.ir

© 2022 The Author (s). This is an Open Access article distributed under the terms of the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

the development of anti-cancer agents, it is imperative to assess the treatment efficacy and also cancer response to chemotherapy.¹¹⁻¹⁴ One of the main purposes of cancer therapy is the induction of apoptosis.¹⁵ Apoptosis is identified as programmed cell death in the damaged and normal tissues.¹⁶ The apoptosis induction in cancerous cells is considered as the main objective of cancer treatment. Indeed, apoptosis is an important regulatory mechanism of normal cells. Definitely, apoptosis dysregulation can cause uncontrolled cell multiplication. This process is described by a series of definite morphological changes with biochemical features that contain intrinsic and extrinsic pathways through a diverse protein which plays a critical function.¹⁷

The MDM2 protein encoded through the mouse double minute 2 (*MDM2*) gene, is considered as the negative regulatory factor of the p53 protein and can preserve the p53 signaling pathway stability. MDM2 amplification has been assessed in numerous human cancers, comprising colon cancer.¹⁸ MDM-2 is an oncogene which its over-expression can exert transformation in cultured cells.¹⁹

Otherwise, Nitric oxide (NO) has been revealed to induce apoptosis by post-translational alterations and has an anti-cancer role.²⁰ On the other hand, as antioxidant enzymes have a critical function in protection of cells against oxidative stress, so, dysregulation of antioxidant enzymes activity, such as glutathione-peroxidase (GPX), are related to cancer.²¹ So, in this study, we investigated the effects of co-treatments of apatinib and piperine with evaluating the some related molecular mechanism in CRC cells.

Materials and Methods

Cell culture

HCT-116 cell line was obtained from the Pasteur Institute in Iran. The culture medium contained 10% fetal bovine serum, 1% penicillin and streptomycin, and Dulbecco's modified Eagle medium. CRC cells were seeded in 96 well cell culture plate and maintained at 37°C in a 5% carbon dioxide in the incubator.

Cell proliferation assay

HCT-116 cells were cultured in 96-well plates at a density of 1×10^4 /well. Following incubation at 37 °C and 5% CO₂ for 24 hours, CRC cells were exposed to increasing concentrations of apatinib (5, 10, 15, 20, 25, 75, 100 μM) and piperine (10, 20, 30, 40, 50, 100, 150, 200 μM). After 48 h of treatment time, a cell viability kit (Kia Zist, Iran) was used to detect cell proliferation. In this regards, 10 μL MTT reagent was added to each well, and the cells were incubated for 3 h at 37°C and 5% CO₂, the supernatants were discarded and a solubilizer was added to each well. The absorption rate was measured by ELISA Reader at 550 nm. The rate of viable cells was determined by measuring the absorbance. Each test in all treatments was repeated at least three times. The IC₅₀ values of each agent were

determined based on MTT assay results, dose-response assessment and with Compusyn software (ComboSyn, Inc., Paramus, NJ 07652, USA).

In all combined treatments, the concentrations of $0.50 \times IC_{50}$, $0.2 \times IC_{50}$, and $0.1 \times IC_{50}$ of both agents were utilized for cell viability assay and all other experimental analysis including GPX activity, MDM-2 gene expression, and NO level assays. The interaction between two therapeutic agents based on Chou²² method and Compusyn software (ComboSyn, Inc., Paramus, US) were evaluated, and the combination index (CI) determined. In this regard, the $CI < 1$, $CI > 1$, and $CI = 1$ display synergism, antagonism, and additive effect, respectively.

In addition, the Fraction affected (Fa) amount (indicating the cell fraction that affected by the combination treatment) and Dose reduction index (DRI) were evaluated.

Real-time polymerase chain reaction (PCR)

Total RNA was extracted from HCT-116 cells (untreated and treated with apatinib and piperine at different concentrations as mentioned above) using an RNA extraction kit based on the kit protocol (GeneAll, South Korea). The First-strand cDNAs were synthesized by GeneAll cDNA synthesis kit (GeneAll, South Korea) and utilized as templates to perform real-time PCR, following the manufacturer's instruction. The MDM2 gene expression was measured by real-time PCR using a Real Q Plus 2x Master Mix Green (Ampliqon, Denmark). Real time-PCR via cDNAs and specific primers were carried out in annealing temperature at 57°C for 30 seconds. Relative MDM2 expression was normalized to β-actin housekeeping gene and calculated by $2^{-\Delta\Delta Ct}$. Error bars in control and treated groups show the Standard deviation.

Nitric oxide assay

In order to assay the NO level in untreated and treated cells with various concentrations of apatinib and piperine in single and combined treatments, the cell supernatants were collected carefully. Assessments of NO level in various treatments were performed based on the kit protocol [ZellBio GmbH (Germany)]. The absorbance of each sample was read at 550 nm and NO levels were detected based on a standard curve.

Glutathione peroxidase activity assay

To evaluate the possible cytotoxic mechanism of apatinib and piperine, the GPX enzyme activity was assessed with the colorimetric method. In this regard, after various treatments as mentioned, cell culture supernatants of each sample were collected carefully, then the GPX activity was determined according to the kit protocol [ZellBio GmbH (Germany)]. In the next step, the absorbance at 412 nm was measured and GPX activity was calculated by this formula:

GPX activity (U/ml) = (OD control - OD sample) / (OD standard - OD blank) × 6000

Statistical analysis

Data were expressed as mean ± standard deviation. All data analysis was performed by one-way ANOVA, followed by the Tukey’s test utilizing GraphPad Prism version 4.0 and SPSS software, V. 10. P-value < 0.05 was considered as significant level.

Results and Discussion

The cytotoxic effects of apatinib and piperine in HCT-116 CRC cells were detected by MTT assay. The potential mechanisms were investigated by assessing the MDM-2 mRNA expression ratio in vitro using the real-time PCR assay. Moreover, the GPX activity and NO levels were evaluated by colorimetric assays.

In the first step, we evaluated the cytotoxic effects of monotherapies at different concentrations of apatinib or piperine after 48 hours treatment and dose-response assessment were performed. Results of cell viability assays in single therapies were presented in Figure 1. Both agents (apatinib or piperine) in single treatments decreased the cellular viability in a concentration-dependent pattern.

Treatment of HCT-116 cells with increasing concentrations of apatinib or piperine exhibited a reduction in the cellular viability in dose-dependent patterns. The IC₅₀ values for apatinib and piperine drugs were equal to 26 and 94 μM, respectively. All combined treatments lead to the synergistic interaction. Indeed, all combined cases showed significant decreased cell viability in comparison with single treatments in same concentrations.

Results of DRI, CI, and Fa for combined treatments at different concentrations were presented in Table 1.

As presented in Figure 2, after combined treatments with apatinib and piperine, the cell viability in terms of 0.5×IC₅₀, 0.2×IC₅₀, and 0.1×IC₅₀ concentrations were decreased significantly compared with untreated control cells (P < 0.05). In 0.5×IC₅₀ and 0.2×IC₅₀ concentrations in combination treatments, the cell viability were lower than that of corresponding monotherapies including apatinib and piperine in IC₅₀ concentrations (P < 0.05).

In order to study the molecular mechanism of apatinib and piperine effects in CRC cells treated with single and combined drugs, the MDM2 gene expression ratio was evaluated by Real-time PCR assay and normalized to β-Actin as a house-keeping gene. The mean fold changes for each treatment and control cells were calculated by 2^{-ΔΔCt}. The results of gene expression levels (Figure 3)




Figure 1. The results of cell viability using MTT assay for Apatinib and Piperine at various concentrations after 48 h treatment. Each value shows the mean ± standard deviation.

showed that the gene expression ratio was decreased in single and combined treatments in compared with control cells (P < 0.05). The MDM2 gene expression ratio in combined treatment groups (at concentrations of 0.5×IC₅₀ and 0.2×IC₅₀) decreased versus single treatments (P < 0.05).

In this presented study, the GPX activity was evaluated by the colorimetric assay. The results (Figure 4) showed that there was a significant decrease in GPX activity in the combined treatment group at 0.5×IC₅₀ concentration of both drugs in comparison with the control group and related single treatments (P < 0.05).

The results of the NO assay (Figure 4) showed that the level of NO in the combination groups at the concentrations of 0.5×IC₅₀ and 0.2×IC₅₀ were more than that of the

Table 1. Results of CI, Fa, and DRI in drug reaction at the various concentrations of Apatinib and Piperine in combined treatments which were measured by Compusyn software. Data showed as mean ± standard deviation.

Combinations	Fa	DRI for Apatinib	DRI for Piperine	CI
Apatinib-Piperine (0.5×IC ₅₀)	0.75 ± 0.043	4.52 ± 0.71	3.71 ± 0.45	0.49 ± 0.07 (synergistic effect)
Apatinib-Piperine (0.2×IC ₅₀)	0.63 ± 0.04	7.57 ± 0.99	6.83 ± 0.68	0.28 ± 0.03 (synergistic effect)
Apatinib-Piperine (0.1×IC ₅₀)	0.38 ± 0.03	7.13 ± 0.69	7.65 ± 0.57	0.27 ± 0.02 (synergistic effect)

Fa, fraction affected; DRI, dose reduction index; CI, combination index.




Figure 2. Cellular inhibitory effects of single and combined therapies with apatinib and piperine at different concentrations after 48 h treatment in HCT-116 CRC cells. Each value shows mean \pm standard deviation). * Significant differences compared to untreated control cells ($P < 0.05$).




Figure 3. The MDM2 gene expression ratio which was assessed by the real-time PCR method in treated and untreated cells with Apatinib and Piperine after 48 h. each column shows the mean fold change \pm standard deviation which was normalized to a housekeeping gene (β -actin) and calculated by $2^{-\Delta\Delta Ct}$. * Significant difference compared to untreated control cells ($P < 0.05$).

untreated control group ($P < 0.05$). In addition, there was an elevation in NO levels in combined groups ($0.5 \times IC_{50}$ and $0.2 \times IC_{50}$ concentrations) compared to monotherapies (apatinib and piperine in IC_{50} concentration).

Combination treatments involve the administration of conventional chemotherapeutic agents together with natural bioactives (typically from a plant). The anticancer

drug combination may be applied to cancer cell cultures.²³ Indeed, there are limitations to the effectiveness of several cancer therapies because of the systemic toxicity. Therefore, Chemotherapy with non-toxic compounds can be a strategy for decreasing cancer incidence. Numerous natural agents have showed chemotherapeutic potential.¹² In this regard; we evaluated the efficacy of apatinib as a chemotherapy drug combined with piperine as a natural agent in the CRC cell line. Our results demonstrated that synergistic effects were observed in the combination treatments of apatinib and piperine at concentrations lower than the IC_{50} values of each agent. Also, the combination treatments regulated the MDM2 gene expression levels and increased NO levels in cell culture, which these effects are related to inducing cytotoxicity. Moreover, combined therapy at the concentration of $0.5 \times IC_{50}$ decreased the GPX enzyme activity that indicates the efficacy of this combination treatment and induction of cytotoxicity.

The p53 is the main transcription factor regulating cellular pathways including apoptosis and cell cycle. It acts as a central defense mechanism toward cancer progression and is controlled by interaction with the MDM2 (oncoprotein). The inhibition of MDM2-p53 interaction displayed a striking treatment strategy for cancer therapy.²⁴

At the molecular level, piperine can affect numerous effector proteins involved in the apoptosis pathway and can stimulate extrinsic and intrinsic apoptosis process. Piperine repressed the cancer development and metastasis in a cancer model.²⁵ In our study, piperine could downregulate the MDM-2 gene expression as an oncogenic mediator.

In a similar study, piperine elevated the anti-proliferative




Figure 4. The levels of NO and GPX activity in HCT-116 cells treated with various concentrations of Apatinib and Piperine in single and combination treatments. Data were expressed as mean \pm standard deviation.

and cytotoxic effects of doxorubicin and paclitaxel in cell line,²⁶ which was parallel to our results.

Piperine has elevated the cytotoxicity of paclitaxel and doxorubicin in cell line. Moreover, piperine in combination with doxorubicin and paclitaxel induced P21 expression. These researchers recommended that the molecular mechanism has to be further assessed to recognize the definite function of piperine.²⁶ Likewise in another study, the isobologram and the CI of the combination of Paclitaxel and piperine showed synergistic effects which were in accordance with our results.²⁷

Correspondingly, in another study similar to our research, apatinib in combinatorial cases showed the anti-cancer effect. Furthermore, apatinib exhibited synergistic interactions with Paclitaxel plus 5-fluorouracil chemotherapeutic agents *in vivo*.²⁸ Also, a related study confirmed that apatinib displayed potentially inhibitory impacts in pancreatic cancerous cells and Astragalus polysaccharide increased the anti-cancer effects of apatinib by decreasing phosphorylation of AKT, and MMP-9.²⁹ In addition, a recent study indicated that in AGS cells Astragalus polysaccharide improved the antitumor efficacy of apatinib by inhibition of AKT signaling pathway.³⁰

Our study found that piperine or apatinib induced cytotoxicity effects in a dose-dependent manner. Nevertheless, when piperine was combined with apatinib, the expression of the MDM-2 as an oncogene was decreased significantly compared with the apatinib and piperine alone treated groups.

It has been indicated that NO induce apoptosis.²⁰ In this regards, piperine and apatinib can increased the NO level in double combinatorial cases including 0.5×IC50 and 0.2×IC50 concentrations versus single therapies.

Therefore, we speculated that apoptosis induction might be associated with the synergetic effects of piperine and apatinib in the present study. Our current study is evidence that indicate pharmacological regulation of the MDM2 gene expression may be related to cytotoxicity, which exerts by these treatments. Piperine in combination with apatinib showed more cytotoxic effects compared to monotherapies by reducing their concentrations in combination treatments. Further studies including clinical trials should be carried out for these new cancer therapies. The combination of apatinib and piperine not only reduced drug concentrations also promoted CRC treatment efficiency.

Conclusion

Overall, in this investigation, piperine as a natural anti-cancer agent is proving efficacious in combination with apatinib at low concentrations, which could be accounting for possible anti-cancer effects of this combination in CRC cells. Based on our results elevating cytotoxic activity of both agents in the combined treatment group might be related to the increased NO level. Nevertheless, the effects

of this combination in cell cycle regulation as well as the decreased expression level of MDM-2 might be examined by further studies.

Acknowledgments

We are thankful to Mr. Mohammad Aziz Rasouli for his technical support.

Ethical Issues

This article does not contain any studies with human subjects or animals performed by authors.

Conflicts of Interest

The authors declared no conflicts of interest.

References

1. Prashant A, Rangaswamy C, Yadav AK, Reddy V, Sowmya MN, Madhunapantula S. *In vitro* anticancer activity of ethanolic extracts of *Piper nigrum* against colorectal carcinoma cell lines. *Int J Appl Basic Med Res*. 2017;7(1):67-72. doi:10.4103/2229-516X.198531
2. Lu W, Ke H, Qianshan D, Zhen W, Guoan X, Honggang Y. Apatinib has anti-tumor effects and induces autophagy in colon cancer cells. *Iran J Basic Med Sci*. 2017; 20(9):990-95. doi:10.22038/IJBMS.2017.9263
3. Mohammadian M, Zeynali-Moghaddam S, Khadem Ansari M H, Rasmi Y, Fathi Azarbayjani A, Kheradmand F. Dihydropyrimidine Dehydrogenase Levels in Colorectal Cancer Cells Treated with a Combination of Heat Shock Protein 90 Inhibitor and Oxaliplatin or Capecitabine. *Adv Pharm Bull*. 2019;9(3):439-444. doi: 10.15171/apb.2019.052.
4. Moradi Z, Mohammadian M, Saberi H, Ebrahimifard M, Mohammadi Z, Ebrahimipour M, et al. Anti-cancer effects of chemotherapeutic agent; 17-AAG, in combined with gold nanoparticles and irradiation in human colorectal cancer cells. *Daru*. 2019;27(1):111-19. doi: 10.1007/s40199-019-00251-w.
5. Zeynali-Moghaddam S, Mohammadian M, Kheradmand F, Fathi-Azarbayjani A, Rasmi Y, Esna-Ashari O, et al. A molecular basis for the synergy between 17-allylamino-17-demethoxy geldanamycin with Capecitabine and Irinotecan in human colorectal cancer cells through VEGF and MMP-9 gene expression. *Gene*. 2019;684:30-38. doi: 10.1016/j.gene.2018.10.016.
6. Martini G, Ciardiello D, Vitiello PP, Napolitano S, Cardone C, Cuomo A, et al. Resistance to anti-epidermal growth factor receptor in metastatic colorectal cancer: What does still need to be addressed?. *Cancer Treat Rev*. 2020 ;86:102023. doi: 10.1016/j.ctrv.2020.102023.
7. Gao X, Liu J, Cho KB, Kedika S, Guo B. Chemopreventive Agent 3,3'-Diindolylmethane Inhibits MDM2 in Colorectal Cancer Cells. *Int J Mol Sci*. 2020;21(13):4642. doi:10.3390/ijms21134642
8. Liao X, Li H, Liu Z, Liao S, Li Q, Liang C, et al. Clinical efficacy and safety of apatinib in patients with advanced colorectal cancer as the late-line treatment. *Medicine (Baltimore)*. 2018; 97(50), e13635. doi: 10.1097/MD.00000000000013635.
9. Yaffe PB, Power Coombs MR, Doucette CD, Walsh M, Hoskin DW. Piperine, an alkaloid from black pepper, inhibits growth of human colon cancer cells via G1 arrest and apoptosis triggered by endoplasmic reticulum stress. *Mol Carcinog*. 2015; 54(10):1070-85. doi: 10.1002/mc.22176. Epub 2014 May 13.
10. Bolat ZB, Islek Z, Demir BN, Yilmaz EN, Sahin F, Ucisik MH. Curcumin- and Piperine-Loaded Emulsomes as Combinational Treatment Approach Enhance the Anticancer Activity of

- Curcumin on HCT116 Colorectal Cancer Model. *Front Bioeng Biotechnol.* 2020;8:50. doi:10.3389/fbioe.2020.00050
11. Esgandari K, Mohammadian M, Zohdiaghdam R, Rastin S J, Alidadi S, Behrouzki Z. Combined treatment with silver graphene quantum dot, radiation, and 17-AAG induces anticancer effects in breast cancer cells. *J Cell physiol.* 2020. doi: 10.1002/jcp.30046.
 12. Sajjadiyan Z, Ghadernejad H, Milani AT, Mohammadian M, Abdolahpour S, Taslimi S, et al. Preparation of silibinin loaded pegylatedniosomal nanoparticles and investigation of its effect on MCF-10A human breast cancer cell lines. *Der Pharm Lett.* 2016; 8;70-5.
 13. Maroufi NF, Vahedian V, Akbarzadeh M, Mohammadian M, Zahedi M, Isazadeh A, et al. The apatinib inhibits breast cancer cell line MDA-MB-231 in vitro by inducing apoptosis, cell cycle arrest, and regulating nuclear factor- κ B (NF- κ B) and mitogen-activated protein kinase (MAPK) signaling pathways. *Breast Cancer.* 2020;27(4):613-620. doi: 10.1007/s12282-020-01055-6..
 14. Mohammadian M, Feizollahzadeh S, Mahmoudi R, Toofani Milani A, Rezapour-Firouzi S, Karimi Douna B. Hsp90 Inhibitor; NVP-AUY922 in Combination with Doxorubicin Induces Apoptosis and Downregulates VEGF in MCF-7 Breast Cancer Cell Line. *Asian Pac J Cancer Prev.* 2020 ; 21(6):1773-78. doi: 10.31557/APJCP.2020.21.6.1773.
 15. Mohamadi N, Kazemi SM, Mohammadian M, Toofani Milani A, Moradi Y, Yasemi M, et al. Toxicity of cisplatin-loaded poly butyl cyanoacrylate nanoparticles in a brain cancer cell line: Anionic polymerization results. *Asian Pac J Cancer Prev.* 2017; 18(3):629-632. doi: 10.22034/APJCP.2017.18.3.629.
 16. Safarzadeh E, Sandoghchian Shotorbani S, Baradaran B. Herbal medicine as inducers of apoptosis in cancer treatment. *Adv Pharm Bull.* 2014;4(Suppl 1):421-427. doi:10.5681/apb.2014.062.
 17. Jan R, Chaudhry GE. Understanding Apoptosis and Apoptotic Pathways Targeted Cancer Therapeutics. *Adv Pharm Bull.* 2019;9(2):205-218. doi:10.15171/apb.2019.024.
 18. Hou H, Sun D, Zhang X. The role of *MDM2* amplification and overexpression in therapeutic resistance of malignant tumors. *Cancer Cell Int.* 2019;19; 216. doi:10.1186/s12935-019-0937-4.
 19. Zhao Y, Yu H, Hu W. The regulation of *MDM2* oncogene and its impact on human cancers. *Acta Biochim Biophys Sin (Shanghai).* 2014;46(3):180-189. doi:10.1093/abbs/gmt147
 20. Vahora H, Khan MA, Alalami U, Hussain A. The potential role of nitric oxide in halting cancer progression through Chemoprevention. *J Cancer Prev.* 2016;21(1):1-12. doi:10.15430/JCP.2016.21.1.1
 21. Khan MA, Chen HC, Wan XX, Tania M, Xu AH, Chen FZ, et al. Regulatory effects of resveratrol on antioxidant enzymes: a mechanism of growth inhibition and apoptosis induction in cancer cells. *Mol Cells.* 2013;35(3):219-25. doi: 10.1007/s10059-013-2259-z. Epub 2013 Feb 26. Erratum in: *Mol Cells.* 2013;35(4):355.
 22. Chou T-C. Drug combination studies and their synergy quantification using the Chou-Talalay method. *Cancer research.* 2010;70(2):440-6. doi: 10.1158/0008-5472.CAN-09-1947.
 23. Redondo-Blanco S, Fernández J, Gutiérrez-Del-Río I, Villar CJ, Lombó F. New Insights toward Colorectal Cancer Chemotherapy Using Natural Bioactive Compounds. *Front Pharmacol.* 2017; 8:109. doi:10.3389/fphar.2017.00109
 24. Nag S, Qin J, Srivenugopal KS, Wang M, Zhang R. The *MDM2*-p53 pathway revisited. *J Biomed Res.* 2013;27(4):254-271. doi:10.7555/JBR.27.20130030
 25. Rather RA, Bhagat M. Cancer Chemoprevention and Piperine: Molecular Mechanisms and Therapeutic Opportunities. *Front Cell Dev Biol.* 2018;6:10. doi:10.3389/fcell.2018.00010
 26. Pushpa Ragini S, Naga Divya AV, Anusha Ch, Kanthaiyah YV. Enhancement of paclitaxel and doxorubicin cytotoxicity in breast cancer cell lines in combination with piperine treatment and analysis of expression of autophagy and apoptosis genes. *J Med Sci Res.* 2014; 2: 62-7. doi: 10.17727/JMSR.2014/2-012.
 27. Motiwala MN, Rangari VD. Combined effect of paclitaxel and piperine on a MCF-7 breast cancer cell line in vitro: evidence of a synergistic interaction. *Synergy.* 2015; 2: 1- 6. doi:10.1016/j.synres.2015.04.001.
 28. Xu Z, Hu C, Chen S, Zhang C, Yu J, Wang X, et al. Apatinib enhances chemosensitivity of gastric cancer to paclitaxel and 5-fluorouracil. *Cancer Manag Res.* 2019;11:4905-4915. doi:10.2147/CMAR.S196372
 29. Wu J, Wang J, Su Q, Ding W, Li T, Yu J, et al. Traditional Chinese medicine Astragalus polysaccharide enhanced antitumor effects of the angiogenesis inhibitor apatinib in pancreatic cancer cells on proliferation, invasiveness, and apoptosis. *Onco Targets Ther.* 2018;11:2685-98. doi: 10.2147/OTT.S157129.
 30. Wu J, Yu J, Wang J, Zhang C, Shang K, Yao X, et al. Astragalus polysaccharide enhanced antitumor effects of Apatinib in gastric cancer AGS cells by inhibiting AKT signalling pathway. *Biomed Pharmacother.* 2018 ;100:176-83. doi: 10.1016/j.biopha.2018.01.140. Epub 2018 Feb 8.