

## Research Article

# The Tonoplast Intrinsic Protein Gene *KvTIP3* is Responsive to Different Abiotic Stresses in *Kosteletzkya virginica*

Xiaohua Liu,<sup>1,2</sup> Jieshan Cheng,<sup>1,2</sup> Fudong Jiang,<sup>3</sup> Meixia Liang,<sup>1,2</sup> Junjie Han,<sup>3</sup> Juan Zhang,<sup>1,2</sup> Hongxia Zhang,<sup>1,2,4</sup> and Xiaoli Tang <sup>1,2</sup>

<sup>1</sup>College of Agriculture, Ludong University, 186 Hongqizhong Road, Yantai, Shandong, China

<sup>2</sup>Key Laboratory of Molecular Module-Based Breeding of High Yield and Abiotic Resistant Plants in Universities of Shandong (Ludong University), 186 Hongqizhong Road, Yantai, Shandong, China

<sup>3</sup>Yantai Academy of Agricultural Sciences, 26 West Gangcheng Street, Yantai, Shandong, China

<sup>4</sup>Institute for Advanced Study of Coastal Ecology, Ludong University, 186 Hongqizhong Road, Yantai, Shandong, China

Correspondence should be addressed to Xiaoli Tang; tangxiaoli.1@163.com

Received 28 July 2019; Revised 18 November 2019; Accepted 4 December 2019; Published 8 January 2020

Academic Editor: Maxim Golovkin

Copyright © 2020 Xiaohua Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In higher plants, aquaporin proteins (AQPs) play important roles in the uptake of water across cell membranes. However, their functions in halophytes are still largely unknown. In this work, we isolated, cloned, and identified *KvTIP3*, a tonoplast intrinsic protein gene from *Kosteletzkya virginica*. Bioinformatic analyses demonstrated that *KvTIP3* encoded a tonoplast protein with the common properties of AQPs. Further multiple sequence alignment and phylogenetic analyses showed that *KvTIP3* shared 65%–82% homology with other AQPs from *Arabidopsis*, cotton, polar, and cocoa. Quantitative real-time PCR (qPCR) analyses revealed that *KvTIP3* was ubiquitously expressed in various tissues such as leaves, stems, and roots, with a predominant expression in roots. In addition, *KvTIP3* transcript was strongly induced by NaCl, low temperature, and ABA in *K. virginica*. Our findings suggest that *KvTIP3* encodes a new AQP possibly involved in multiple abiotic stress responses in *K. virginica*, and *KvTIP3* could be used as a potential candidate gene for the improvement of plants resistant to various abiotic stresses.

## 1. Introduction

Water is one of the essential requirements for all the organisms living on Earth [1, 2]. As a principal member of the biological kingdom, plants have evolved a series of mechanisms to absorb, transport, and utilize water. Unlike animals, plants cannot escape away from the adverse environment, such as drought and salt stress. Therefore, they must ensure the balance of water at both cellular and whole plant levels. AQPs are a family of channel proteins existing in all living organisms and play crucial roles in regulating the movement of water across the cell membrane [3]. For example, 13 AQPs were identified in human genome [4]. 35 and 36 AQPs were identified in *Arabidopsis* and maize, respectively [5, 6]. In addition, the number of AQPs isolated was 28 in *Vitis vinifera* L. [7], 28 in *Beta vulgaris* [8], 55 in *Populus trichocarpa* [9], 41 in *Solanum lycopersicum* [10], 63

in bamboo [11], 66 in *Glycine max* [12], and 71 in cotton [13].

AQPs are integral membrane proteins working in the form of tetramers and each monomer acts as an independent water channel [14]. According to the phylogenetic positions and subcellular localizations, plant AQPs are subdivided into four groups: plasma membrane intrinsic proteins (PIPs), tonoplast intrinsic proteins (TIPs), nodulin 26-like intrinsic proteins (NIPs), and small basic intrinsic proteins (SIPs) [15]. PIPs and TIPs are the most abundant proteins located in the plasma membrane and tonoplast, respectively [15]. Recently, three more groups, uncharacterized intrinsic proteins (XIPs), GlpF-like intrinsic proteins (GIPs), and hybrid intrinsic proteins (HIPs), were also proposed [16, 17]. Therefore, AQPs consist of seven subfamilies. TIPs are AQPs predominantly located at the tonoplast [15]. Structurally, TIPs possess the

conserved domains and motifs of AQPs, the six transmembrane helices with N-terminus and C-terminus located in the cytosol, and two NPA (asparagine-proline-alanine) motifs [7, 12]. The transmembrane helices constitute the channel for water and other substrate molecules. The conserved NPA motifs determine the substrate specificity [18]. In addition to transporting water across the membrane, TIPs can also facilitate the transport of other small molecules such as CO<sub>2</sub>, glycerol, NH<sub>3</sub>, arsenite, and silicon [19, 20]. Therefore, TIPs not only play a key role in water uptake but also function in many other aspects during the growth and development of plant.

Recently, an increasing number of researches revealed that TIPs played important functions in plant response to abiotic stresses. In *Arabidopsis*, TIP3;1, TIP3;2, and TIP4;1 were found to be involved in seed dormancy and germination in response to water stress [21]. In the halophyte *Thellungiella salsuginea* (salt cress), *TsTIP1;2* was involved in multiple stresses responses [22]. Recently, AQPs were reported to be involved in abiotic stress response in the halophyte *Eutrema salsugineum* [23]. In *Mesembryanthemum crystallinum*, osmotic stress can induce the relocation of McTIP1;2 from the tonoplast to other membranes [24]. Consistently, expression of *CsTIP2;1* from citrus in tobacco improved the growth and antioxidant capacity of transgenic plants under stress conditions [25]. Similarly, expression of tomato *SITIP2;2* and jojoba *ScPIP1* enhanced the tolerance of transgenic plants to various abiotic stresses [26]. However, the functions of TIPs are multifaceted. The same environmental stimulus can lead to both upregulated and down-regulated expression of different TIPs. Similarly, overexpression of TIPs does not always cause a positive effect on transgenic plants. For example, expression of *GsTIP2;1*, which was responsive to abiotic stresses, depressed salt and dehydration resistance in *Arabidopsis* [27]. Therefore, AQPs could function in various environmental stress responses.

*K. virginica*, a typical halophyte, is a new species with fine salt-tolerant characteristics at both physiological and molecular levels [28]. As an extremophile plant, *K. virginica* is very tolerant to adverse environmental stresses. It could grow and reproduce successfully in soil containing 0.3 to 2.5% sodium [29]. Previously, we examined the global gene expression profiles of *K. virginica* in response to salt stress [28]. Many gene expressions and pathways showed discrepancy. In this work, we isolated and identified a tonoplast aquaporin encoding gene *KvTIP3* from *K. virginica*, which showed the most significant gene expression variation as exhibited by transcriptomic analyses. Our results demonstrate that *KvTIP3* is a typical TIP encoding gene with a responsive expression to different abiotic stresses.

## 2. Materials and Methods

**2.1. Plant Materials.** *K. virginica* seeds were collected from the Yellow River Delta, Shandong Province, China. Seeds were surface sterilized and seedlings were cultivated as described before [30]. Two-week-old seedlings with the same

morphology were collected and used for the following experiments as described previously [31, 32].

**2.2. Abiotic Stress Treatments.** Two-week-old seedlings were subjected to high temperature, low temperature, high salinity, osmotic stress, and ABA treatments as described previously [30, 33].

**2.3. RNA Isolation and cDNA Synthesis.** Total RNA was extracted and analyzed as described previously [30]. The RNA samples were used to synthesize cDNA with TransScript One-Step gDNA Removal and cDNA Synthesis SuperMix (Transgen, China). Primers used for cDNA synthesis were random primer and anchored oligo (dT)<sub>18</sub>. Reaction was carried out by incubating the reaction mixture at 42°C for 30 min, and then at 85°C for 5 min. Finally, cDNA was examined with Agilent 2100 BioAnalyzer to confirm its quality and quantity.

**2.4. Cloning and Bioinformatic Analyses of *KvTIP3*.** The full length of *KvTIP3* gene was amplified by RACE technique according to the user's manual (SMART™ RACE cDNA Amplification Kit). Gene-specific primers (GSP-F: 5'-TAGTAAACCG CACGGACCACAGA-3' and GSP-R: 5'-CCAACATCTTAGTAGGCGGACCATT -3') and nested primers (NGSP-F: 5'-TTCCTCCAAGCAAGGCCAAG-3' and NGSP-R: 5'-GCCTTGGTAGGTTG-GAGGTGGGA-3') were designed according to the sequence of cDNA fragments in our previous transcriptomic sequencing data, which was deposited in Transcriptome Shotgun Assembly (TSA) Sequence Database with the accession number-GCJL00000000 [29]. The obtained gene fragment was ligated into the pEASY-T5 Zero Cloning Vector (Transgen, China) after purification and sequencing.

**2.5. Quantitative Real-Time RT-PCR (qPCR) Analyses.** qPCR was carried out with the instrument ABI Prism-SYMBOL 0 \f “Times New Roman” \s 127500 FAST (Applied Biosystems, Foster City, CA). The reaction system was performed using the SYBR Green Real-Time Selected Master Mix (Applied Biosystems by Life technologies), and the reaction condition was performed following the user's manual. cDNA was diluted to the required concentration (<50 ng/μL), and gene-specific primers were designed with the Beacon Designer (BD) 7.0 software based on the RACE results. Gene-specific primers used for qPCR are 5'-ATGCCAACTCGTAGATAC-3' and 5'-CTAGTAATCTT-CAGGAG C-3'. The reaction volume was 20 μL with a composition of 2 μL diluted cDNA template, 10 μL 2 × SYBR Master Mix, 1.5 μL primers, and 6.5 μL double distilled water. A reference gene, *18SrRNA*, which was the most stably expressed gene among the four selected reference genes (*18SrRNA*, *ACT*, *TUA*, and *EF*) under salt stress condition, was used as an internal control [34]. qPCR amplification was performed as following: initial denaturation at 95°C for 2 min, 40 cycles of 15 s at 95°C for denaturation, and 1 min at 60°C for annealing and extension.

Fluorescence signal was conducted at the temperature between 60°C and 90°C. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiment, was strictly followed throughout our experiment [35]. In addition, to guarantee the accuracy of the experiment, the primer specificity of qPCR was insured by the typical melting curve, amplification plot, and the single product of RT-PCR. cDNA was diluted to suitable concentrations to guarantee the C<sub>q</sub> values in the optimum range. For all the experiments, each sample was performed in triplicates and three biological replicates were performed.

**2.6. Data Analysis.** All the data were mean values  $\pm$  SD of three experiments. For qPCR analysis, the  $\Delta\Delta C_q$  method was used. ANOVA was applied to analyze the significant differences, and the *P* value was kept below 0.05. Sigma Plot 12.0 was used for plotting.

### 3. Results

**3.1. Isolation and Sequence Analysis of KvTIP3.** To understand the possible function of AQPs in abiotic stress tolerance in halophyte, a *K. virginica* homolog of tonoplast intrinsic protein gene, *KvTIP3*, was isolated (GenBank accession no. KT732779). The 1084-bp *KvTIP3* consists of a 154-bp 5' untranslated region (UTR), a 768-bp encoding region, and a 161-bp 3' UTR, encoding a 256 amino acid protein (KvTIP3) with a calculated molecular mass of 27 kD and isoelectric point of 6.58, as predicted by ProtParam (<http://www.expasy.org/tools/>).

**3.2. Bioinformatic Analysis of KvTIP3.** We further performed bioinformatic analyses. PsortII (<http://www.psort.org/>) prediction revealed that KvTIP3 protein was located in the secreted pathway with a probability of 94.4%. Further topological prediction with TMHMM (<http://www.cbs.dtu.dk/services/TMHMM-2.0/>) and TMPred ([http://www.ch.embnet.org/software/TMPRED\\_form.html](http://www.ch.embnet.org/software/TMPRED_form.html)) demonstrated that KvTIP3 contained six transmembrane helices with its N-terminus and C-terminus located in the cytoplasm (Figures 1(a) and 1(b)). It is well known that the activities of AQPs are regulated by their phosphorylation status. For example, the Snf1-related protein kinase 2.6 (SnRK2.6) was able to phosphorylate a cytosolic PIP2;1 peptide at Ser-121 to regulate the response of guard cells to ABA signaling [36]. Similarly, Cys116 and Cys118 have been proved to be the phosphorylation sites in TIP proteins [37]. We also examined the possible phosphorylation sites in KvTIP3 with NetPhos 2.0 (<http://www.cbs.dtu.dk/services/NetPhos/>) and found that KvTIP3 also contained 7 potential phosphorylation sites, including 4 Ser, 2 Thr, and 1 Tyr (Figure 1(c)). Moreover, we constructed a phylogenetic tree including the typical TIPs in *Arabidopsis* and other closely related organisms. We observed that KvTIP3 was clustered to the *Arabidopsis* AtTIP3 subgroup and was 82% homologous to TIP3-2 in *Gossypium hirsutum*, one of the most closely related organisms to *K. virginica* (Figure 2(d)). In addition, KvTIP3 shared 65%–78% amino acid sequence homology

with tonoplast aquaporins from other higher plants (Figure 2). KvTIP3, like TIPs from other higher plant, contained two NPA motifs and formed the cavity of AQPs to act as a selective barrier for the transport of water or small molecules [38].

**3.3. KvTIP3 Was Induced by Salt Stress in *K. virginica* Seedlings.** As a first step to know the expression profile of *KvTIP3* in *K. virginica*, we examined its relative expression levels in different tissues of two-week-old seedlings under normal growth condition by qPCR (Figure 3(a)). *KvTIP3* was ubiquitously expressed in various tissues including leaves, stems, and roots, with a predominant expression in the roots (Figure 3(b)). Based on the information extracted from the transcriptomic sequencing database (accession number-GCJL00000000) that *KvTIP3* cDNA fragments were significantly accumulated upon salt stress treatment, we performed qPCR to investigate the transcript levels of *KvTIP3* in *K. virginica* under both normal and salt stress conditions. Two-week-old seedlings were treated with 0, 200, 300, or 400 mM NaCl for 24 h. Although no significant phenotype change was observed after different treatments, salt stress significantly induced the expression of *KvTIP3* in a concentration-dependent manner, with the most significant induction with 300 mM NaCl (Figure 3(c)). When treated with 300 mM NaCl for different time periods, the transcript level of *KvTIP3* increased after 6 h and reached a maximum level at 12 h followed by a gradual decrease at 24 h (Figure 3(d)).

**3.4. Effects of Various Abiotic Stresses on the Expression of KvTIP3.** For the past few years, an increasing number of reports have revealed the important roles of AQPs in abiotic stress responses [39–41]. Thus, we tested the expression of *KvTIP3* under a few abiotic stresses and ABA treatments (Figures 4(a)–4(d)). Similar to the salt stress treatment, both low temperature and ABA treatments upregulated the expression of *KvTIP3*. The expression of *KvTIP3* increased after the seedlings were kept at 4°C for 24 hours (Figure 4(a)), whereas the expression of *KvTIP3* increased within 6 hours after the seedlings were treated with 100  $\mu$ M ABA (Figure 4(b)). Different from these observations, the expression of *KvTIP3* was downregulated when the seedlings were kept at 42°C or treated with 15% PEG (Figures 4(c) and 4(d)). All these results suggest that *KvTIP3* is an abiotic stress and ABA inducible or responsive gene.

### 4. Discussion

Previous studies have demonstrated that AQPs play important roles in plant growth, development, and in plant resistance to adverse environmental stresses. In Tulip (*Tulipa gesneriana*), both TgTIP1;1 and TgTIP1;2 were involved in petal development by regulating the absorption and transport of water [42]. Under osmotic or salt stress condition, AQPs in cucumber seedlings were able to regulate root and leaf hydraulic properties to respond to the environmental stimulus [41]. In this work, we isolated a novel AQP




FIGURE 1: The secondary structure, phosphorylation sites, and phylogenetic analysis of KvTIP3. (a, b) Transmembrane domain analysis of KvTIP3 with TMHMM and TMpred. (c) Phylogenetic analyses of KvTIP3 with other closely related TIPs proteins. All the sequences were extracted from the NCBI database. The phylogenetic tree was constructed by Mega 7.0 program using the neighbor-joining (NJ) method. (d) The potential phosphorylation sites of KvTIP3 predicted with NetPhos 2.0.

encoding gene *KvTIP3* from *K. virginica*. Acting as multi-functional membrane channels, AQPs have very highly conserved structure [7, 18]. Sequence analyses of *KvTIP3* revealed that it was one of the homologs of AQPs. The

molecular weights of the reported AQPs ranged from 23 to 31 kDa (243–302 amino acid residues). We found that *KvTIP3* gene encoded a 27 kDa AQP that consisted of 256 amino acid residues.




FIGURE 2: Amino acid sequence alignment of KvTIP3, AtTIP1-1, AtTIP1-2, AtTIP1-3, AtTIP2-1, AtTIP2-2, AtTIP2-3, AtTIP3-1, AtTIP3-2, AtTIP4-1, and AtTIP5-1 from *Arabidopsis thaliana*, GaTIP3-2 from *Gossypium arboreum*, GrTIP3-2 from *Gossypium raimondii*, PtTIP from *Populus trichocarpa*, and TcTIP from *Theobroma cacao*. Different colors in background display the different degrees of conserved sequences.

Similar to the AQPs from other plant species, KvTIP3 also contained six membrane-spanning  $\alpha$ -helices, as predicted with both TMHMM and TMPred software programs (Figures 1(a) and 1(b)). The water channel activities of AQPs were activated and deactivated via the phosphorylation and dephosphorylation of their phosphorylation sites [43]. The Snf1-related protein kinase 2.6 (SnRK2.6) was reported to be able to phosphorylate AQPs [36]. The

C-termini of several PIPs were also discovered to be phosphorylated by a sugar-induced receptor kinase [44]. We found that KvTIP3 contained 7 potential phosphorylation sites, implying that it may also play its functions via phosphorylation mechanism (Figure 1(c)). Further analyses indicated that KvTIP3 was homologous to and shared amino acid sequence homology with other plant AQPs (Figure 1(d)). It is well known that the conserved NPA




FIGURE 3: Expression analyses of *KvTIP3*. (a) Phenotype of a two-week-old *K. virginica* seedling used for salt stress or ABA treatment. (b) Expression analysis of *KvTIP3* in the leaves, stems, and roots of two-week-old *K. virginica* seedlings grown under normal growth condition. The expression level in leaves was set as the standard value 1. (c) Expression profiles of *KvTIP3* under different salt stress conditions. Two-week-old *K. virginica* seedlings were treated with 0, 200, 300, and 400 mM NaCl for 24 hours. (d) Expression profiles of *KvTIP3* under salt stress condition. Two-week-old *K. virginica* seedlings were treated with 300 mM NaCl for 0, 6, 12, and 24 hours. The mean values of three biological replicates were used for data analysis. The results of qPCR were normalized with the internal control gene, 18S rRNA. Error bar indicates SD ( $n=3$ ).

motif in AQPs functions as a selective filter for the substrate specificity [44]. Sequence analysis showed that *KvTIP3* contained two NPA motifs, indicating that the *KvTIP3* was a typical AQP (Figure 2).

The expression pattern of AQPs may affect their biological functions in the growth, development, and response to environmental stress of plants. We observed that the expression level of *KvTIP3* in roots was over 270-folds higher than in leaves and stems (Figure 3(b)). Similar expression pattern of AQPs was also observed in *Arabidopsis* [45], rice [46], grape [47], and barley [48]. In addition, the *KvLEA* gene which we reported previously also showed a similar expression pattern, indicating that these genes play important roles in roots of plants to improve salt resistance [30]. Staying the same with our transcriptome analysis, *KvTIP3* could be induced obviously by NaCl treatment [28]. However, different from *KvLEA* gene [30], 300 mM NaCl led to the most drastic accumulation to *KvTIP3* rather than 400 mM (Figure 3(c)). As for the different time of duration treatments, *KvTIP3* gave the

same expression profiles with *KvLEA*, proving that both *KvTIP3* and *KvLEA* were salt stress responsive genes of *K. virginica* (Figure 3(d)).

We also examined the expression levels of *KvTIP3* in response to different abiotic stresses and ABA treatment. *KvTIP3* was upregulated by low temperature and ABA but downregulated by high temperature and osmotic stress (Figures 4(a)–4(d)). This is consistent with previous studies that the responses of AQPs under adverse environment were diverse [49]. The responses of *KvTIP3* under high temperature and osmotic stress were not obvious (Figures 4(c) and 4(d)). Therefore, we speculated that *KvTIP3* may not be involved in high temperature response. However, TIPs were reported to take part in the response to osmotic stress [12]. In this research, the result showed that there should be other *KvTIPs* that participate in osmotic stress response in *K. virginica* rather than *KvTIP3*. Besides, ABA treatment gave rise to the most drastic accumulation to *KvTIP3* among these treatments (Figure 4(b)). For instance, the expression of *KvTIP3* reached as much as 180-




FIGURE 4: Expression profiles of *KvTIP3* upon different stresses or ABA treatments. (a) Expression profiles of *KvTIP3* upon low temperature treatment. (b) Expression profiles of *KvTIP3* upon ABA treatment. (c) Expression profiles of *KvTIP3* upon high temperature treatment. (d) Expression profiles of *KvTIP3* upon osmotic stress treatment. Error bar indicates SD ( $n = 3$ ).

fold after 6h ABA treatment. And the expression of *KvTIP3* increased rapidly as soon as after 2 h ABA treatment; hence, we speculated that *KvTIP3* may be involved in ABA-mediated stress responses in *K. virginica*. Actually, the intersections between ABA and AQPs have already been reported in other species. In *Arabidopsis*, AQPs functioned in ABA-triggered stomatal closure [36], and the vacuolar aquaporins were able to maintain seed longevity under the control of ABA [50]. In addition, ABA also could influence the phosphorylation of PIPs to regulate their activities [51]. Therefore, the exact mechanisms between *KvTIP3* and ABA deserve further investigation in the following research.

## 5. Conclusions

In this report, we characterized a novel tonoplast intrinsic protein gene *KvTIP3* from *K. virginica*. Bioinformatic analyses revealed that *KvTIP3* is a typical AQP with conserved transmembrane helices and NPA motifs. Gene expression profiles displayed that it was predominately expressed in roots and was responsive to various abiotic stresses. Our study on *KvTIP3* will enrich the gene resources and lay foundation for further study on AQPs. *KvTIP3* could be used as a potential candidate gene for the improvement of plants resistant to various abiotic stresses in the future.

## Data Availability

The data used to support the findings of this study are included within the article.

## Conflicts of Interest

The authors declare that there are no conflicts of interest.

## Authors' Contributions

Xiaohua Liu and Jieshan Cheng contributed equally to this work.

## Acknowledgments

The authors thank Ms. Jessie Zhang (Faculty of Health Sciences, McMaster University, Canada) for her critical reading and editing of this paper. This work has been jointly supported by the following grants: the National Natural Science Foundation of China (31901987); the Key R&D Project of Shandong Province (2017NC210012 and 2018GNC110007); the National Mega Project of GMO Crops of China (2016ZX08004-002-006); the Modern Agricultural Industry Technology System Innovation Team of

Shandong Province of China (SDAIT-02-05); and the National Natural Science Foundation of China (31870576).

## References

- [1] F. Chaumont and S. D. Tyerman, "Aquaporins: highly regulated channels controlling plant water relations," *Plant Physiology*, vol. 164, no. 4, pp. 1600–1618, 2014.
- [2] R. Kaldenhoff and M. Fischer, "Aquaporins in plants," *Acta Physiologica*, vol. 187, no. 1-2, pp. 169–176, 2006.
- [3] A. Shekoofa and T. R. Sinclair, "Aquaporin activity to improve crop drought tolerance," *Cells*, vol. 7, no. 9, Article ID 7090123, 2018.
- [4] A. R. Fox, L. C. Maistriaux, and F. Chaumont, "Toward understanding of the high number of plant aquaporin isoforms and multiple regulation mechanisms," *Plant Science*, vol. 264, pp. 179–187, 2017.
- [5] F. Chaumont, F. Barrieu, E. Wojcik, M. J. Chrispeels, and R. Jung, "Aquaporins constitute a large and highly divergent protein family in maize," *Plant Physiology*, vol. 125, no. 3, pp. 1206–1215, 2001.
- [6] U. Johanson, M. Karlsson, I. Johansson et al., "The complete set of genes encoding major intrinsic proteins in *Arabidopsis* provides a framework for a new nomenclature for major intrinsic proteins in plants," *Plant Physiology*, vol. 126, no. 4, pp. 1358–1369, 2001.
- [7] R. Fouquet, C. Léon, N. Ollat, and F. Barrieu, "Identification of grapevine aquaporins and expression analysis in developing berries," *Plant Cell Reports*, vol. 27, no. 9, pp. 1541–1550, 2008.
- [8] W. Kong, S. Yang, Y. Wang, M. Bendahmane, and X. Fu, "Genome-wide identification and characterization of aquaporin gene family in *Beta vulgaris*," *PeerJ*, vol. 5, Article ID e3747, 2017.
- [9] A. Gupta and R. Sankaramakrishnan, "Genome-wide analysis of major intrinsic proteins in the tree plant *Populus trichocarpa*: characterization of XIP subfamily of aquaporins from evolutionary perspective," *BMC Plant Biology*, vol. 9, no. 1, pp. 134–150, 2009.
- [10] S. Reuscher, M. Akiyama, C. Mori, K. Aoki, D. Shibata, and K. Shiratake, "Genome-wide identification and expression analysis of aquaporins in tomato," *PLoS One*, vol. 8, no. 11, Article ID e79052, 2013.
- [11] H. Y. Sun, S. N. Wang, Y. F. Lou et al., "Whole-genome and expression analysis of bamboo aquaporin genes reveal their functions involved in maintaining diurnal water balance in bamboo shoots," *Cells*, vol. 7, no. 7, Article ID 7110195, 2018.
- [12] D. Y. Zhang, M. Kumar, L. Xu et al., "Genome-wide identification of Major Intrinsic Proteins in Glycine soja and characterization of GmTIP2;1 function under salt and water stress," *Scientific Reports*, vol. 7, no. 1, Article ID 4106, 2017.
- [13] W. Park, B. E. Scheffler, P. J. Bauer, and B. T. Campbell, "Identification of the family of aquaporin genes and their expression in upland cotton (*Gossypium hirsutum* L.)," *BMC Plant Biology*, vol. 10, no. 1, pp. 142–150, 2010.
- [14] M. C. Berny, D. Gilis, M. Rooman, and F. Chaumont, "Single mutations in the transmembrane domains of maize plasma membrane aquaporins affect the activity of monomers within a heterotetramer," *Molecular Plant*, vol. 9, no. 7, pp. 986–1003, 2016.
- [15] C. Maurel, Y. Boursiac, D. T. Luu, V. Santoni, Z. Shahzad, and L. Verdoucq, "Aquaporins in plants," *Physiological Reviews*, vol. 95, no. 4, pp. 1321–1358, 2015.
- [16] G. P. Bienert, M. D. Bienert, T. P. Jahn, M. Boutry, and F. Chaumont, "Solanaceae XIPs are plasma membrane aquaporins that facilitate the transport of many uncharged substrates," *The Plant Journal*, vol. 66, no. 2, pp. 306–317, 2011.
- [17] D. Lopez, M. B. Amira, D. Brown et al., "The *Hevea brasiliensis* XIP aquaporin subfamily: genomic, structural and functional characterizations with relevance to intensive latex harvesting," *Plant Molecular Biology*, vol. 91, no. 4-5, pp. 375–396, 2016.
- [18] R. Kapilan, M. Vaziri, and J. J. Zwiazek, "Regulation of aquaporins in plants under stress," *Biological Research*, vol. 1, no. 4, 2018.
- [19] Z. He, H. Yan, Y. Chen et al., "An aquaporin PvTIP4;1 from *Pteris vittata* may mediate arsenite uptake," *New Phytologist*, vol. 209, no. 2, pp. 746–761, 2016.
- [20] X. Q. Zhao, N. Mitani, N. Yamaji, R. F. Shen, and J. F. Ma, "Involvement of silicon influx transporter OsNIP2;1 in selenite uptake in rice," *Plant Physiology*, vol. 153, no. 4, pp. 1871–1877, 2010.
- [21] S. Footitt, R. Clewes, M. Feeney, W. E. Finch-Savage, and L. Frigerio, "Aquaporins influence seed dormancy and germination in response to stress," *Plant Cell and Environment*, vol. 2019, pp. 1–15, 2019.
- [22] L.-L. Wang, A.-P. Chen, N.-Q. Zhong et al., "The *Thellungiella salsuginea* tonoplast aquaporin TsTIP1;2 functions in protection against multiple abiotic stresses," *Plant and Cell Physiology*, vol. 55, no. 1, pp. 148–161, 2014.
- [23] W. Qian, X. Yang, J. Li, R. Luo, X. Yan, and Q. Pang, "Genome-wide characterization and expression analysis of aquaporins in salt cress (*Eutrema salsugineum*)," *PeerJ*, vol. 7, Article ID 7664, 2019.
- [24] R. Vera-Estrella, B. J. Barkla, H. J. Bohnert, and O. Pantoja, "Novel regulation of aquaporins during osmotic stress," *Plant Physiology*, vol. 135, no. 4, pp. 2318–2329, 2004.
- [25] C. P. S. Martins, D. M. Neves, L. C. Cidade et al., "Expression of the citrus CsTIP2;1 gene improves tobacco plant growth, antioxidant capacity and physiological adaptation under stress conditions," *Planta*, vol. 245, no. 5, pp. 951–963, 2017.
- [26] X. Wang, F. Gao, J. Bing et al., "Overexpression of the jojoba aquaporin gene, *ScPIPI*, enhances drought and salt tolerance in transgenic *Arabidopsis*," *International Journal of Molecular Sciences*, vol. 20, no. 1, p. 153, 2019.
- [27] X. Wang, Y. Li, W. Ji et al., "A novel *Glycine soja* tonoplast intrinsic protein gene responds to abiotic stress and depresses salt and dehydration tolerance in transgenic *Arabidopsis thaliana*," *Journal of Plant Physiology*, vol. 168, no. 11, pp. 1241–1248, 2011.
- [28] X. L. Tang, H. Y. Wang, C. Y. Shao, and H. B. Shao, "Global gene expression of *Kosteletzkya virginica* seedlings responding to salt stress," *Plos One*, vol. 10, no. 4, Article ID e0124421, 2015.
- [29] H. Y. Wang, X. L. Tang, H. L. Wang, and H. B. Shao, "Physiological responses of *Kosteletzkya virginica* to coastal wetland soil," *Scientific World Journal*, vol. 2015, Article ID 354581, 2015.
- [30] X. L. Tang, H. Y. Wang, L. Y. Chu, and H. B. Shao, "KvLEA, a new isolated late embryogenesis abundant protein gene from *Kosteletzkya virginica* responding to multiabiotic stresses," *BioMed Research International*, vol. 2016, Article ID 9823697, 2015.
- [31] N. Ye, G. Zhu, Y. Liu, Y. Li, and J. Zhang, "ABA controls H<sub>2</sub>O<sub>2</sub> accumulation through the induction of OsCATB in rice leaves under water stress," *Plant and Cell Physiology*, vol. 52, no. 4, pp. 689–698, 2011.



- [32] M. H. Cui, K. S. Yoo, S. Hyoung et al., "An *Arabidopsis* R2R3-MYB transcription factor, AtMYB20, negatively regulates type 2C serine/threonine protein phosphatases to enhance salt tolerance," *FEBS Letters*, vol. 587, no. 12, 2013.
- [33] Z. Y. Ni, Z. Hu, Q. Y. Jiang, and H. Zhang, "GmNFYA3, a target gene of miR169, is a positive regulator of plant tolerance to drought stress," *Plant Molecular Biology*, vol. 82, no. 2, pp. 113–129, 2013.
- [34] X. Tang, H. Wang, C. Shao, and H. Shao, "Reference gene selection for qPCR normalization of *Kosteletzkya virginica* under salt stress," *Biomed Research International*, vol. 2015, pp. 1–8, 2015.
- [35] S. A. Bustin, V. Benes, J. A. Garson et al., "The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments," *Clinical Chemistry*, vol. 55, no. 4, pp. 611–622, 2009.
- [36] A. Grondin, O. Rodrigues, L. Verdoucq, S. Merlot, N. Leonhardt, and C. Maurel, "Aquaporins contribute to ABA-triggered stomatal closure through OST1-mediated phosphorylation," *The Plant Cell*, vol. 27, no. 7, pp. 1945–1954, 2015.
- [37] M. J. Daniels, F. Chaumont, T. E. Mirkov, and M. J. Chrispeels, "Characterization of a new vacuolar membrane aquaporin sensitive to mercury at a unique site," *The Plant Cell*, vol. 8, no. 4, pp. 587–599, 1996.
- [38] P. Mut, C. Bustamante, G. Martínez et al., "A fruit-specific plasma membrane aquaporin subtype PIP1;1 is regulated during strawberry (*Fragaria × ananassa*) fruit ripening," *Physiologia Plantarum*, vol. 132, no. 4, pp. 538–551, 2008.
- [39] M. Moshelion, O. Halperin, R. Wallach, R. Oren, and D. A. Way, "Role of aquaporins in determining transpiration and photosynthesis in water-stressed plants: crop water-use efficiency, growth and yield," *Plant, Cell & Environment*, vol. 38, no. 9, pp. 1785–1793, 2015.
- [40] A. Pou, H. Medrano, J. Flexas, and S. D. Tyerman, "A putative role for TIP and PIP aquaporins in dynamics of leaf hydraulic and stomatal conductances in grapevine under water stress and re-watering," *Plant, Cell & Environment*, vol. 36, no. 4, pp. 828–843, 2013.
- [41] Z.-J. Qian, J.-J. Song, F. Chaumont, and Q. Ye, "Differential responses of plasma membrane aquaporins in mediating water transport of cucumber seedlings under osmotic and salt stresses," *Plant, Cell & Environment*, vol. 38, no. 3, pp. 461–473, 2015.
- [42] A. K. Azad, R. Hanawa, T. Ishikawa, Y. Sawa, and H. Shibata, "Expression profiles of aquaporin homologues and petal movement during petal development in *Tulipagesneriana*," *Physiologia Plantarum*, vol. 148, no. 3, pp. 397–407, 2013.
- [43] W. Wei, E. Alexandersson, D. Gollack, A. J. Miller, P. O. Kjellbom, and W. Fricke, "HvPIP1;6, a barley (*Hordeum vulgare* L.) plasma membrane water channel particularly expressed in growing compared with non-growing leaf tissues," *Plant and Cell Physiology*, vol. 48, no. 8, pp. 1132–1147, 2007.
- [44] X. N. Wu, C. Sanchez Rodriguez, H. Pertl-Obermeyer, G. Obermeyer, and W. X. Schulze, "Sucrose-induced receptor kinase SIRK1 regulates a plasma membrane aquaporin in *Arabidopsis*," *Molecular & Cellular Proteomics*, vol. 12, no. 10, pp. 2856–2873, 2013.
- [45] E. Alexandersson, L. Fraysse, S. Sjövall-Larsen et al., "Whole gene family expression and drought stress regulation of aquaporins," *Plant Molecular Biology*, vol. 59, no. 3, pp. 469–484, 2005.
- [46] J. Sakurai, F. Ishikawa, T. Yamaguchi, M. Uemura, and M. Maeshima, "Identification of 33 rice aquaporin genes and analysis of their expression and function," *Plant and Cell Physiology*, vol. 46, no. 9, pp. 1568–1577, 2005.
- [47] R. K. Vandeleur, G. Mayo, M. C. Sheldon, M. Gilliam, B. N. Kaiser, and S. D. Tyerman, "The role of plasma membrane intrinsic protein aquaporins in water transport through roots: diurnal and drought stress responses reveal different strategies between isohydric and anisohydric cultivars of grapevine," *Plant Physiology*, vol. 149, no. 1, pp. 445–460, 2009.
- [48] T. Knipfer, M. Besse, J.-L. Verdeil, and W. Fricke, "Aquaporin-facilitated water uptake in barley (*Hordeum vulgare* L.) roots," *Journal of Experimental Botany*, vol. 62, no. 12, pp. 4115–4126, 2011.
- [49] J. Madrid-Espinoza, N. Brunel-Saldias, F. P. Guerra, A. Gutiérrez, and A. Del Pozo, "Genome-wide identification and transcriptional regulation of aquaporin genes in bread wheat (*Triticum aestivum* L.) under water stress," *Gene*, vol. 9, no. 10, Article ID 497, 2018.
- [50] Z. Mao and W. Sun, "Arabidopsis seed-specific vacuolar aquaporins are involved in maintaining seed longevity under the control of ABSCISIC ACID INSENSITIVE 3," *Journal of Experimental Botany*, vol. 66, no. 15, pp. 4781–4794, 2015.
- [51] K. G. Kline, G. A. Barrett-Wilt, and M. R. Sussman, "In planta changes in protein phosphorylation induced by the plant hormone abscisic acid," *Proceedings of the National Academy of Sciences*, vol. 107, no. 36, pp. 15986–15991, 2010.