



Colocalization of pain and reduced intraepidermal nerve fiber density in individuals with HIV-associated sensory neuropathy

Imraan G. Patel^a, Peter R. Kamerman^{a,b,*}

Abstract

Introduction: There is poor correlation between decreases in intraepidermal nerve fiber density (IENFD) and the presence of pain in HIV-associated sensory neuropathy (HIV-SN) and other painful distal symmetrical polyneuropathies.

Objectives: We investigated whether in individuals with HIV-SN, having pain at the ankle skin biopsy site was associated with lower IENFD compared to when there was no pain at the ankle biopsy site.

Methods: We recruited 15 individuals with symptomatic HIV-SN. Nine had pain at the site where the ankle biopsy was taken, whereas 6 did not. Skin punch biopsies for IENFD quantification were taken from the ankle and the thigh. Contrasts between the 2 groups were made using the overlap of confidence interval (CI) method.

Results: Intraepidermal nerve fiber density was substantially lower in the group that had pain at the site of the ankle biopsy compared with the other group (6.6 [CI: 5.3–7.2] vs 3.3 [CI: 10.0–15.0] fibers/mm). However, there was no group differences at the thigh biopsy site (15.6 [CI: 15.0–15.9] vs 16.2 [CI: 14.5–17.8] fibers/mm). When taking the ratio of ankle IENFD:thigh IENFD, the point estimate for the pain at the ankle group (0.43 [CI: 0.36–0.48]) was about half that of the other group (0.81 [CI: 0.68–0.87]).

Conclusion: Thus, colocalization of pain to the ankle is associated with meaningful decreases in ankle IENFD.

Keywords: HIV, Neuropathy, Neuropathic pain, Intraepidermal nerve fiber density

1. Introduction

HIV-associated sensory neuropathy (HIV-SN) is distal symmetrical polyneuropathy associated with HIV infection and its treatment.⁷ Like other peripheral neuropathies, it is often but not always painful.^{4,8,15} There have been efforts to correlate the presence of pain in HIV-SN to objective measures of a peripheral nerve lesion.

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

^a Brain Function Research Group, School of Physiology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa, ^b School of Pharmacy and Biomedical Sciences, Faculty of Health Sciences, Curtin University, Perth, Australia

*Corresponding author. Address: Peter R. Kamerman, Brain Function Research Group, School of Physiology, Faculty of Health Sciences, University of the Witwatersrand, 7 York Rd, Parktown 2193, South Africa. Tel.: +27 (0)11 717 2363. E-mail address: peter.kamerman@wits.ac.za (P.R. Kamerman).

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.painrpts.com).

Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of The International Association for the Study of Pain. This is an open access article distributed under the Creative Commons Attribution-ShareAlike License 4.0 (CC BY-SA) which allows others to remix, tweak, and build upon the work, even for commercial purposes, as long as the author is credited and the new creations are licensed under the identical terms.

PR9 4 (2019) e778

<http://dx.doi.org/10.1097/PR9.0000000000000778>

Quantitative sensory testing on the foot has shown that compared to nonpainful HIV-SN, painful HIV-SN is associated with mechanical allodynia and mechanical hyperalgesia.¹ Although intraepidermal nerve fiber density (IENFD) is reduced in HIV-SN and is associated with pain intensity in those individuals with pain, no associations have been identified between IENFD and the presence of symptomatic HIV-SN.^{13,16,18} However, lower ankle IENFD was associated with increased risk of transitioning from asymptomatic to symptomatic HIV-SN.⁶ We hypothesized that this tangential evidence for an association between IENFD and the presence of pain in HIV-SN results from the anatomical separation of where pain may be occurring (typically the feet),¹² and where distal skin biopsies for IENFD quantification are taken (the ankle).^{10,11} We therefore undertook an exploratory study assessing whether in individuals with HIV-SN, having pain at the ankle biopsy site was associated with lower IENFD at the ankle compared to having no pain or pain restricted to the feet.

2. Methods

The study was approved by the Human Research Ethics Committee of the University of the Witwatersrand.

2.1. Study population

We recruited a convenience sample of people living with HIV/AIDS and who were attending the Green House Pharmacy at

Chris-Hani Baragwanath Hospital, Johannesburg, South Africa. Study inclusion criteria were: (1) being on stable combination antiretroviral therapy for at least 6 months; (2) having a current CD4 T-cell count >350 cells/mm³; (3) fulfilling our diagnostic criteria for the presence of distal symmetrical polyneuropathy; (4) the onset of signs or symptoms of neuropathy occurred after the initiation of combination antiretroviral therapy; (5) not having diabetes mellitus, and (6) not having any current skin lesions on the lower legs. Thirty-five people provided informed consent and were screened. Fifteen met the inclusion requirements.

2.2. Neuropathy assessment

We assessed for symptomatic neuropathy in the lower limbs using the ACTG Brief Peripheral Neuropathy Screen (BPNS),² which has a case definition of the bilateral presence of at least one neurological sign (reduced or absent vibration detection in the great toes and absent ankle reflexes), and at least one symptom of polyneuropathy (pain, numbness, and paresthesias) in a consistent neuroanatomical pattern.

Once neuropathy had been diagnosed, the presence, severity, and location of any symptoms of pain were assessed using a guided interview. Pain intensity was assessed using an 11-point numerical pain rating scale. Depending on the outcome of the assessment of pain, participants were grouped into a “pain” group ($n = 9$) or a “no pain” group ($n = 6$). Participants in the “no pain” group met our case definition for neuropathy, but did not report experiencing pain at the ankle or thigh biopsy sites (2 participants had no pain at all, but had numbness or paresthesias, and 4 had pain that was limited to the soles of their feet). Participants in the “pain group” met our case definition for neuropathy and they had pain that extended proximally from the foot to at least the ankle biopsy site, but not past the knee.

We did not perform fine-scale mapping of the neuropathy (eg, pinprick sensitivity or brush sensitivity) to the area of the ankle pain, and therefore our diagnostic certainty is limited to possible neuropathic pain. The results of the study need to be considered in the light of this diagnostic certainty.

2.3. Skin biopsies

Three-millimeter (3 mm) skin punch biopsies were taken from the ankle (10 cm above the lateral malleolus) and from the lateral aspect of the proximal thigh. The thigh biopsies served as a reference site because this site is only affected very late in the progression of HIV-associated sensory neuropathy.¹⁷ We followed published methods for the biopsy procedure, tissue fixing and preparation, panaxonal marker protein gene product 9.5 immunohistochemical staining, and fiber density quantification.¹¹ The single assessor (I.G.P.) was blinded to the source of the biopsy (“pain” vs “no pain,” and thigh vs ankle). A detailed description of all procedures is provided in Supplement 1 (available at <http://links.lww.com/PR9/A53>).

2.4. Data analysis

We assessed the following: (1) the proportion of ankle biopsies with fiber densities below the normal range, (2) the fiber densities at the ankle and at the thigh, and (3) the ankle:thigh fiber density ratios. Because of the small sample size, we chose to conduct exploratory analysis using the overlap of confidence intervals (CIs) method. That is, failure of CIs to overlap indicates a significant and meaningful difference. The CI method of significance testing requires the confidence level of the intervals be adjusted to maintain a 5% error

rate.^{5,9} Bias-corrected accelerated CIs were generated using bootstrapping, with 100000 replicates. All analyses were performed in the R Statistical Environment (R v3.5.2).¹⁴ A full analysis script is provided in the supplementary material (Supplement 2, available at <http://links.lww.com/PR9/A53>).

3. Results

3.1. Description of the groups

The sex ratio (female:male) was 3:3 in the “no pain” group and 7:2 in the “pain” group. Participants in the 2 groups were of similar age (“pain” group [years]: mean = 41, SD = 10; “no pain” group: mean = 43, SD = 4; 95% CI for the difference in location = -5 to 11 years), and had similar current CD4 T-cell counts (“pain” group [cells/mm³]: median = 469, range = 353–653; “no pain” group: median = 543, range = 368–770; 95% CI for the difference in location = -90 to 205 cells/mm³). In the “pain” group, median pain intensity was 10 (range: 7–10), and the 4 participants with foot pain in the “no pain” group had a median pain intensity of 5 (range: 1–10).

3.2. Intraepithelial nerve fiber density

Figure 1 shows IENFD data for the “pain” group and the “no pain” group at the ankle and thigh, and the ankle:thigh IENFD ratio for the groups. All participants in the “pain” group (9/9) had ankle fiber densities below the fifth percentile for healthy adults, whereas only 2 (33%) of individuals in the “no pain” group (2/6) had values below the fifth percentile.¹¹

At the ankle (Fig. 1: Ankle), the point estimates and CIs of the 2 groups show no overlap, such that upper confidence limit in the “pain” group was about 2.5 fibers/mm lower than the lower limit of the CI for the “no pain” group, indicating that the data are compatible with a meaningful difference in IENFD between the 2 groups. At the thigh, there was substantial overlap in the CIs for the 2 groups’ IENFD at the thigh (Fig. 1: Thigh). Thus, at the reference site, the data are compatible with the 2 groups having similar IENFD. When expressed as the ratio of ankle IENFD to thigh IENFD, the “pain” group had a lower ratio compared to that of the “no pain” group, with no overlap in point estimates and CIs of the 2 groups, indicating a statistically significant and a meaningful difference (Fig. 1: Ankle:thigh).

4. Discussion

In this exploratory study, we assessed whether having pain at the ankle biopsy site was associated with lower IENFD at the ankle compared to having no pain or pain restricted to the feet. We report that previous failures to identify an association between IENFD changes and the presence of painful symptoms in HIV-SN may have resulted from the anatomical separation of where pain was occurring, and where distal skin biopsies for IENFD quantification were taken. More broadly, our findings may, in part, explain the poor diagnostic performance of IENFD in HIV-SN, and ankle IENFD may best correlate with the presence of symptoms when the symptoms colocalize to the place the biopsy is taken from.

Our data are consistent with the findings of a recent meta-analysis on the association between IENFD and features of distal symmetrical polyneuropathies.⁸ The meta-analysis found low levels of association between IENFD and other features of neuropathy, except for when measurements and IENFD assessment were colocalized. In this case, the measures were cold detection threshold and warm detection threshold. Taken together with our data, the results of the meta-analysis have broad implications for research on distal symmetrical polyneuropathies




Figure 1. Mean and confidence interval (CI) of intraepidermal fiber densities (IENFDs) for the “pain” group (pain at biopsy site: yes) and the “no pain” group (pain at biopsy site: no) at the ankle (top left panel) and the thigh (top right panel), and the ankle:thigh IENFD ratio (bottom left panel). Confidence intervals were calculated using bootstrapping and the confidence limits reflect 95% confidence intervals adjusted to maintain a 5% long-term error rate (see the data analysis section for details). Absence of overlap between confidence intervals indicates the data are compatible with a statistically significant and a meaningful difference between groups. At the ankle, there was no overlap of the point estimates and confidence intervals, with the “pain” group having a mean of 6.6 (85.9% CI: 5.3–7.2) fibers/mm, whereas the “no pain” group had a mean of 13.3 (85.9% CI: 10.0–15.0) fibers/mm. So, the “pain” group had a point estimate of about half the fiber density of the “no pain” group, and the upper limit of the confidence interval for the “pain” group was about 2.5 fibers/mm different from that of the lower limit of the confidence interval of the “no pain” group. By contrast, there was substantial overlap in the confidence intervals at the thigh, with the “pain” group having a mean of 15.6 (88.4% CI: 15.0–15.9) fibers/mm, and the “no pain” had a similar mean of 16.2 (88.4% CI: 14.5–17.8) fibers/mm. The overlap between confidence intervals, and point estimates by confidence intervals indicates that the data are compatible with no notable difference between the 2 groups at the reference site. When expressed as the ankle:thigh IENFD ratio, the “pain” group had a mean ratio of 0.43 (83.6% CI: 0.36–0.48), which is about half that of the “no pain” group, which had a mean ratio of 0.81 (83.6% CI: 0.68–0.87). Thus, the ratio data reinforce the findings from the ankle data.

in terms of making sure that measurements of function, symptoms, and morphology target the same anatomical area.

Our study was only a small exploratory study, which included a heterogeneous “no pain” group. Moreover, our phenotyping did not include direct assessment of the skin area at the site of the ankle pain (eg, quantitative sensory testing), and thus our phenotypic grading at the biopsy site is that of possible neuropathic pain.³ Nevertheless, we believe that we have analyzed the data conservatively, and that our findings are not spurious. Indeed, our findings are consistent with the presumed neurobiology, where morphological changes should associate with symptoms in a circumscribed neuroanatomical pattern.³ As such, we believe that there is merit in the study, and that it provides impetus for further research on the association between IENFD and colocalization of other characteristics of distal symmetrical polyneuropathies.

Disclosures

The authors have no conflict of interest to declare.

The research was funded by the Rated Researchers Programme of the National Research Foundation of South Africa, and the Medical Faculty Research Endowment Fund of the Faculty of Health Sciences, University of the Witwatersrand.

Appendix A. Supplemental digital content

Supplemental digital content associated with this article can be found online at <http://links.lww.com/PR9/A53>.

Article history:

Received 25 March 2019

Received in revised form 7 June 2019

Accepted 9 July 2019

References

- [1] Bouhassira D, Attal N, Willer JC, Brasseur L. Painful and painless peripheral sensory neuropathies due to HIV infection: a comparison using quantitative sensory evaluation. *PAIN* 1999;80:265–72.

- [2] Cherry CL, Wesselingh SL, Lal L, McArthur JC. Evaluation of a clinical screening tool for HIV-associated sensory neuropathies. *Neurology* 2005;65:1778–81.
- [3] Finnerup NB, Haroutounian S, Kamerman P, Baron R, Bennett DLH, Bouhassira D, Cruccu G, Freeman R, Hansson P, Nurmikko T, Raja SN, Rice ASC, Serra J, Smith BH, Treede RD, Jensen TS. Neuropathic pain: an updated grading system for research and clinical practice. *PAIN* 2016; 157:1599–606.
- [4] Freeman R, Baron R, Bouhassira D, Cabrera J, Emir B. Sensory profiles of patients with neuropathic pain based on the neuropathic pain symptoms and signs. *PAIN* 2014;155:367–76.
- [5] Ryan GW, Leadbetter SD. On the misuse of confidence intervals for two means in testing for the significance of the difference between the means. *J Mod Appl Stat Methods* 2002;1:56.
- [6] Herrmann DN, McDermott MP, Henderson D, Chen L, Akowuah K, Schifitto G. Epidermal nerve fiber density, axonal swellings and QST as predictors of HIV distal sensory neuropathy. *Muscle Nerve* 2004;29: 420–7.
- [7] Kamerman PR, Wadley AL, Cherry CL. HIV-associated sensory neuropathy: risk factors and genetics. *Curr Pain Headache Rep* 2012; 16:226–36.
- [8] Karlsson P, Hincker AM, Jensen TS, Freeman R, Haroutounian S. Structural, functional, and symptom relations in painful distal symmetric polyneuropathies: a systematic review. *PAIN* 2019;160:286–97.
- [9] Knol MJ, Pestman WR, Grobbee DE. The (mis)use of overlap of confidence intervals to assess effect modification. *Eur J Epidemiol* 2011;26:253–4.
- [10] Lauria G, Cornblath DR, Johansson O, McArthur JC, Mellgren SI, Nolano M, Rosenberg N, Sommer C; European Federation of Neurological Societies. EFNS guidelines on the use of skin biopsy in the diagnosis of peripheral neuropathy. *Eur J Neurol* 2005;12:747–58.
- [11] Lauria G, Hsieh ST, Johansson O, Kennedy WR, Leger JM, Mellgren SI, Nolano M, Merkies ISJ, Polydefkis M, Smith AG, Sommer C, Valls-Solé J. European Federation of Neurological Societies/Peripheral Nerve Society Guideline on the use of skin biopsy in the diagnosis of small fiber neuropathy. Report of a joint task force of the European Federation of Neurological Societies and the Peripheral Nerve Society. *Eur J Neurol* 2010;17:903–e49.
- [12] Phillips TJC, Brown M, Ramirez JD, Perkins J, Woldeamanuel YW, Williams ACdeC, Orengo C, Bennett DLH, Bodi I, Cox S, Maier C, Krumova EK, Rice ASC. Sensory, psychological, and metabolic dysfunction in HIV-associated peripheral neuropathy: a cross-sectional deep profiling study. *PAIN* 2014;155:1846–60.
- [13] Polydefkis M, Yiannoutsos CT, Cohen BA, Hollander H, Schifitto G, Clifford DB, Simpson DM, Katzenstein D, Shriver S, Hauer P, Brown A, Haidich AB, Moo L, McArthur JC. Reduced intraepidermal nerve fiber density in HIV-associated sensory neuropathy. *Neurology* 2002;58:115–19.
- [14] R Core Team. R: a language and environment for statistical computing. Vienna, Austria, 2018 Available at: <http://www.R-project.org>. Accessed December 20, 2018.
- [15] Sadosky A, McDermott AM, Brandenburg NA, Strauss M. A review of the epidemiology of painful diabetic peripheral neuropathy, postherpetic neuralgia, and less commonly studied neuropathic pain conditions. *Pain Pract* 2008;8:45–56.
- [16] Skopelitis E, Aroni K, Kontos AN, Konstantinou K, Kokotis P, Karandreas N, Kordossis T. Early detection of subclinical HIV sensory polyneuropathy using intraepidermal nerve fibre density quantification: association with HIV stage and surrogate markers. *Int J STD AIDS* 2007;18:856–60.
- [17] Wulff EA, Wang AK, Simpson DM. HIV-associated peripheral neuropathy: epidemiology, pathophysiology and treatment. *Drugs* 2000;59:1251–60.
- [18] Zhou L, Kitch DW, Evans SR, Hauer P, Raman S, Ebenezer GJ, Gerschenson M, Marra CM, Valcour V, Diaz-Arrastia R, Goodkin K, Millar L, Shriver S, Asmuth DM, Clifford DB, Simpson DM, McArthur JC; NARC and ACTG A5117 Study Group. Correlates of epidermal nerve fiber densities in HIV-associated distal sensory polyneuropathy. *Neurology* 2007;68:2113–19.