

Acridone alkaloids with cytotoxic and antimalarial activities from *Zanthoxylum simullans* Hance

Chao Wang, Jinfu Wan¹, Zhinan Mei, Xinzhou Yang

College of Pharmacy, South-Central University for Nationalities, ¹Institute of Yunnan Baiyao Group Co. Ltd, China

Submitted: 03-12-2012

Revised: 23-01-2013

Published: 07-02-2014

ABSTRACT

Background: *Zanthoxylum simullans* Hance is a popular natural spice belonging to the Rutaceae family and it is one of the common prescribed herbs in traditional Chinese medicine. **Materials and Methods:** The chemical constituents were mainly isolated and purified by silica gel column chromatography and semi-preparative High Performance Liquid Chromatography. Their structures were identified by comparing the spectral data with those reported in the literature. Cytotoxic activities for the isolated acridone alkaloids were evaluated against two prostate cancer cell lines PC-3M and Lymph Node Carcinoma of Prostate (LNCaP), and their antimalarial activities were tested against two different strains of the parasite *Plasmodium falciparum* 3D7, and Dd2. **Results:** The root bark MeOH extract of *Z. simullans* Hance afforded β -sitosterol, 4-methoxy benzoic acid, daucosterol, and five acridone alkaloids, normelicopidine, normelicopine, melicopine, melicopidine, and melicopicine. All five acridone alkaloids were isolated from this plant for the first time and exhibited certain cytotoxic and antimalarial activities *in vitro*. **Conclusion:** Normelicopidine was the most active against PC-3M, LNCaP and Dd2 with IC₅₀ values of 12.5, 21.1, and 18.9 ug/mL respectively.

Key words: Acridone alkaloids, antimalarial, cytotoxicity, *Zanthoxylum simullans* Hance

Access this article online

Website:

www.phcog.com

DOI:

10.4103/0973-1296.126669

Quick Response Code:



INTRODUCTION

The genus *Zanthoxylum*, family Rutaceae comprises about 250 species of deciduous, evergreen trees and shrubs distributed world-wide around the tropical and subtropical areas. It is economically important because of their alimentary, industrial, and medicinal applications.^[1] *Zanthoxylum simullans* Hance, a popular natural spice, has long been prescribed in traditional Chinese medicine for the treatment of stomach ache, toothache, intestinal worms, eczema, and pruritus.^[2] Extensive phytochemical investigations have been carried out on this plant, along with the study of its biological activities, especially, antiplatelet aggregation.^[3-7] Most alkaloids isolated from this plant exhibited strong antiplatelet aggregation activities, except for zanthosimuline, a new acridone alkaloid found in the root bark. It has demonstrated a general cytotoxic response when evaluated against a variety of human cancer cell lines and P388 cells.^[4] In this paper, we conducted a search for more bioactive constituents from *Z. simullans* Hance.

Address for correspondence:

Dr. Xinzhou Yang, College of Pharmacy, South-Central University for Nationalities, 708 Minzu Road, Wuhan 430074, P. R. China.
E-mail: yxzyxz01@163.com

MATERIALS AND METHODS

Plant material

The root barks of *Z. simullans* Hance were collected in September 2009 at Jingxi county, Guangxi province, China and identified by Dr. Jingquan Yuan, Associate Professor of Guangxi Medicinal Herb Garden. A voucher specimen (No. 20090018Z) has been deposited in the Herbarium of South-Central University for Nationalities, Wuhan, P. R. China.

General

Melting points were determined on a Kofler melting point apparatus and are uncorrected. Ultraviolet (UV) spectra were determined on a Shimadzu UV-250 spectrometer. ¹H and ¹³C NMR (Carbon-13 Nuclear Magnetic Resonance) spectra were recorded on a Bruker DRX-400 spectrometer with Tetramethylsilane (TMS) as an internal standard. The chemical shifts (δ) were reported in ppm, and coupling constants (J) were given in Hz. The ESI-MS and EI-MS (Electrospray Ionization Mass Spectrometry and Electron Impact Mass Spectrometry) data were recorded on the Quadrupole Time-of-flight (Q-TOF) Mass Spectrometry Micro Liquid Chromatography-Tandem Mass Spectrometry (LC-MS-MS) and Finnigan MAT-95

mass spectrometers. Semi-preparative HPLC system: Two PrepStar SD-1 solvent delivery modules, a ProStar UV-Vis 320 detector and a ProStar 701 Fraction Collector (Varian, Walnut Creek, CA, USA); a Sunfire RP-18 column (5 μ m; 250 mm \times 20 mm, waters) for semi-preparative isolation. The following types of silica gel were used: 300-400 mesh for column chromatography and GF₂₅₄ for Thin Layer Chromatography (TLC) (Qing Dao Hai Yang Chemical Group Co.). Sephadex LH 20 (GE Healthcare) was also used for column chromatography. Cell lines PC-3M, LNCaP and HEK293 were purchased from the American Type Culture Collection. Parasite strains Dd2 and 3D7 were from the Queensland Institute of Medical Research.

Extraction

Air-dried root barks of *Z. simullans* Hance (200 g) were ground and then extracted by maceration at room temperature with MeOH in a 1 L conical flask mounted on a shaking table (3 \times 0.6 L, 4 h each). The solvents were evaporated at reduced pressure to yield 18.6 g MeOH extract.

Isolation

The MeOH extract was subjected to CC over silica gel (500 g) eluted gradiently with CH₂Cl₂-MeOH (25:1-1:2) to afford fractions 1-7. Fraction 2 (2.1 g) was rechromatographed on silica gel (120 g) with CH₂Cl₂-MeOH (20:1-10:1) to give compounds 1 (135 mg), 2 (27 mg) and 5 (87 mg). The crude alkaloids (0.9 g) derived from fraction 3 (1.6 g) after filtration over Sephadex LH 20 with MeOH were further submitted to semi-preparative HPLC eluted with a linear gradient of 10-70% aq. MeOH in 0.5% TFA at 9.0 mL/min for 120 min to yield successively compounds 4 (12 mg, t_R 32.5 min), 5 (39 mg, t_R 36.1 min), 6 (5 mg, t_R 47.7 min), 7 (16 mg, t_R 64.6 min), 8 (78 mg, t_R 68.9 min).

Compound 1, colorless crystal (MeOH), mp, 140-142°C; Electron Impact Mass Spectrometry (EIMS) m/z 414 [M]⁺, 396, 381, 354, 329, 303, 273, 255; ¹H-NMR (CDCl₃): δ 5.33 (1H, d, J = 5.0 Hz, H-6), 3.50 (1H, m, H-3), 0.99 (3H, s, H-19), 0.92 (3H, d, J = 8.1 Hz, H-21), 0.84 (3H, t, J = 7.7 Hz, H-29), 0.82 (3H, d, J = 8.7 Hz, H-27), 0.80 (3H, d, J = 8.7 Hz, H-26), 0.66 (3H, s, 18-Me).

Compound 2, colorless crystal (EtOH), mp, 183-185°C; EIMS m/z 152 [M]⁺, 135, 122, 109, 92, 84, 56; ¹H-NMR (CDCl₃): δ 8.42 (2H, dd, J = 8.6, 2.0 Hz, H-2, H-6), 7.06 (2H, dd, J = 8.6, 2.0 Hz, H-3, H-5), 3.68 (3H, s, OMe); ¹³C-NMR (CDCl₃): 125.1 (C-1), 132.3 (C-2, C-6), 114.1 (C-3, C-5), 163.6 (C-4), 168.8 (C-7), 55.4 (OMe).

Compound 3, white powder; ESIMS m/z 599 [M + Na]⁺, 576 [M]⁺, 413 [M-glu-1]⁺; EIMS m/z 414 [M]⁺, 396, 381, 354, 329, 303, 273, 255, 231, 213; ¹H-NMR (DMSO-d₆):

δ 5.34 (1H, m, H-6), 4.92 (1H, d, J = 8.0 Hz, Glu, H-1'), 3.50 (1H, m, H-3), 0.99 (3H, s, H-19), 0.92 (3H, d, J = 8.1 Hz, H-21), 0.84 (3H, t, J = 7.7 Hz, H-29), 0.82 (3H, d, J = 8.7 Hz, H-27), 0.80 (3H, d, J = 8.7 Hz, H-26), 0.66 (3H, s, 18-Me); ¹³C-NMR (DMSO-d₆): 36.7 (C-1), 29.3 (C-2), 76.8 (C-3), 38.2 (C-4), 140.3 (C-5), 120.1 (C-6), 31.3 (C-7, C-8), 50.5 (C-9), 36.1 (C-10), 22.5 (C-11), 41.7 (C-12), 45.0 (C-13), 56.1 (C-14), 24.8 (C-15), 27.7 (C-16), 55.3 (C-17), 11.6 (C-18), 19.0 (C-19), 35.4 (C-20), 18.8 (C-21), 33.0 (C-22), 25.3 (C-23), 49.5 (C-24), 28.5 (C-25), 19.6 (C-26), 12.9 (C-27), 23.8 (C-28), 12.0 (C-29), 100.6 (Glu, C-1'), 73.3 (C-2'), 76.8 (C-3'), 69.9 (C-4'), 76.6 (C-5'), 60.9 (C-6').

Compound 4, orange needles (CHCl₃), mp, 208-210°C; UV (EtOH) λ_{max} (nm): 219, 250, 272, 304, 322, 393; ESI-MS: m/z 621 [2M + Na]⁺, 599 [2M + H]⁺, 300 [M + H]⁺; ¹H-NMR (DMSO-d₆, 400 MHz): 14.7 (1H, s, OH), 8.32 (1H, d, J = 8.9 Hz, H-8), 7.79 (1H, dd, J = 8.7, 7.6 Hz, H-6), 7.76 (1H, d, J = 8.7 Hz, H-5), 7.64 (1H, d, J = 8.9, 7.6 Hz, H-7), 6.05 (2H, s, O-CH₂-O), 4.03 (3H, s, OCH₃), 3.86 (3H, s, N-CH₃); ¹³C-NMR (DMSO-d₆, 100 MHz): 175.9 (C-9), 145.1 (C-3), 144.4 (C-5a), 137.3 (C-4a), 136.6 (C-1), 135.4 (C-2), 133.5 (C-6), 130.1 (C-4), 126.3 (C-8), 124.1 (C-8a), 122.1 (C-7), 117.1 (C-5), 113.5 (C-1a), 102.7 (O-CH₂-O), 61.2 (4-OMe), 42.0 (N-Me).

Compound 5, red needles (CHCl₃), mp, 234-236°C; UV (EtOH) λ_{max} (nm): 222, 250, 277, 313, 423; ESI-MS: m/z 322 [M + Na]⁺, 300 [M + H]⁺; EIMS m/z 299 [M]⁺ (83), 285 (8), 284 (100), 254 (31), 170 (16), 158 (31), 115 (17); ¹H-NMR (DMSO-d₆, 400 MHz): 14.9 (1H, s, OH), 8.37 (1H, d, J = 7.9 Hz, H-8), 7.82 (1H, dd, J = 8.4, 7.7 Hz, H-6), 7.78 (1H, d, J = 7.7 Hz, H-5), 7.67 (1H, d, J = 8.4, 7.9 Hz, H-7), 5.99 (2H, s, O-CH₂-O), 4.01 (3H, s, OCH₃), 3.97 (3H, s, N-CH₃); ¹³C-NMR (DMSO-d₆, 100 MHz): 176.2 (C-9), 147.9 (C-1), 143.5 (C-3), 144.1 (C-5a), 133.9 (C-6), 133.1 (C-2), 132.1 (C-4), 127.5 (C-4a), 127.3 (C-8), 122.5 (C-8a), 121.8 (C-7), 116.2 (C-5), 115.1 (C-1a), 102.4 (O-CH₂-O), 61.1 (2-OMe), 40.6 (N-Me).

Compound 6, light yellow needles (CHCl₃), mp, 176-178°C; UV (EtOH) λ_{max} (nm): 215, 251, 271, 302, 404; EIMS m/z 313 [M]⁺ (75), 299 (28), 298 (100), 284 (21), 280 (10), 270 (50), 268 (13), 255 (10), 254 (8), 169 (8); ¹H-NMR (DMSO-d₆, 400 MHz): δ_H 8.53 (1H, dd, J = 9.0 Hz, H-8), 7.90 (1H, dd, J = 8.7, 7.7 Hz, H-6), 7.75 (1H, d, J = 8.7 Hz, H-5), 7.58 (1H, d, J = 9.0, 7.7 Hz, H-7), 6.08 (2H, s, O-CH₂-O), 4.01, 3.98 (2 CH₂H, s, 2OCH₃), 3.97 (3H, s, N-CH₃); ¹³C-NMR (DMSO-d₆, 100 MHz): 176.1 (C-9), 149.8 (C-1), 143.8 (C-3), 143.1 (C-5a), 134.1 (C-6), 133.6 (C-2), 131.9 (C-4), 127.1 (C-4a), 127.0 (C-8), 122.9 (C-8a), 121.6 (C-7), 115.5 (C-5), 114.3 (C-1a), 102.3 (O-CH₂-O), 62.4 (1-OMe), 61.4 (2-OMe), 39.2 (N-Me).

Compound 7, light yellow prismatic crystal (EtOEt), mp, 121-122°C; UV (EtOH) λ_{\max} (nm): 221, 252, 276, 302, 325, 399; ESI-MS: m/z 649 [2M + Na]⁺, 336 [M + Na]⁺, 314 [M + H]⁺; MS m/z : 313 [M]⁺ (100), 298 (59), 285 (25), 280 (28), 271 (13), 270 (74), 269 (9), 268 (25), 255 (16), 253 (10), 252 (50), 240 (9), 224 (14), 212 (7); ¹H-NMR (DMSO-d₆, 400 MHz): δ_{H} 8.26 (1H, d, J = 8.7 Hz, H-8), 7.89 (1H, dd, J = 8.5, 7.4 Hz, H-6), 7.81 (1H, d, J = 8.5 Hz, H-5), 7.63 (1H, d, J = 8.7, 7.4 Hz, H-7), 6.02 (2H, s, O-CH₂-O), 4.05, 3.84 (2 CH₃H, s, 2OCH₃), 3.82 (3H, s, N-CH₃); ¹³C-NMR (DMSO-d₆, 100 MHz): 176.3 (C-9), 145.6 (C-3), 144.7 (C-5a), 137.6 (C-1), 137.2 (C-4a), 135.7 (C-2), 133.3 (C-6), 129.4 (C-4), 126.1 (C-8), 123.8 (C-8a), 121.7 (C-7), 116.9 (C-5), 113.9 (C-1a), 103.1 (O-CH₂-O), 61.4 (1-OMe), 61.1 (4-OMe), 42.2 (N-Me).

Compound 8, yellow prismatic crystal (Acetone), mp 133-134°C; UV λ_{\max} (nm): 203 (sh), 218, 268, 310 (sh), 397; EIMS m/z : 329 [M]⁺, 314 (100), 270, 256, 228, 157, 149, 143, 77; ¹H-NMR (DMSO-d₆, 400 MHz): δ_{H} 8.41 (1H, d, J = 8.9 Hz, H-8), 7.85 (1H, dd, J = 8.4, 7.7 Hz, H-6), 7.73 (1H, d, J = 8.4 Hz, H-5), 7.59 (1H, d, J = 8.9, 7.7 Hz, H-7), 4.11, 4.02, 3.97, 3.80 (4 s, 4OCH₃), 3.91 (3H, s, NCH₃); ¹³C-NMR (DMSO-d₆, 100 MHz): 176.4 (C-9), 152.6 (C-1), 149.7 (C-3), 144.9 (C-5a), 139.4 (C-4), 138.5 (C-4a), 136.7 (C-2), 134.1 (C-6), 126.7 (C-8), 123.5 (C-8a), 121.9 (C-7), 117.2 (C-5), 114.2 (C-1a), 62.2 (1-OMe), 61.9 (2-OMe), 61.9 (3-OMe), 62.1 (4-OMe), 42.1 (N-Me).

Identification of the isolated compounds [Figure 1] was based on comparison of UV, MS, ¹H-NMR and ¹³C-NMR data to previously reported ones as β -sitosterol 1,^[8] 4-methoxy benzoic acid 2,^[8] daucosterol 3^[9] and five acridone alkaloids, normelicopidine 4,^[10] normelicopine 5,^[11] melicopine 6,^[11] melicopidine 7^[11] and melicopine 8.^[12] It is noteworthy to mention that all these acridone alkaloids were isolated from this plant for the first time.

Cytotoxicity assay

Compounds 4-8 were added to Falcon 384 well black/clear tissue treated assay plates containing 3000 adherent cells/well (HEK293) in an assay volume of 45 μ L. The plates were incubated for 72 h at 37°C and 5% CO₂. After incubation the supernatant was aspirated from the wells and 40 μ L of 10% Alamar Blue added per well. Plates were incubated for a further 5-6 h and measured for fluorescence at 535 nm excitation and 590 nm emission using a VICTOR II (PerkinElmer, Waltham, MA). The % inhibition of cell proliferation was calculated using Dimethylsulfoxide (DMSO) and 10 μ M 5-Fluorouracil control data. IC₅₀ values were obtained by plotting % inhibition against log dose using the Prism4




Figure 1: Structures for compounds 4-8

graphing package and nonlinear regression with variable slope plot.

Antimalarial assay

Compounds 4-8 were incubated in the presence of 2 or 3% parasitemia (3D7 or Dd2) and 0.3% hematocrit in a total assay volume of 50 μ L, for 72 h at 37°C and 5% CO₂ in poly-Dlysine-coated Cell Carrier imaging plates. After incubation plates were stained with DAPI in the presence of saponin and Triton X-100 and incubated for a further 5 h at RT in the dark before imaging on the Evotec OPERA High Throughput Screening confocal imaging system (PerkinElmer, Waltham, MA). The digital images obtained were analyzed using the Perkin-Elmer Acapella spot detection software, where fluorescent spots that fulfilled the criteria established for a stained parasite were counted. The % inhibition of parasite replication was calculated using DMSO and 2 μ M artemisinin control data.

RESULTS AND DISCUSSION

Cytotoxicity [Table 1] of five acridone alkaloids isolated from this herb was evaluated against two prostate cancer cell lines PC-3M and LNCaP. Preliminary toxicity toward human cells was investigated using a human embryonic kidney cell line (HEK293). The results revealed that all five compounds exhibited potential cytotoxicity against PC-3M and LNCaP with IC₅₀ value range between 12 μ g/mL and 65 μ g/mL, and showed no cytotoxicity against HEK293 up to 100 μ g/mL. Compound 4 was the most active against PC-3M and LNCaP with IC₅₀ values of 12.5 and 21.1 μ g/mL respectively.

Antimalarial activity [Table 2] for the acridone alkaloids has also been tested against a chloroquine-sensitive (3D7) and a chloroquine-resistant (Dd2) *Plasmodium falciparum* strain. All Compounds displayed IC₅₀ values in a range of 18-42 μ g/mL

Table 1: Cytotoxic activities of compounds 4-8 (IC₅₀, ug/mL; mean±SD, n=3)

Compound	Cell lines		
	PC-3M	LNCaP	HEK293
4	12.5±1.9	21.1±2.6	>100
5	36.3±2.4	44.6±2.7	>100
6	47.9±3.6	37.8±3.4	>100
7	36.1±2.2	28.9±3.1	>100
8	51.4±4.3	64.2±4.7	ND
5-FUa	7.5±1.8	9.3±1.5	14.7±0.4

5-FU: 5-fluorouracil was used as positive control; ND: Not determined; SD: Standard deviation. PC-3M and LNCaP: two prostate cancer cell lines; IC₅₀: 50% inhibitory concentration

Table 2: Antimalarial and cytotoxic activity for compounds 4-8 (IC₅₀, ug/mL; mean±SD, n=3)

Compound	IC ₅₀ (ug/mL)			Selectivity index
	Dd2	3D7	HEK293	
4	18.9±2.2	NT	>100	>5.0 (Dd2)
5	25.9±2.6	31.2±3.3	>100	>3.8 (Dd2), >3.2 (3d7)
6	33.7±1.9	29.7±2.8	>100	>2.9 (Dd2), >3.3 (3d7)
7	21.6±2.5	25.5±1.9	>100	>4.6 (Dd2), >3.9 (3d7)
8	41.2±2.4	37.7±3.1	>100	>2.4 (Dd2), >2.6 (3d7)
Artemisinin	0.0056	0.0056	0% @ 0.56	>100 (Dd2, 3D7)
Chloroquine	0.067	0.013	0% @ 1.03	>16 (Dd2), >80 (3D7)

NT: Not tested; SD: Standard deviation

against 3D7 and Dd2. Compound 4 showed the strongest activity against Dd2 with IC₅₀ value of 18.9 ug/mL.

CONCLUSION

Acridone alkaloids are abundant in the Rutaceae family, especially occurring in citrus plants, and many of them showed interesting pharmaceutical properties such as anticancer, antitumor, antiviral, antimalarial, and antiallergic activities. Our investigation on the root bark of *Z. simullans* Hance has led to the isolation of five acridone alkaloids, normelicopidine, normelicopine, melicopine, melicopidine, and melicopicine from this plant for the first time. All tested compounds exhibited certain cytotoxic and antimalarial activities *in vitro*, and compound 4 was the most active against PC-3M, LNCaP and Dd2 with IC₅₀ values of 12.5, 21.1 and 18.9 ug/mL respectively.

ACKNOWLEDGMENTS

The work was financially supported by the Talented Faculty Foundation of South-Central University for Nationalities (YZ211002 to X.Z. Yang) and National Natural Science Foundation of China (81102798 to X.Z. Yang).

REFERENCES

- Seidemann J. World Spice Plants: Economic Usage, Botany, Taxonomy. Berlin: Springer-Verlag; 2005. p. 399-402.
- Xu GJ, He HX, Xu LS, Jin RY. Zhonghua Bencao. Vol. 7. Beijing: Chinese Medicinal Science and Technology Publishing House; 1999. p. 506.
- Chen IS, Wu SJ, Lin YC, Tsai IL, Seki H, Ko FN, et al. Dimeric 2-quinolone alkaloid and antiplatelet aggregation constituents of *Zanthoxylum simulans*. Phytochemistry 1994a; 36:237-9.
- Chen IS, Wu SJ, Tsai IL. Chemical and bioactive constituents from *Zanthoxylum simulans*. J Nat Prod 1994;57:1206-11.
- Chen IS, Wu SJ, Leu YL, Tsai IW, Wu TS. Alkaloids from root bark of *Zanthoxylum simulans*. Phytochemistry 1996;24:217-9.
- Chen IS, Tsai IW, Teng CM, Chen JJ, Chang YL, Ko FN, et al. Pyranoquinoline alkaloids from *Zanthoxylum simulans*. Phytochemistry 1997;46:525-9.
- Yang YP, Cheng MJ, Teng CM, Chang YL, Tsai IL, Chen IS. Chemical and anti-platelet constituents from formosan *Zanthoxylum simulans*. Phytochemistry 2002;61:567-572.
- Pan ZH, Luo L, Li ZQ, Zhang LL, Dong Q, Ma GY. The chemical constituents from *Pittosporum podocarpum* Gagnep. J Yunnan Univ (Natural Sciences) 2012;34:450-3.
- Yang W, Tang CM, Li X, Zhou Y, Wang L, Li L. Study on the chemical constituents of *Myrica esculenta*. J Yunnan Univ (Natural Sciences) 2011;33:453-7.
- Tillequin F, Baudouin G, Koch M, Sévenet T. Plants of new caledonia LXIV, alkaloids of leaves of *Bauerella simplicifolia* ssp. neo-scotica. J Nat Prod 1980;43:498-502.
- Funayama S, Cordell GA. Chemistry of acronyline IV. Minor constituents of acronine and the phytochemistry of the genus *Acronychia*. J Nat Prod 1984;47:285-291.
- Lwande W, Gebreyesus T, Chapya A, Macfoy C, Hassanali A, Okech M. 9-Acridone insect antifeedant alkaloids from *teclea trichocarpa* bark. Int J Trop Insect Sci 1983;4:393-5.

Cite this article as: Wang C, Wan J, Mei Z, Yang X. Acridone alkaloids with cytotoxic and antimalarial activities from *Zanthoxylum simulans* Hance. Phcog Mag 2014;10:73-6.

Source of Support: The work was financially supported by the Talented Faculty Foundation of South-Central University for Nationalities (YZ211002 to X.Z. Yang) and National Natural Science Foundation of China (81102798 to X.Z. Yang), **Conflict of Interest:** None declared.