

RESEARCH ARTICLE

Renin-Angiotensin System Inhibitors, Type 2 Diabetes and Fibrosis Progression: An Observational Study in Patients with Nonalcoholic Fatty Liver Disease

Serena Pelusi¹, Salvatore Petta², Chiara Rosso³, Vittorio Borroni¹, Anna Ludovica Fracanzani¹, Paola Dongiovanni¹, Antonio Craxi², Elisabetta Bugianesi³, Silvia Fargion¹, Luca Valenti^{1*}

1 Internal Medicine and Metabolic Diseases, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Università degli Studi di Milano, Milano, Italy, **2** Gastroenterology, Università di Palermo, Palermo, Italy, **3** Gastroenterology, Dept. of Medical Sciences, Università di Torino, Torino, Italy

* luca.valenti@unimi.it



OPEN ACCESS

Citation: Pelusi S, Petta S, Rosso C, Borroni V, Fracanzani AL, Dongiovanni P, et al. (2016) Renin-Angiotensin System Inhibitors, Type 2 Diabetes and Fibrosis Progression: An Observational Study in Patients with Nonalcoholic Fatty Liver Disease. PLoS ONE 11(9): e0163069. doi:10.1371/journal.pone.0163069

Editor: Vincent Wong, The Chinese University of Hong Kong, HONG KONG

Received: August 6, 2016

Accepted: September 1, 2016

Published: September 20, 2016

Copyright: © 2016 Pelusi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: LV was funded by MyFirst AIRC (Italian Association for Cancer Research) grant n.16888 and Ricerca corrente Fondazione IRCCS Ca' Granda Milano, Molecular Medicine Grant Fondazione-INGM. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Background

The clinical determinants of fibrosis progression in nonalcoholic fatty liver disease (NAFLD) are still under definition.

Aim

To assess the clinical determinants of fibrosis progression rate (FPR) in NAFLD patients with baseline and follow-up histological evaluation, with a special focus on the impact of pharmacological therapy.

Methods

In an observational cohort of 118 Italian patients from tertiary referral centers, liver histology was evaluated according to Kleiner. Independent predictors of FPR were selected by a stepwise regression approach.

Results

Median follow-up was 36 months (IQR 24–77). Twenty-five patients (18%) showed some amelioration, 63 (53%) had stability, 30 (25%) had progression of fibrosis. Patients with non-alcoholic steatohepatitis (NASH) had similar demographic and anthropometric features, but a higher prevalence of type 2 diabetes (T2D; $p = 0.010$), and use of renin-angiotensin axis system (RAS) inhibitors ($p = 0.005$). Fibrosis progression was dependent of the length of follow-up, and was associated with, but did not require, the presence of NASH ($p < 0.05$). Both fibrosis progression and faster FPR were independently associated with higher APRI score at follow-up, absence of treatment with RAS inhibitors, and T2D diagnosis at baseline

Competing Interests: The authors have declared that no competing interests exist.

($p < 0.05$). There was a significant interaction between use of RAS inhibitors and T2D on FPR ($p = 0.002$). RAS inhibitors were associated with slower FPR in patients with ($p = 0.011$), but not in those without ($p = \text{NS}$) T2D.

Conclusions

NASH is not required for fibrosis progression in NAFLD, whereas T2D seems to drive fibrogenesis independently of hepatic inflammation. Use of RAS inhibitors may contrast fibrosis progression especially in high-risk patients affected by T2D.

Introduction

Nonalcoholic fatty liver disease (NAFLD) is commonly held as the hepatic manifestation of obesity and insulin resistance. Due to the worldwide epidemics of obesity and type 2 diabetes (T2D), NAFLD is projected to become the leading cause of hepatocellular carcinoma and end-stage liver disease within the next ten years[1]. Despite NAFLD affects nearly one third of the population, progressive liver disease remains a relatively rare complication of this condition[1]. Cross-sectional studies have identified severity of overweight, T2D, muscle fitness, dietary factors, lack of use of lipid lowering drugs such as statins, and genetic predisposition as risk factors for advanced disease [2–5]. However, the clinical determinants of progression of fibrosis, the main determinant of liver-related outcomes and overall mortality[6,7], are still under definition. Indeed, data from prospective studies are still very limited[8,9]. Overall evidence suggests that when steatosis is associated with hepatocellular damage and necroinflammation, that is nonalcoholic steatohepatitis (NASH), higher AST/ALT ratio, and in the presence of hyperglycemia, fibrosis progression rate (FPR) is faster[8–10]. Yet, some individuals with simple steatosis have fast-progressing disease, especially when gain weight or develop T2D [9,11]. Furthermore, arterial hypertension has also been associated with faster FPR[12]. This suggests that neuro-hormonal alterations associated with this condition, and in particular activation of the renin-angiotensin system (RAS), directly favors steatosis, inflammation and fibrogenesis via enhanced activation of hepatic stellate cells, whereas RAS inhibits contrast this process[13–20]. In keeping, RAS inhibitors such as ACE-inhibitors or angiotensin receptor blockers have been associated with improvement of liver damage[21], even if evidence is controversial[22]. Furthermore, in cross-sectional studies RAS inhibition protected from severe fibrosis in patients with hypertension and NAFLD[23], and was associated with reduced liver stiffness in patients with chronic kidney disease [24]

Aim of this study was therefore to assess the clinical determinants of FPR in an ethnically homogeneous cohort of Italian patients with histological diagnosis of NAFLD, with a special focus on the impact of pharmacological therapy.

Methods

Patients

In the study retrospective data collected from 118 consecutive patients from Italian ancestry with clinical and histological diagnosis of NAFLD were prospectively evaluated. Patients were followed-up at three tertiary referral centers in Italy (Milan, $n = 67$, 57%, Palermo, $n = 32$, 27%, and Turin, $n = 19$, 16%), for whom a baseline and a follow-up liver biopsy and clinical data were available between January 1992 and June 2015.

In all patients other liver diseases were ruled out by standard assessment [2,25], and alcohol intake (evaluated by a questionnaire) had to be lower than 30/20 g/day in males/females, respectively. Patients with decompensated cirrhosis, hepatocellular carcinoma, and current use of steatosis inducing drugs were also excluded.

In all subjects, first biopsy was performed for suspected NASH in the presence of persistently elevated liver enzymes, or a long history of NAFLD associated with severe insulin resistance. Follow-up control biopsy was routinely offered to all compliant patients at five years, or indicated when alterations in the clinical picture or imaging suggested progressive liver disease. We also included patients randomized to iron depletion [26] or vitamin D supplementation (<http://www.webaisf.org/studi-e-ricerche/studi-in-corso.aspx>) vs. lifestyle changes alone in open label trials, as these treatments were not demonstrated to influence fibrosis progression. Patients randomized to active arms in pharmacological studies, where the investigational product was shown to improve liver histology, or who underwent bariatric surgery procedures between the two biopsies (n = 13) were excluded.

The study was carried out in accordance with the principles of the Helsinki Declaration, and with local and national laws. Approval was obtained from the hospital Internal Review Boards and Ethics Committees of the Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico Milano, Azienda Ospedaliera Universitaria Citta' della Salute e della Scienza Torino and Azienda Ospedaliera Universitaria Policlinico Palermo and written informed consent was obtained from all patients. Clinical and laboratory assessment is described in details in the Supplementary methods.

Histological analysis

Slides were coded and read by one expert pathologist at each center, who was unaware of patients' identity and history. A minimum 15mm-length of the biopsy specimen or the presence of at least 10 complete portal tracts was required [27]. Clinically significant steatosis was defined as steatosis involving $\geq 5\%$ of hepatocytes [28]. Diagnosis of NASH was based on the presence of steatosis with both lobular necroinflammation and ballooning [28,29]. Disease activity and fibrosis stage were assessed according to the NAFLD activity score (NAS) and staging [28]. We previously observed a good correlation for liver fibrosis assessment among the centers involved in the study [30].

Statistical analysis

For descriptive statistics, continuous traits were summarized as means \pm SD. Highly skewed variables, were summarized as medians and interquartile range, and log-transformed before analysis. Categorical variables are shown as percentages. Baseline and follow-up clinical features of patients were compared by chi-square and paired t-test, as required. Independent predictors of fibrosis progression (increase in at least one stage) were determined by logistic regression, considering as independent variables those significant at univariate analysis. FPR was calculated by taking the ratio between the difference of fibrosis stage and the time (months) between the baseline and follow-up biopsy, and it was treated as a continuous variable. A multivariate regression model with a stepwise regression procedure was set to identify the strongest predictors of FPR, among all variables considered in the study (enlisted in supplementary material). A significance level of 0.1 was defined to allow a variable into the model, and a significance level of 0.25 was defined for a variable to stay into the model. A generalized linear model was then fit to examine the independent predictors of FPR, excluding highly correlated variables to avoid collinearity. In this final model, product terms between variables were evaluated to investigate the interaction between risk factors on FPR.

Statistical analyses were carried out with JMP 12.0 (SAS Institute, Cary, NC) and SPSS 21.0 (IBM, Burbank, NJ). A two-sided P value <0.05 was considered statistically significant.

Results

Study cohort and clinical evolution at follow-up

The baseline clinical features of patients included in the study are shown in Table 1, left column. They were mostly middle-aged men or post-menopausal women, overweight or obese, with a high prevalence of metabolic alterations defining metabolic syndrome and/or altered liver enzymes (Table 1). Forty-two % had histological NASH.

At baseline liver biopsy, 16% of patients were on statins, 22% on RAS inhibitors (14 on ACE-inhibitors and 12 on angiotensin receptor blockers). Indication for RAS inhibitors was treatment of hypertension and/or micro-albuminuria in T2D in 14, and treatment of hypertension without T2D in 12. RAS inhibitors were not indicated for prevention of liver fibrosis progression in this cohort.

Table 1. Clinical features of 118 Italian patients with NAFLD, who underwent a follow-up liver biopsy.

Clinical features	Baseline	Follow-up	p value
Sex, F	45 (38)	45 (38)	1.00
Age, years	47±12	51±11	<0.001
BMI, Kg/m ²	30.6±6.6	29.0±7.2	0.006
T2D, yes	29 (25)	32 (27)	0.66
Glucose, mg/dl	98±25	102±25	0.13
Total cholesterol, mg/dl	194±44	189±43	0.066
HDL cholesterol, mg/dl	47±14	49±13	0.054
Triglycerides, mg/dl	111 {78–161}	111 {73–154}	0.078
Arterial hypertension, yes	38 (32)	54 (46)	0.033
ALT, IU/ml	50 {20–83}	40 {21–60}	0.002
AST, IU/ml	33 {23–51}	27 {21–36}	0.077
GGT, IU/ml	45 {28–80}	35 {19–63}	0.45
Ferritin (ng/mL)	234 {88–506}	138 {73–334}	0.012
Platelets (x10 ⁹ /L)	224 ± 71	224 ± 67	0.9
NASH, yes	49 (42)	47 (40)	0.79
APRI score	0.7 ± 1.7	0.4 ± 0.3	0.11
FIB4 score	1.3 ± 1.1	1.3 ± 0.9	0.82
NFS	-1.7 ± 1.7	-1.6 ± 1.6	0.79
RAS inhibitors, yes	26 (22)	36 (31)	0.14
Beta-blockers, yes	14 (12)	19 (16)	0.35
Calcium-antagonists, yes	12 (10)	11 (9)	0.81
Diuretics, yes	9 (8)	9 (8)	1.00
Metformin, yes	22 (19)	30 (26)	0.21
Statins, yes	16 (14)	28 (24)	0.045
Omega-3, yes	7 (6)	8 (7)	0.79
Vitamin E, yes	4 (3)	6 (5)	0.52
Iron depletion, yes	0	12 (10)	<0.001
Length of follow-up, months	36 {24–77}		-

Data are shown as mean±SD, frequency (%), median {IQR}, as required. BMI: body mass index; T2D: type 2 diabetes; RAS: renin angiotensin system. Less than three patients (per drug class) were on glitazones, fibrates, and GLP-1 agonists/DPP-4 inhibitors.

doi:10.1371/journal.pone.0163069.t001

Median follow-up was 36 months (IQR 24–77, range 7–196), for a total of 6,509 months. Clinical features of patients at follow-up are shown in [Table 1](#), right column. Patients lost on average 1.6 Kg/m² of body mass ($p = 0.006$), which was associated with reduced ALT levels ($p = 0.002$). The prevalence of arterial hypertension ($p = 0.033$), but not that of T2D, increased at follow-up.

At the end of the observation there was an increase in the prevalence of treatment with statins ($p = 0.045$). An additional 10% of patients had started RAS inhibitors (7 ACE-inhibitors and 3 angiotensin receptor blockers; 2 with T2D, and 8 with hypertension without T2D; $p = \text{NS}$). Twelve patients with baseline hyperferritinemia underwent iron depletion by phlebotomy during follow-up ($p < 0.001$).

Fibrosis evolution during follow-up

The prevalence of histological NASH did not change significantly at follow-up ([Table 1](#); $p = \text{NS}$), although average NAS score decreased (median 4, IQR 2–5 at baseline vs. 3, IQR 2–5 at follow-up; $p = 0.026$). Non-invasive fibrosis scores did not significantly change at follow-up ($p = \text{NS}$; [Table 1](#)).

Clinical features of patients stratified by the presence of baseline NASH are presented in [Table 2](#). Patients with NASH had a higher prevalence of T2D ($p = 0.010$), use of RAS inhibitors ($p = 0.005$), and in the subgroup of patients for whom data were available, of prevalence of *PNPLA3* 148M/M risk genotype ($p = 0.023$).

Evolution of liver fibrosis according to baseline stage is presented in [Table 3](#), upper panel. Of 118 patients, 25 (18%) showed some amelioration, 63 (53%) had stability, 30 (25%) had progression of fibrosis. Of note, 5 patients had progression to cirrhosis. Mean FPR was -0.002 ± 0.040 . Evolution of fibrosis in patients stratified according to NASH at baseline is presented in [Table 3](#), middle and bottom panels, respectively. Fibrosis stage was more severe in patients with NASH at baseline ($p < 0.001$). FPR was non-significantly lower in patients with than in those without baseline NASH (-0.011 ± 0.054 vs. $+0.004 \pm 0.033$; $p = 0.056$), and it was not associated with presence of necroinflammation without NASH ($p > 0.5$). No single baseline histological feature of liver damage was able to predict FPR ([S1 Table](#)).

Clinical predictors of fibrosis progression

Baseline and follow-up clinical features of patients without baseline cirrhosis ($n = 10$) stratified by progression status are presented in [Table 4](#). Progression of fibrosis was associated with length of follow-up ($p = 0.027$), lower HDL, absence of use of RAS inhibitors at baseline, AST and ALT levels at baseline and follow-up, APRI score and NASH at follow-up ($p < 0.05$). After correction for duration of observation ([Table 4](#)), progression was associated with NASH at baseline and follow-up, T2D at baseline and metformin use (likely a proxy of more severe T2D) at follow-up, and AST and APRI score at follow-up ($p < 0.05$). There was a non-significant trend for a protective effect of use of RAS inhibitors. Among patients without NASH at baseline ($n = 69$), 6 of 15 (43%) progressors developed NASH at follow-up, vs. 8/46 (15%) of non-progressors ($p = 0.044$).

The association of the changes of clinical variables during follow-up with fibrosis progression is shown in [S2 Table](#). None was associated with progression, even if at unadjusted analysis development of new hypertension was associated with progression (8/30, 27% vs. 7/78, 9%; $p = 0.028$). The noninvasive independent predictors of progression at multivariate logistic regression analysis are shown in [S3 Table](#). Duration of observation, lack of use of RAS inhibitors, baseline NASH, T2D, and follow-up APRI score were associated with the likelihood of progression ($p < 0.05$).

Table 2. Clinical features associated with presence of NASH at baseline evaluation.

Clinical features	NASH (n = 49)	Non-NASH (n = 69)	p value
Sex, F	21 (43)	24 (35)	0.37
Age, years	48±12	47±12	0.49
BMI, Kg/m ²	30.4±4.1	31.0±8.0	0.41
T2D, yes	18 (37)	11 (16)	0.010
Glucose, mg/dl	108±34	92±13	0.002
Total cholesterol, mg/dl	190±40	197±46	0.33
HDL cholesterol, mg/dl	47±15	48±15	0.67
Triglycerides, mg/dl	135±77	130±72	0.71
Arterial hypertension, yes	20 (41)	18 (26)	0.091
ALT, IU/ml	57 {41–95}	47 {24–77}	0.86
AST, IU/ml	37 {27–62}	31 {21–40}	0.60
GGT, IU/ml	54 {31–93}	42 {26–72}	0.45
Ferritin (ng/mL)	196 {51–408}	292 {75–523}	0.11
Platelets (x10 ⁹ /L)	237 ± 74	217 ± 70	0.16
NASH, yes	49 (42)	47 (40)	0.79
APRI score	0.5 ± 0.3	0.8 ± 2.2	0.37
FIB4 score	1.3 ± 0.8	1.3 ± 1.3	0.82
NFS	-1.7 ± 1.5	-1.7 ± 1.5	0.73
RAS inhibitors, yes	17 (35)	9 (13)	0.005
Beta-blockers, yes	7 (14)	7 (10)	0.49
Calcium-antagonists, yes	8 (16)	4 (6)	0.062
Diuretics, yes	4 (8)	5 (7)	0.85
Metformin, yes	16 (33)	6 (9)	0.001
Statins, yes	8 (16)	8 (12)	0.59
Omega-3, yes	4 (8)	3 (4)	0.39
Vitamin E, yes	4 (3)	6 (5)	0.52
Iron depletion, yes	3 (6)	9 (13)	0.22
PNPLA3 148 M/M	12/27 (44)	13/62 (21)	0.023
FPR, stage/month	-0.01 ± 0.05	+0.004 ± 0.03	0.080
Follow-up, months	30 {7–144}	38 {9–196}	0.010

Data are shown as mean±SD, frequency (%), median {IQR}, as required. BMI: body mass index; T2D: type 2 diabetes; RAS: renin angiotensin system; FPR: fibrosis progression rate.

doi:10.1371/journal.pone.0163069.t002

Independent predictors of FPR

The independent predictors of FPR are shown in Table 5. Among variables selected by stepwise regression, faster FPR was associated with higher APRI score at follow-up (p = 0.005), absence of treatment with RAS inhibitors (p = 0.009), T2D diagnosis at baseline (p = 0.025). There was a significant interaction between use of RAS inhibitors and T2D (p = 0.002).

The impact of RAS inhibitors use on FPR in patients stratified by the presence of T2D at diagnosis is shown in Fig 1. RAS inhibitors were associated with slower FPR in patients with (p = 0.011), but not in those without (p = 0.52) T2D.

Discussion

In this study, we evaluated the clinical determinants of FPR in a relatively large prospective cohort of 118 Italian patients with histological NAFLD, with a special focus on the impact of

Table 3. Evolution of liver fibrosis by baseline disease stage in 118 Italian patients with NAFLD.

		Overall					
		Follow-up					Total =
Baseline		Stage 0	Stage 1	Stage 2	Stage 3	Stage 4	
	Stage 0	21 (18)	8 (7)	4 (3)	3 (2)	0	36 (31)
	Stage 1	6 (5)	21 (18)	6 (5)	2 (2)	0	35 (30)
	Stage 2	1 (1)	6 (5)	9 (8)	2 (2)	2 (2)	20 (17)
	Stage 3	1 (1)	5 (4)	3 (2)	5 (4)	3 (2)	17 (14)
	Stage 4	0	1 (1)	2 (2)	0	7 (6)	10 (8)
Total =		29 (25)	41 (35)	24 (20)	12 (10)	12 (10)	118
		NASH					
		Follow-up					Total =
Baseline		Stage 0	Stage 1	Stage 2	Stage 3	Stage 4	
	Stage 0	1 (2)	3 (6)	2 (4)	1 (2)	0	7 (14)
	Stage 1	1 (2)	6 (12)	3 (6)	1 (2)	0	11 (22)
	Stage 2	1 (2)	3 (6)	5 (10)	2 (2)	1 (2)	12 (25)
	Stage 3	0	4 (8)	2 (4)	4 (8)	2 (4)	12 (25)
	Stage 4	0	1 (2)	2 (4)	0	4 (8)	7 (14)
Total =		3 (6)	17 (35)	14 (29)	8 (16)	7 (14)	49
		Non-NASH					
		Follow-up					Total =
Baseline		Stage 0	Stage 1	Stage 2	Stage 3	Stage 4	
	Stage 0	20 (18)	5 (7)	2 (3)	2 (2)	0	29 (42)
	Stage 1	5 (5)	15 (18)	3 (5)	1 (2)	0	24 (35)
	Stage 2	1 (1)	3 (5)	4 (8)	0	1 (2)	8 (12)
	Stage 3	1 (1)	1 (4)	1 (2)	1 (4)	1 (2)	5 (7)
	Stage 4	0	0	0	0	3 (6)	3 (4)
Total =		26 (38)	24 (35)	10 (14)	4 (6)	5 (7)	69

doi:10.1371/journal.pone.0163069.t003

pharmacological therapies. This is especially relevant for the clinical management because fibrosis is the main prognostic indicator in patients with NAFLD [6,7]

In line with previous results[8], we observed that progression of fibrosis did not require the presence of baseline NASH, despite NASH was related to more severe fibrosis at baseline and associated with progression of fibrosis. However, follow-up liver biopsy was not systematically performed in patients without baseline NASH. Therefore, an indication bias may have led to selection of patients with worsening metabolic status, liver enzymes, and noninvasive predictors of liver damage. Indeed, in about half of the cases, progressors developed NASH at follow-up. Notwithstanding, data suggest that preventive strategies and clinical trials should not only focus on individuals with histological severe inflammation and hepatocellular ballooning. It is therefore imperative to identify other risk factors and disease mechanism contributing to disease evolution.

One may be represented by T2D and the severity of insulin resistance, which, in line with data reported in a previous UK study evaluating a comparable number of patients[8], was identified as a predictor of fibrosis progression in the present cohort. There is ample literature on the role of insulin resistance and hyperglycemia in the progression of NAFLD[3,31,32]. However, the mechanisms linking metabolic abnormalities with fibrosis progression independently of NASH and inflammation, even in patients with normal liver enzymes[33], need further clarification. Indeed, this may represent a different form of NAFLD progressing to hepatic

Table 4. Clinical features associated with fibrosis progression at baseline and follow-up evaluation in 108 patients with NAFLD without F4 fibrosis at baseline.

	Non-progressors (n = 78)	Progressors (n = 30)	p value	p value*
Follow-up, months	36 {24–72}	60 {30–120}	0.027	1.00
BASELINE				
Sex, F	27 (35)	11 (37)	0.83	0.28
Age, years	47±11	45±13	0.33	0.87
BMI, Kg/m ²	30.7±8.0	29.9±8.4	0.55	0.75
T2D, yes	15 (19)	8 (27)	0.43	0.034
Glucose, mg/dl	98±27	98±21	0.98	0.15
Total cholesterol, mg/dl	200±47	188±33	0.16	0.12
HDL cholesterol, mg/dl	49±14	43±13	0.026	0.14
Triglycerides, mg/dl	129±70	137±86	0.75	0.88
Arterial hypertension, yes	25 (32)	6 (20)	0.24	0.40
ALT, IU/ml	47 {26–72}	72 {39–116}	0.024	0.17
AST, IU/ml	30 {23–39}	40 {26–55}	0.037	0.35
GGT, IU/ml	44 {25–80}	45 {32–65}	0.55	0.46
Ferritin (ng/mL)	161 {72–504}	335 {191–543}	0.12	0.35
Platelets (x10 ⁹ /L)	229 ± 51	215 ± 75	0.38	0.51
NASH, yes	27 (35)	15 (50)	0.18	0.037
APRI score	0.4 ± 0.3	1.2 ± 3.3	0.072	0.079
FIB4 score	1.1 ± 0.7	1.3 ± 1.3	0.18	0.060
NFS	-1.8 ± 1.5	-1.9 ± 1.8	0.92	0.53
RAS inhibitors, yes	20 (26)	2 (7)	0.028	0.059
Beta-blockers, yes	8 (10)	3 (10)	1.00	0.74
Calcium-antagonists, yes	7 (9)	0	0.19	0.99
Diuretics, yes	6 (8)	1 (3)	0.67	0.69
Metformin, yes	11 (14)	6 (20)	0.55	0.14
Statins, yes	10 (13)	3 (10)	1.00	0.98
Omega-3, yes	5 (6)	1 (3)	1.00	0.64
Vitamin E, yes	2 (3)	1 (3)	1.00	0.92
FOLLOW-UP				
Age, years	52±11	51±11	0.81	0.67
BMI, Kg/m ²	29.0±6.6	27.8±9.0	0.48	0.42
T2D, yes	17 (22)	9 (30)	0.45	0.094
Glucose, mg/dl	98±22	102±24	0.35	0.10
Total cholesterol, mg/dl	191±41	186±48	0.64	0.66
HDL cholesterol, mg/dl	49±12	46±13	0.26	0.37
Triglycerides, mg/dl	111±54	131±78	0.20	0.27
Arterial hypertension, yes	32 (41)	14 (47)	0.66	0.64
ALT, IU/ml	40 {20–52}	53 {29–82}	0.022	0.13
AST, IU/ml	27 {19–33}	35 {24–46}	0.006	0.014
GGT, IU/ml	28 {15–58}	42 {26–62}	0.36	0.29
Ferritin (ng/mL)	119 {70–296}	237 {90–427}	0.077	0.19
Platelets (x10 ⁹ /L)	220 ± 74	231 ± 67	0.49	0.40
NASH, yes	22 (28)	17 (57)	0.008	0.012
APRI score	0.3 ± 0.2	0.5 ± 0.4	0.018	0.008
FIB4 score	1.1 ± 0.7	1.5 ± 1.2	0.16	0.053
NFS	-1.9 ± 1.4	-1.8 ± 1.7	0.82	0.36

(Continued)

Table 4. (Continued)

	Non-progressors (n = 78)	Progressors (n = 30)	p value	p value*
RAS inhibitors, yes	24 (31)	6 (20)	0.34	0.25
Beta-blockers, yes	24 (14)	5 (17)	0.74	0.50
Calcium-antagonists, yes	6 (8)	0	0.18	0.99
Diuretics, yes	6 (8)	1 (3)	0.67	0.69
Metformin, yes	15 (19)	10 (33)	0.14	0.043
Statins, yes	10 (13)	3 (10)	0.60	0.24
Omega-3, yes	2 (3)	3 (10)	0.13	0.54
Vitamin E, yes	2 (3)	1 (3)	1.00	0.37
Iron depletion, yes	11 (14)	1 (3)	0.17	0.14

Data are shown as mean±SD, frequency (%), median {IQR}, as required.

* p value adjusted for duration of observation at logistic regression analysis.

BMI: body mass index; T2D: type 2 diabetes; RAS: renin angiotensin system.

doi:10.1371/journal.pone.0163069.t004

complications by inflammation-independent pathways, for which specific noninvasive biomarkers and therapeutic approaches should be developed.

Remarkably, fibrosis progression was associated with the length of the observation, being more frequent in patients observed for an average 5-years period. This supports the EASL-EASD-EASO guidelines that suggest to consider re-biopsy after this timeframe[29]. We also identified the absolute value of APRI score, not reported in a previous cohort[8], as the most reliable predictor of fibrosis progression. These results need however replication.

An important novel aspect that could be addressed in the present database, but could not be evaluated in a previous meta-analysis and in a recent study [8,9], was the impact of therapies on FPR. A major finding was that patients in treatment with RAS inhibitors had slower FPR, and lower likelihood of fibrosis progression. Furthermore, use of RAS inhibitors resulted independently associated with lack of fibrosis progression also at logistic regression analysis considering variables associated at univariate analysis, providing an independent confirmation of this association by an alternative approach. This is in line with accumulating evidence indicating that angiotensin is involved in hepatic stellate cells activation, while RAS inhibitors contrast fibrogenesis in experimental models[13–20], and their use was associated with protection from fibrosis in cross-sectional studies in patients with NAFLD and kidney disease[23,24]. Interestingly, the protective effect of RAS inhibitors was particularly evident in patients with T2D, but not in patients with arterial hypertension in general. This would suggest that activation of RAS represents a specific feature driving both kidney [34] and hepatic fibrogenesis in T2D.

Table 5. Independent predictors of FPR in 118 Italian patients with NAFLD (including 10 with F4 fibrosis at baseline).

FPR predictor	Estimate±SE	p value
APRI at f-up, per unit	+0.04±0.01	0.005
RAS inhibitors at baseline or f-up	-0.012±0.004	0.009
T2D at baseline, yes	+0.010±0.004	0.025
Beta-blockers at baseline or f-up	-0.008±0.005	0.14
BMI variation, Kg/m ²	+0.001±0.001	0.14
Hypertension at baseline or f-up	+0.006±0.005	0.24

SE: standard error; F-up: follow-up.

doi:10.1371/journal.pone.0163069.t005




Fig 1. Impact of RAS inhibitors on FPR in patients stratified by the presence of T2D at baseline.

doi:10.1371/journal.pone.0163069.g001

Noteworthy, activation of the neuro-vegetative system is associated with induction of RAS activation, and the two have a synergic role in the pathogenesis of arterial hypertension and kidney dysfunction in overweight individuals[35]. More widespread use of RAS inhibitors in patients with baseline NASH may have contributed to the slow FPR observed in this subgroup despite severe histological activity. On the other hand, we could not detect any significant effect of statins [2,36], or iron depletion [37] on FPR.

Limitations of this study include that the sample size did not allow to test the effect of specific therapeutic molecules on FPR, and to test for a dose-response effect. Furthermore, findings may not be extended to other ethnic groups and populations with different genetic and lifestyle risk factors. This study does not report results of a randomized trial, therefore the association of therapeutic approaches with liver disease evolution should be considered as hypothesis generating, and not as a proof of efficacy. In addition, treatment of T2D has recently evolved, but in the present cohort only a few patients were taking oral hypoglycemic drugs other than metformin. Larger collaborative studies are needed to better characterize the impact of new as well as “old” pharmacological treatments on the progression of liver disease in NAFLD.

In conclusion, our data confirm that NASH is not required for fibrosis progression in NAFLD, whereas T2D seems to drive fibrogenesis independently of hepatic inflammation. Most importantly, we could show for the first time in a prospective evaluation of retrospectively collected data by two independent approaches that use of RAS inhibitors may contrast fibrosis progression especially in high-risk patients affected by T2D. These findings further reinforce the indication of use RAS inhibitors to treat arterial hypertension or initial kidney disease in T2D patients in the presence of NAFLD.

Supporting Information

S1 File. Supplementary methods.

(DOCX)

S2 File. Supplementary references.

(DOCX)

S1 Table. Impact of baseline histological features of FPR in 118 Italian patients with NAFLD.

(DOCX)

S2 Table. Association of the changes of clinical variables during follow-up with fibrosis progression.

(DOCX)

S3 Table. Independent predictors of fibrosis progression at multivariate logistic regression analysis in 108 Italian patients with NAFLD without F4 fibrosis at baseline.

(DOCX)

Author Contributions

Conceptualization: LV S. Pelusi.

Data curation: S. Pelusi CR VB PD ALF.

Formal analysis: EB S. Petta S. Pelusi.

Funding acquisition: LV SF EB AC.

Methodology: LV S. Pelusi.

Visualization: LV S. Pelusi.

Writing – original draft: EB S. Petta S. Pelusi.

Writing – review & editing: EB S. Petta S. Pelusi.

References

1. Petta S, Valenti L, Bugianesi E, Targher G, Bellentani S, Bonino F. A "systems medicine" approach to the study of non-alcoholic fatty liver disease. *Dig Liver Dis* 2016; 48(3):333–42. doi: [10.1016/j.dld.2015.10.027](https://doi.org/10.1016/j.dld.2015.10.027) PMID: [26698409](https://pubmed.ncbi.nlm.nih.gov/26698409/)
2. Dongiovanni P, Petta S, Mannisto V, Mancina RM, Pipitone R, Karja V et al. Statin use and nonalcoholic steatohepatitis in at risk individuals. *J Hepatol* 2015; 63:705–12. doi: [10.1016/j.jhep.2015.05.006](https://doi.org/10.1016/j.jhep.2015.05.006) PMID: [25980762](https://pubmed.ncbi.nlm.nih.gov/25980762/)
3. Dongiovanni P, Rametta R, Meroni M, Valenti L. The role of insulin resistance in nonalcoholic steatohepatitis and liver disease development—a potential therapeutic target? *Expert Rev Gastroenterol Hepatol* 2015:1–14.

4. Dongiovanni P, Valenti L. Genetics of nonalcoholic fatty liver disease. *Metabolism* 2015.
5. Marchesini G, Brizi M, Bianchi G, Tomassetti S, Bugianesi E, Lenzi M et al. Nonalcoholic fatty liver disease: a feature of the metabolic syndrome. *Diabetes* 2001; 50(8):1844–50. PMID: [11473047](#)
6. Angulo P, Kleiner DE, Dam-Larsen S, Adams LA, Bjornsson ES, Charatcharoenwitthaya P et al. Liver Fibrosis, but No Other Histologic Features, Is Associated With Long-term Outcomes of Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2015; 149(2):389–97 e10. doi: [10.1053/j.gastro.2015.04.043](#) PMID: [25935633](#)
7. Ekstedt M, Hagstrom H, Nasr P, Fredrikson M, Stål P, Kechagias S et al. Fibrosis stage is the strongest predictor for disease-specific mortality in NAFLD after up to 33 years of follow-up. *Hepatology* 2015; 61(5):1547–54. doi: [10.1002/hep.27368](#) PMID: [25125077](#)
8. McPherson S, Hardy T, Henderson E, Burt AD, Day CP, Anstee QM. Evidence of NAFLD progression from steatosis to fibrosing-steatohepatitis using paired biopsies: implications for prognosis and clinical management. *J Hepatol* 2015; 62(5):1148–55. doi: [10.1016/j.jhep.2014.11.034](#) PMID: [25477264](#)
9. Singh S, Allen AM, Wang Z, Prokop LJ, Murad MH, Loomba R. Fibrosis progression in nonalcoholic fatty liver vs nonalcoholic steatohepatitis: a systematic review and meta-analysis of paired-biopsy studies. *Clin Gastroenterol Hepatol* 2015; 13(4):643–54.e1-9; quiz e39-40. doi: [10.1016/j.cgh.2014.04.014](#) PMID: [24768810](#)
10. Ekstedt M, Franzen LE, Mathiesen UL, Thorelius L, Holmqvist M, Bodemar G et al. Long-term follow-up of patients with NAFLD and elevated liver enzymes. *Hepatology* 2006; 44(4):865–73. PMID: [17006923](#)
11. Pais R, Charlotte F, Fedchuk L, Bedossa P, Lebray P, Poynard T et al. A systematic review of follow-up biopsies reveals disease progression in patients with non-alcoholic fatty liver. *J Hepatol* 2013; 59(3):550–6. doi: [10.1016/j.jhep.2013.04.027](#) PMID: [23665288](#)
12. Singh S, Khera R, Allen AM, Murad MH, Loomba R. Comparative effectiveness of pharmacological interventions for non-alcoholic steatohepatitis: A systematic review and network meta-analysis. *Hepatology* 2015.
13. Matthew Morris E, Fletcher JA, Thyfault JP, Rector RS. The role of angiotensin II in nonalcoholic steatohepatitis. *Mol Cell Endocrinol* 2013; 378(1–2):29–40. doi: [10.1016/j.mce.2012.04.013](#) PMID: [22579612](#)
14. Jonsson JR, Clouston AD, Ando Y, Kelemen LI, Horn MJ, Adamson MD et al. Angiotensin-converting enzyme inhibition attenuates the progression of rat hepatic fibrosis. *Gastroenterology* 2001; 121(1):148–55. PMID: [11438504](#)
15. Bataller R, Gabele E, Parsons CJ, Morris T, Yang L, Schoonhoven R et al. Systemic infusion of angiotensin II exacerbates liver fibrosis in bile duct-ligated rats. *Hepatology* 2005; 41(5):1046–55. PMID: [15841463](#)
16. Hirose A, Ono M, Saibara T, Nozaki Y, Masuda K, Yoshioka A et al. Angiotensin II type 1 receptor blocker inhibits fibrosis in rat nonalcoholic steatohepatitis. *Hepatology* 2007; 45(6):1375–81. PMID: [17518368](#)
17. Moreno M, Gonzalo T, Kok RJ, Sancho-Bru P, van Beuge M, Swart J et al. Reduction of advanced liver fibrosis by short-term targeted delivery of an angiotensin receptor blocker to hepatic stellate cells in rats. *Hepatology* 2010; 51(3):942–52. doi: [10.1002/hep.23419](#) PMID: [20044807](#)
18. Yang L, Bataller R, Dulyx J, Coffman TM, Ginès P, Rippe RA et al. Attenuated hepatic inflammation and fibrosis in angiotensin type 1a receptor deficient mice. *J Hepatol* 2005; 43(2):317–23. PMID: [15964094](#)
19. Sturzeneker MC, Ioshii SO, Villela Baroncini LA, Precoma DB. Olmesartan severely weakened the development of NASH in an animal model of hypercholesterolemia. *Atherosclerosis* 2011; 216(1):97–102. doi: [10.1016/j.atherosclerosis.2011.01.047](#) PMID: [21338989](#)
20. Wu Y, Ma KL, Zhang Y, Wen Y, Wang GH, Hu ZB et al. Lipid disorder and intrahepatic renin-angiotensin system activation synergistically contributes to non-alcoholic fatty liver disease. *Liver Int* 2016.
21. Yokohama S, Yoneda M, Haneda M, Okamoto S, Okada M, Aso K et al. Therapeutic efficacy of an angiotensin II receptor antagonist in patients with nonalcoholic steatohepatitis. *Hepatology* 2004; 40(5):1222–5. PMID: [15382153](#)
22. Torres DM, Jones FJ, Shaw JC, Williams CD, Ward JA, Harrison SA. Rosiglitazone versus rosiglitazone and metformin versus rosiglitazone and losartan in the treatment of nonalcoholic steatohepatitis in humans: a 12-month randomized, prospective, open-label trial. *Hepatology* 2011; 54(5):1631–9. doi: [10.1002/hep.24558](#) PMID: [21748770](#)
23. Goh GB, Pagadala MR, Dasarathy J, Unalp-Arida A, Sargent R, Hawkins C et al. Renin-angiotensin system and fibrosis in non-alcoholic fatty liver disease. *Liver Int* 2015; 35(3):979–85. doi: [10.1111/liv.12611](#) PMID: [24905085](#)

24. Orlic L, Mikolasevic I, Lukenda V, Anic K, Jelic I, Racki S. Nonalcoholic fatty liver disease and the renin-angiotensin system blockers in the patients with chronic kidney disease. *Wien Klin Wochenschr* 2015; 127(9–10):355–62. doi: [10.1007/s00508-014-0661-y](https://doi.org/10.1007/s00508-014-0661-y) PMID: [25412597](https://pubmed.ncbi.nlm.nih.gov/25412597/)
25. Dongiovanni P, Petta S, Maglio C, Fracanzani AL, Pipitone R, Mozzi E et al. Transmembrane 6 superfamily member 2 gene variant disentangles nonalcoholic steatohepatitis from cardiovascular disease. *Hepatology* 2015; 61(2):506–14. doi: [10.1002/hep.27490](https://doi.org/10.1002/hep.27490) PMID: [25251399](https://pubmed.ncbi.nlm.nih.gov/25251399/)
26. Valenti L, Fracanzani AL, Dongiovanni P, Rovida S, Rametta R, Fatta E et al. A randomized trial of iron depletion in patients with nonalcoholic fatty liver disease and hyperferritinemia. *World J Gastroenterol* 2014; 20(11):3002–10. doi: [10.3748/wjg.v20.i11.3002](https://doi.org/10.3748/wjg.v20.i11.3002) PMID: [24659891](https://pubmed.ncbi.nlm.nih.gov/24659891/)
27. Colloredo G, Guido M, Sonzogni A, Leandro G. Impact of liver biopsy size on histological evaluation of chronic viral hepatitis: the smaller the sample, the milder the disease. *J Hepatol* 2003; 39(2):239–44. PMID: [12873821](https://pubmed.ncbi.nlm.nih.gov/12873821/)
28. Kleiner DE, Brunt EM, Van Natta M, Behling C, Contos MJ, Cummings OW et al. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology* 2005; 41(6):1313–21. PMID: [15915461](https://pubmed.ncbi.nlm.nih.gov/15915461/)
29. European Association for the Study of the Liver. Electronic address eee, European Association for the Study of D, European Association for the Study of O. EASL-EASD-EASO Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease. *J Hepatol* 2016.
30. Petta S, Valenti L, Marra F, Grimaudo S, Tripodo C, Bugianesi E et al. MERTK rs4374383 polymorphism affects the severity of fibrosis in non-alcoholic fatty liver disease. *J Hepatol* 2016; 64(3):682–90. doi: [10.1016/j.jhep.2015.10.016](https://doi.org/10.1016/j.jhep.2015.10.016) PMID: [26596542](https://pubmed.ncbi.nlm.nih.gov/26596542/)
31. Marchesini G, Bugianesi E, Forlani G, Cerrelli F, Lenzi M, Manini R et al. Nonalcoholic fatty liver, steatohepatitis, and the metabolic syndrome. *Hepatology* 2003; 37(4):917–23. PMID: [12668987](https://pubmed.ncbi.nlm.nih.gov/12668987/)
32. Rosso C, Mezzabotta L, Gaggini M, Salomone F, Gambino R, Marengo A et al. Peripheral insulin resistance predicts liver damage in nondiabetic subjects with nonalcoholic fatty liver disease. *Hepatology* 2016; 63(1):107–16. doi: [10.1002/hep.28287](https://doi.org/10.1002/hep.28287) PMID: [26473614](https://pubmed.ncbi.nlm.nih.gov/26473614/)
33. Fracanzani AL, Valenti L, Bugianesi E, Andreoletti M, Colli A, Vanni E et al. Risk of severe liver disease in nonalcoholic fatty liver disease with normal aminotransferase levels: a role for insulin resistance and diabetes. *Hepatology* 2008; 48(3):792–8. doi: [10.1002/hep.22429](https://doi.org/10.1002/hep.22429) PMID: [18752331](https://pubmed.ncbi.nlm.nih.gov/18752331/)
34. Ziyadeh FN, Wolf G. Pathogenesis of the podocytopathy and proteinuria in diabetic glomerulopathy. *Curr Diabetes Rev* 2008; 4(1):39–45. PMID: [18220694](https://pubmed.ncbi.nlm.nih.gov/18220694/)
35. Hall JE, do Carmo JM, da Silva AA, Wang Z, Hall ME. Obesity-induced hypertension: interaction of neurohumoral and renal mechanisms. *Circ Res* 2015; 116(6):991–1006. doi: [10.1161/CIRCRESAHA.116.305697](https://doi.org/10.1161/CIRCRESAHA.116.305697) PMID: [25767285](https://pubmed.ncbi.nlm.nih.gov/25767285/)
36. Ekstedt M, Franzen LE, Mathiesen UL, Holmqvist M, Bodemar G, Kechagias S. Statins in non-alcoholic fatty liver disease and chronically elevated liver enzymes: a histopathological follow-up study. *J Hepatol* 2007; 47(1):135–41. PMID: [17400325](https://pubmed.ncbi.nlm.nih.gov/17400325/)
37. Dongiovanni P, Fracanzani AL, Fargion S, Valenti L. Iron in fatty liver and in the metabolic syndrome: a promising therapeutic target. *J Hepatol* 2011; 55(4):920–32. doi: [10.1016/j.jhep.2011.05.008](https://doi.org/10.1016/j.jhep.2011.05.008) PMID: [21718726](https://pubmed.ncbi.nlm.nih.gov/21718726/)