



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

WITHDRAWN: Soluble urokinase plasminogen activator receptor and its complicated role in hemodialysis (HD) patients with Covid-19 infection

Ioannis Griveas*, Evangelos Pratilas

Nephrology Department, Army Share Fund Hospital of Athens, 417 NIMTS, Greece

This article has been withdrawn at the request of the author(s) and/or editor. The Publisher apologizes for any inconvenience this may cause.

The full Elsevier Policy on Article Withdrawal can be found at <https://www.elsevier.com/about/our-business/policies/article-withdrawal>.

*Corresponding author. 362 Kifisias Ave, 15233, Chalandri, Athens, Greece.
E-mail address: giannisgriv@hotmail.com (I. Griveas).