

Research article

Open Access

## Spinocerebellar ataxia type 17: Report of a family with reduced penetrance of an unstable Gln<sub>49</sub> TBP allele, haplotype analysis supporting a founder effect for unstable alleles and comparative analysis of SCA17 genotypes

Christine Zühlke\*<sup>1</sup>, Andreas Dalski<sup>1</sup>, Eberhard Schwinger<sup>1</sup> and Ulrich Finckh<sup>2</sup>

Address: <sup>1</sup>Institute of Human Genetics, University of Lübeck, Ratzeburger Allee 160, 23538 Lübeck, Germany and <sup>2</sup>Institute of Human Genetics, Universitätsklinikum Eppendorf, Butenfeld 42, 22529 Hamburg, Germany

Email: Christine Zühlke\* - [zuehlke@uni-luebeck.de](mailto:zuehlke@uni-luebeck.de); Andreas Dalski - [dalski@medinf.mu-luebeck.de](mailto:dalski@medinf.mu-luebeck.de); Eberhard Schwinger - [schwinger@uni-luebeck.de](mailto:schwinger@uni-luebeck.de); Ulrich Finckh - [finckh@labmed.de](mailto:finckh@labmed.de)

\* Corresponding author

Published: 01 July 2005

Received: 10 February 2005

*BMC Medical Genetics* 2005, **6**:27 doi:10.1186/1471-2350-6-27

Accepted: 01 July 2005

This article is available from: <http://www.biomedcentral.com/1471-2350/6/27>

© 2005 Zühlke et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Spinocerebellar ataxia type 17 (SCA17), a neurodegenerative disorder in man, is caused by an expanded polymorphic polyglutamine-encoding trinucleotide repeat in the gene for TATA-box binding protein (TBP), a main transcription factor. Observed pathogenic expansions ranged from 43 – 63 glutamine (Gln) codons (Gln<sub>43–63</sub>). Reduced penetrance is known for Gln<sub>43–48</sub> alleles. In the vast majority of families with SCA17 an expanded CAG repeat interrupted by a CAA CAG CAA element is inherited stably.

**Results:** Here, we report the first pedigree with a Gln<sub>49</sub> allele that is a) not interrupted, b) unstable upon transmission, and c) associated with reduced penetrance or very late age of onset. The 76-year-old father of two SCA17 patients carries the Gln<sub>49</sub> TBP allele but presents without obvious neurological symptoms. His children with Gln<sub>53</sub> and Gln<sub>52</sub> developed ataxia at the age of 41 and 50. Haplotype analysis of this and a second family both with uninterrupted expanded and unstable pathological SCA17 alleles revealed a common core genotype not present in the interrupted expansion of an unrelated SCA17 patient. Review of the literature did not present instability in SCA17 families with expanded alleles interrupted by the CAA CAG CAA element.

**Conclusion:** The presence of a Gln<sub>49</sub> SCA17 allele in an asymptomatic 76-year-old male renews the discussion of reduced penetrance and genotypes producing very late disease onset. In SCA17, uninterrupted expanded alleles of TBP are associated with repeat instability and a common founder haplotype. This suggests for uninterrupted expanded alleles a mutation mechanism and some clinical genetic features distinct from those alleles interrupted by a CAA CAG CAA element.

### Background

The spinocerebellar ataxias (SCAs), a group of autosomal

dominantly inherited human disorders with mainly adult age of onset, are caused by progressive neurodegeneration

and significant cerebellar dysfunction, but also involve other regions of the central or peripheral nervous system. Clinically and even histopathologically the differentiation between SCA subtypes may be rather difficult. Genetically, for 25 types causative mutations in various genes are known or defined by linkage to distinct chromosomal regions. For seven SCAs expanded CAG trinucleotide repeats have been discovered [1]. This kind of mutation seems to be specific for the human species.

SCA17 [OMIM: 607136], a rare type of SCA with a variety of clinical features, is caused by an expanded CAG repeat in *TBP*, the gene of the TATA box-binding protein [TBP; OMIM: 600075]. TBP forms the DNA-binding subunit of the universally essential RNA polymerase II transcription factor D. TBP has been mapped close to the telomeric region at chromosome 6q27 [2]. An imperfect repetitive and polymorphic polyglutamine encoding CAG triplet sequence is part of the coding region of TBP [3]. Initially, 20 different alleles coding for 29 to 42 glutamine residues (Gln) have been identified [4] with the most common alleles containing 32 to 39 Gln codons. The polyglutamine-encoding DNA sequence of *TBP* wildtype alleles can be subdivided into several regions including two polymorphic (CAG)<sub>n</sub> stretches: (CAG)<sub>3</sub> (CAA)<sub>3</sub> (CAG)<sub>n</sub> CAA CAG CAA (CAG)<sub>n</sub> CAA CAG. In patients with SCA17, the polymorphic CAG sequence encoding the polyglutamine stretch of TBP is expanded heterozygous. Alleles with 43 - 48 Gln codons represent a zone of incomplete penetrance or very late age of onset [5,6].

Disease causing alleles with up to 63 [7] Gln codons were identified. Clinically, SCA17 may mimic a broad spectrum of neuropsychiatric diseases, including Parkinson and Huntington disease, major psychosis, and multitudinous cerebellar syndromes [8-10]. Neuropathological findings include cerebellar, cortical, and subcortical atrophy, Purkinje cell loss, gliosis [11], and neuronal intranuclear inclusions immunopositive for TBP and polyglutamine.

The clinical relevance of Gln<sub>43-48</sub> encoding *TBP* alleles is not obvious. A Gln<sub>43</sub> allele was considered responsible for clinical symptoms in a 64-year-old patient with ataxia and progressive mental deterioration [6]. Gln<sub>44</sub> repeats were found in a patient with gait ataxia and behavioral changes and a disease onset at 29 [12] and another patient with early onset cerebellar ataxia (EOCA) and suspected Friedreich ataxia [5]. Regarding the EOCA phenotype in the latter patient, Gln<sub>44</sub> was considered a large normal allele. On the other hand, alleles for Gln<sub>46</sub> [12] and Gln<sub>48</sub> [13] showed reduced penetrance in two families. Therefore, Gln<sub>43-48</sub> encoding alleles could present intermediate alleles with incomplete penetrance.

In contrast to the majority of other polyglutamine diseases, the SCA17 repeat expansion shows meiotic stability upon transmission. To date, only two pedigrees are known in which instability of the repetitive sequence has been observed, irrespectively of the gender of the transmitting parent. In one case, the repeat increased by one triplet after maternal transmission [5], while in the other case an increase of 13 triplets and marked anticipation was associated with paternal inheritance [14].

Here, we describe a SCA17 family from northern Germany with a) repeat instability upon paternal transmission and b) reduced penetrance or very late onset in association with an uninterrupted CAG<sub>41</sub> sequence (Gln<sub>49</sub> allele). Meiotic instability is not common in SCA17 pedigrees. Therefore, we performed haplotype analysis in two unrelated families looking for a founder allele associated with repeat instability at the SCA17 locus. In addition, we included a patient homozygous for a SCA17 allele with a CAG CAA CAG interrupting element to reveal independent mutation events for the different repeat compositions.

## Results


### Phenotype and genotype

In family A, two of four offspring of unaffected parents (74- and 76-year-old) developed an adult onset ataxia. The 43-year-old index patient reported onset of gait disturbances accompanied by a slowly progressive dysarthria at the age of 41. Brain magnetic resonance image analysis performed approximately half a year after clinical onset revealed cortical cerebellar atrophy and a mildly reduced brain volume. His 56-year-old sister noticed symptoms of gait ataxia, dysarthria, and disturbed handwriting at the age of 50. The 76-year-old father of the siblings reported that a neurological examination had revealed no abnormality. Both parents presented without obvious neurological signs. In order to clarify the repeat status of *TBP* and the inheritance of the disease in the family, both agreed in molecular genetic analysis for SCA17 but rejected the offer of a detailed clinical neurological examination.

Molecular genetic analyses revealed pathogenic alleles for the *TBP* gene: 53 repeats for the male patient with age of onset at 41 and 52 repeats for his sister with symptoms starting at 50 years of age. Their clinically non-affected 76-year-old father carries an elongated allele of 49 repeats. Therefore, this allele displays reduced penetrance or very late onset, as well as instability upon transmission. Sequence analysis revealed loss of the CAA CAG CAA interruption in the expanded alleles.

### Haplotype analysis

Recently, we described a family (B) with repeat instability and missing CAA triplets at the expanded alleles [5,10]. Genealogic data for both families (A,B), as far as available,



**Figure 1**  
**Pedigrees and haplotype analysis.** Pedigrees of families A and B, including six-locus haplotypes based on five 6q-telomeric microsatellite markers (D6S1719, D6S264, D6S281, D6S446, D6S1590) and closely downstream located polymorphic poly-Gln encoding trinucleotide stretch within *TBP*. Alleles of the microsatellites are denoted by numbers 1 – 8, *TBP* alleles are denoted by the number of Gln-encoding trinucleotides. SCA17-linked alleles are boxed as well as the disease haplotype D6S446 and D6S1590.

revealed no relationship between the families. But Northern German ancestry of both families suggested the possibility of a common founder. This was assessed by haplotype analysis (figure 1) including the unrelated patient H carrying homozygous an expansion in the *TBP* gene [15]. In both families A and B, the same haplotype consisting of allele 5 of D6S446 and allele 2 of D6S1590 (5\_2) was located on the chromosome bearing the expanded *TBP* allele co-segregating with SCA17, respectively. Patient H did not share this haplotype, she is homozygous for haplotype 2\_4. Among the markers analyzed, D6S446 and D6S1590 were the ones most closely located to *TBP*. Therefore, the data suggest the possibility of a common SCA17 founder allele of the variably

expanded *TBP* allele in families A and B with further instability upon transmission.

**Allele frequency**

To estimate the probability of a founder effect, the allele frequencies of D6S446 and D6S1590 were determined in 96 anonymous control subjects of the geographic region of families A and B. For D6S446 six alleles were found, for D6S1590 seven. Allele 5 of D6S446 is present in 43 out of 184 chromosomes (23%), allele 2 of D6S1590 in 92 (50%). The computed maximum likelihood frequencies (using Arlequin software version 2.001) of the three most common two-locus haplotypes of D6S446 and D6S1590 with frequencies > 0.1 are 0.219 (6\_2), 0.163 (6\_1), and

**Table 1: Review of Gln Repeats and SCA17 Alleles of 16 Unrelated Cases. Duplicated elements are boxed, instability by maternal (m) or paternal (p) inheritance.**

Gln	Case	Repeat Composition	Comments	Ref.
49-53	familial	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>41-45</sub> CAA CAG	instability (p)	here
63	sporadic	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>9</sub> CAA CAG CAA (CAG) <sub>9</sub> (CAA) <sub>3</sub> (CAG) <sub>9</sub> CAA CAG CAA (CAG) <sub>19</sub> CAA CAG	duplication	[7]
47	familial	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>8</sub> CAA CAG CAA (CAG) <sub>28</sub> CAA CAG		[16]
47	sporadic	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>6</sub> CAA CAG CAA (CAG) <sub>30</sub> CAA CAG		
48	familial	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>6</sub> CAA CAG CAA (CAG) <sub>31</sub> CAA CAG		
55	familial	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>9</sub> CAA CAG CAA (CAG) <sub>16</sub> CAA CAG CAA (CAG) <sub>16</sub> CAA CAG	duplication	
51	familial	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>9</sub> CAA CAG CAA (CAG) <sub>31</sub> CAA CAG		[5]
53 - 55	familial	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>45-47</sub> CAA CAG	instability (m)	
48	familial	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>9</sub> CAA CAG CAA (CAG) <sub>28</sub> CAA CAG	reduced penetrance	[13]
47	sporadic	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>9</sub> CAA CAG CAA (CAG) <sub>27</sub> CAA CAG	homozygous mutation	[15]
46	familial	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>9</sub> CAA CAG CAA (CAG) <sub>26</sub> CAA CAG		[21]
43		(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>9</sub> CAA CAG CAA (CAG) <sub>26</sub> CAA CAG		[6]
53 - 66	familial	(CAG) <sub>3</sub> (CAA) <sub>4</sub> (CAG) <sub>44/57</sub> CAA CAG	instability (p)	[14]
44	sporadic	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>9</sub> CAA CAG CAA (CAG) <sub>24</sub> CAA CAG		[12]
46	sporadic	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>11</sub> CAA CAG CAA (CAG) <sub>24</sub> CAA CAG	reduced penetrance	
48	familial	(CAG) <sub>3</sub> (CAA) <sub>3</sub> (CAG) <sub>6</sub> CAA CAG CAA (CAG) <sub>31</sub> CAA CAG	homozygous mutation	[23]

0.153 (5<sub>2</sub>). Thus, the three most common haplotypes, including the haplotype 5<sub>2</sub> co-segregating with SCA17 in families A and B account for 53.5% of all haplotypes. The 17 remaining haplotypes predicted had computed frequencies from 0.3-6.8%. Maximum likelihood frequency of haplotype 4<sub>2</sub> (patient H) is 0.042. The likelihood ratio test suggested a highly significant linkage disequilibrium between D6S446 and D6S1590 ( $p < 5 \times 10^{-6}$ ). Observed and maximum likelihood counts of genotypes and haplotypes did not deviate significantly from Hardy Weinberg equilibrium. Therefore, the likelihood of two unrelated subjects sharing haplotype 5<sub>2</sub> is estimated to be ~8%. The combined likelihood that two unrelated subjects affected by SCA17 share haplotype 5<sub>2</sub> and that this haplotype co-segregates with SCA17 in both families is ~2%. This strongly supports the view of a possible founder haplotype of the SCA17 causing alleles in families A and B.

## Discussion

In comparison to other spinocerebellar ataxias, some genetic features in SCA17 are remarkable: The majority of pathological alleles contains an interspersed CAA CAG CAA element separating the (CAG)<sub>n</sub> sequence into two parts (table 1). In three SCA17 pedigrees known to date the pathogenic allele lacks the CAA CAG CAA interruption and is associated with instability and variable repeat expansion upon transmission. Two of the three families are of northern German origin and were investigated in the study presented. In both families, SCA17 co-segregates with an expanded, uninterrupted, and instable (CAG)<sub>n</sub> in *TBP* located on chromosomes sharing a haplotype in the close centromeric neighborhood of *TBP*, respectively (fig-

ure 1). Although the unstable repetitive sequence is linked with one of the three more commonly prevalent haplotypes, statistics point to a nominally significant likelihood that our observation reflects a founder effect rather than coincidence by chance.

In some cases, the expanded glutamine stretch may be the consequence of an intragenic duplication. So, the first published case with SCA17 is a Japanese girl with a de novo duplication event in her paternal allele [7]. In addition, doubling of a stretch of 19 Gln codons was found in a three-generation SCA17 pedigree [16]. Expanded polyglutamine stretches arising from duplications represent rare events in SCAs. Similarly, a pathogenic elongation of the polyalanine part within the polyadenylate binding protein nuclear 1 resulting from duplication has been described for oculopharyngeal muscular dystrophy [17]. The identification of these mutations gives support to the hypothesis of unequal crossing-over as one possible molecular mechanism for repeat expansions. In SCA17, such rare unequal crossing-over events may have been the causative mechanism underlying both the loss of the interspersed CAA CAG CAA element and expansion of (CAG)<sub>n</sub>. As discussed for SCA2 [18] the presence of CAA interruptions in SCA17 alleles breaks the repetitive sequence into shorter homogenous triplet tracts and may thus protect it from instability by reducing the slippage between the complementary strands.

In SCA17, there is a broad range of intermediate *TBP* alleles Gln<sub>43-48</sub> associated with reduced penetrance. In other SCA types, such intermediate alleles are rarely found [19,20] or even unknown. However, intermediate alleles

represent a well-known finding in patients with Huntington disease [21]. We found (CAG/CAA)<sub>49</sub> but no visible symptoms at the age of 76 in the father of two SCA17 patients with expanded *TBP* alleles (CAG/CAA)<sub>52-53</sub>. This large allele, (CAG/CAA)<sub>49</sub>, has not been found in control samples. It may be of reduced penetrance, associated with very late age of onset and/or low expression of clinical signs. In addition, we performed haplotype analysis for a sporadic patient with homozygosity of the intermediate allele (CAG/CAA)<sub>47</sub> and a rather progressive course [15]. Here, we cannot exclude the possibility of an additive effect of two intermediate alleles with respect to pathogenicity. In this patient, (CAG/CAA)<sub>47</sub> is interrupted by CAA CAG CAA and linked with D6S446/D6S1590 haplotype 2\_4. Thus, both the type of *TBP* expansion and the SCA17 linked haplotypes differ between the sporadic homozygous and the familial cases presented and do not point to a single founder. An extended analysis of intra-genetic markers in larger numbers of SCA17 families with different types of expansions could reveal "mutation prone" founder alleles differing with respect to the type of mutation.

## Conclusion

The extraordinary high variability of the clinical expression of pathological *TBP* repeat expansions in SCA17 is further complicated by the occurrence of genetically unstable repeats in association with the lack of an interspersing CAA CAG CAA element. The degree of reduced penetrance and very low expression of symptoms refer to strong modifying factors including potent influence of the genetic background. Unless SCA17 is a rare type of dominantly inherited ataxia, the repeat expansion within the *TBP* gene arose at unrelated genotypes. This has to be taken into account both in molecular genetic diagnostics and in genetic counseling.

## Methods

### Molecular analysis and polymorphic markers

Genomic DNA was prepared from peripheral blood leukocytes using standard protocols. SCA17 alleles were amplified by PCR as described [7], separated on 6% denaturing polyacrylamide gels, and visualized by silver staining. For sequencing, PCR products were separated under non-denaturing conditions using the dHPLC-HT-system (WAVE Transgenomic). The eluted fragments were sequenced using dye terminators and the automated capillary sequencer Avant 3100 (Applied Biosystems). For haplotype analysis, the highly polymorphic microsatellite markers D6S1719, D6S264, D6S281, D6S446, and D6S1590 were used as described [15]. Maximum likelihood haplotype frequencies of two-locus haplotypes based on markers D6S446 and D6S1590 genotyped in 96 unrelated anonymous control subjects and linkage dise-

quilibrium were analyzed using Arlequin software version 2.001 [22].

## Competing interests

The author(s) declare that they have no competing interests.

## Authors' contributions

CZ conceived of the study and developed the concept, interpreted the data and contributed substantially to the manuscript (corresponding author). AD carried out the molecular genetic studies. ES participated in the design of the study and helped to draft the manuscript. UF participated in the design and performed the statistical analysis.

All authors read and approved the final manuscript.

## Acknowledgements

We would like to thank Ulrike Gehlken for excellent technical assistance and the German Heredo-Ataxia Society (DHAG), whose cooperation is essential in our work.

## References

- Margolis RL: **The spinocerebellar ataxias: order emerges from chaos.** *Curr Neurol Neurosci Rep* 2002, **2**:447-456.
- Imbert G, Trottier Y, Beckmann J, Mandel JL: **The gene for the TATA binding protein (TBP) that contains a highly polymorphic protein coding CAG repeat maps to 6q27.** *Genomics* 1994, **21**:667-668.
- Polymeropoulos MH, Rath DS, Xiao H, Merrill CR: **Trinucleotide repeat polymorphism at the human transcription factor IID gene.** *Nucleic Acids Res* 1991, **19**:4307.
- Gostout B, Liu Q, Sommer SS: **'Cryptic' repeating triplets of purines and pyrimidines (cRRY(i)) are frequent and polymorphic: analysis of coding cRRY(i) in the proopiomelanocortin (POMC) and TATA-binding protein (TBP) genes.** *Am J Hum Genet* 1993, **52**:1182-1190.
- Zühlke C, Hellenbroich Y, Dalski A, Kononowa N, Hagenah J, Vierge P, Riess O, Klein C, Schwinger E: **Different types of repeat expansion in the TATA-binding protein gene are associated with a new form of inherited ataxia.** *Eur J Hum Genet* 2001, **9**:160-164.
- Silveira I, Miranda C, Guimaraes L, Moreira MC, Alonso I, Mendonca P, Ferro A, Pinto-Basto J, Coelho J, Ferreirinha F, Poirier J, Parreira E, Vale J, Januario C, Barbot C, Tuna A, Barros J, Koide R, Tsuji S, Holmes SE, Margolis RL, Jardim L, Pandolfo M, Coutinho P, Sequeiros J: **Trinucleotide repeats in 202 families with ataxia: a small expanded (CAG)<sub>n</sub> allele at the SCA17 locus.** *Arch Neurol* 2002, **59**:623-629.
- Koide R, Kobayashi S, Shimohata T, Ikeuchi T, Maruyama M, Saito M, Yamada M, Takahashi H, Tsuji S: **A neurological disease caused by an expanded CAG trinucleotide repeat in the TATA-binding protein gene: a new polyglutamine disease?** *Hum Molec Genet* 1999, **8**:2047-2053.
- De Michele G, Maltecca F, Carella M, Volpe G, Orio M, De Falco A, Gombia S, Servadio A, Casari G, Filla A, Bruni A: **Dementia, ataxia, extrapyramidal features, and epilepsy spectrum in two Italian families with spinocerebellar ataxia type 17.** *Neurol Sci* 2003, **24**:166-167.
- Bauer P, Laccone F, Rolfs A, Wullner U, Bosch S, Peters H, Liebscher S, Scheible M, Epplen JT, Weber BH, Holinski-Feder E, Weirich-Schwaiger H, Morris-Rosendahl DJ, Andrich J, Riess O: **Trinucleotide repeat expansion in SCA17/TBP in white patients with Huntington's disease-like phenotype.** *J Med Genet* 2004, **41**:230-232.
- Hagenah JM, Zühlke Ch, Hellenbroich Y, Heide W, Klein Ch: **Focal dystonia as presenting sign of SCA17.** *Mov Dis* 2004, **19**:217-220.

11. Rolfs A, Koeppen AH, Bauer I, Bauer P, Buhlmann S, Topka H, Schols L, Riess O: **Clinical features and neuropathology of autosomal dominant spinocerebellar ataxia (SCA17).** *Ann Neurol* 2003, **54**:367-375.
12. Stevanin G, Fujigasaki H, Lebre AS, Camuzat A, Jeannequin C, Dode C, Takahashi J, San C, Bellance R, Brice A, Durr A: **Huntington's disease-like phenotype due to trinucleotide repeat expansions in the TBP and JPH3 genes.** *Brain* 2003, **126**:1599-1603.
13. Zühlke C, Gehlken U, Hellenbroich Y, Schwinger E, Bürk K: **Phenotypic variability of expanded alleles in the TATA-binding protein gene. Reduced penetrance in SCA17?** *J Neurol* 2003, **250**:161-163.
14. Maltecca F, Filla A, Castaldo I, Coppola G, Fragassi NA, Carella M, Bruni A, Cocozza S, Casari G, Servadio A, De Michele G: **Intergenerational instability and marked anticipation in SCA-17.** *Neurology* 2003, **61**:1441-1443.
15. Zühlke C, Spranger M, Spranger S, Voigt R, Lanz M, Gehlken U, Hinrichs F, Schwinger E: **SCA17 caused by homozygous repeat expansion in TBP due to partial isodisomy 6.** *Eur J Hum Genet* 2003, **11**:629-632.
16. Nakamura K, Jeong SY, Uchihara T, Anno M, Nagashima K, Nagashima T, Ikeda S, Tsuji S, Kanazawa I: **SCA17, a novel autosomal dominant cerebellar ataxia caused by an expanded polyglutamine in TATA-binding protein.** *Hum Molec Genet* 2001, **10**:1441-1448.
17. van der Sluijs BM, van Engelen BG, Hoefsloot LH: **Oculopharyngeal muscular dystrophy (OPMD) due to a small duplication in the PABPN1 gene.** *Hum Mutat* 2003, **21**:553.
18. Choudhry S, Mukerji M, Srivastava AK, Jain S, Brahmachari SK: **CAG repeat instability at SCA2 locus : anchoring CAA interruptions and linked single nucleotide polymorphisms.** *Hum Mol Genet* 2001, **10**:2437-2446.
19. Zühlke C, Dalski A, Hellenbroich Y, Bubel S, Schwinger E, Bürk K: **Spinocerebellar ataxia type I (SCA1): phenotype-genotype correlation studies in intermediate alleles.** *Eur J Hum Genet* 2002, **10**:204-209.
20. Andrew SE, Goldberg YP, Hayden MR: **Rethinking genotype and phenotype correlations in polyglutamine expansion disorders.** *Hum Mol Genet* 1997, **6**:2005-2010.
21. McNeil SM, Novelletto A, Srinidhi J, Barnes G, Kornbluth I, Altherr MR, Wasmuth JJ, Gusella JF, MacDonald ME, Myers RH: **Reduced penetrance of the Huntington's disease mutation.** *Hum Mol Genet* 1997, **6**:775-779.
22. Schneider S, Roessler D, Excofier L: **Arlequin: A software for population genetics data analysis. Ver 2.000.** Genetics and Biometry Lab, Dept. of Anthropology, University of Geneva; 2000.
23. Toyoshima Y, Yamada M, Onodera O, Shimohata M, Inenaga C, Fujita N, Morita M, Tsuji S, Takahashi H: **SCA17 homozygote showing Huntington's disease-like phenotype.** *Ann Neurol* 2004, **55**:281-286.

### Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2350/6/27/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

