

# A multicenter, open-label, phase III study of Abcetin in Gaucher disease: Erratum

In the article, “A multicenter, open-label, phase III study of Abcetin in Gaucher disease”,<sup>[1]</sup> which appears in Volume 96, Issue 45 of *Medicine*, Table 1 contained incorrect information and the correct table appears below.

**Table 1**

**Demographics and Baseline Clinical Characteristics of the Per Protocol Population.**

No.	Gender	Ethnicity	Age at enrollment (years)	Height (cm)	Weight (kg)	Hemoglobin (g/dL)	Platelet ( $\times 10^3/\mu\text{L}$ )	Spleen volume, MN	Liver volume, MN	Genotype	Enzyme activity (1–5 $\mu\text{mol/g/h}$ )
1	Male	Egyptian	15	137 (<3 <sup>rd</sup> P)	31 (<3 <sup>rd</sup> P)	8.9	129	56.32	1.23	NA	0.34*
2	Male	Egyptian	9	122 (3 <sup>rd</sup> P)	25 (<3 <sup>rd</sup> P)	10.1	120	12.24	0.69	NA	0.42*
3	Male	Egyptian	6	108 (10 <sup>th</sup> P)	17 (5 <sup>th</sup> P)	8.9	44	43.18	2.08	NA	0.2*
4	Male	Egyptian	8	114.2 (<3 <sup>rd</sup> P)	23 (25 <sup>th</sup> P)	10.3	193	17.26	1.03	NA	0.5*
5	Male	Egyptian	2	79 (<3 <sup>rd</sup> P)	10 (<3 <sup>rd</sup> P)	9.8	258	7.2	1.34	L444P/L444P	0.49*
6	Male	Egyptian	2	82 (5 <sup>th</sup> P)	11 (3 <sup>rd</sup> P)	10.3	110	22.09	2.25	L444P/N370S	0.49*
7	Male	Egyptian	2	83 (25 <sup>th</sup> P)	14 (50 <sup>th</sup> –75 <sup>th</sup> P)	8.1	74	45.04	2.09	L444P/N370S	1.6**
Mean $\pm$ SD (range)	Male	Egyptian	6.3 $\pm$ 4.86 (2–15)	103.6 $\pm$ 03.61 (79–137)	18.7 $\pm$ 8.7 (10–31)	9.5 $\pm$ 0.86 (8.1–10.3)	132.6 $\pm$ 72.27 (44–258)	29.0 $\pm$ 18.91 (7.2–56.32)	1.5 $\pm$ 0.61 (0.69–2.25)		

NA = not available, P = percentile, SD = standard deviation.

Normal range of GBA activity in peripheral leukocytes.

\* 1–5  $\mu\text{mol/g/h}$  and.

\*\* 9.3  $\mu\text{mol/g/h}$

## Reference

[1] Lee B-H, Abdalla A-F, Choi J-H. A multicenter, open-label, phase III study of Abcetin in Gaucher disease. *Medicine*. 96;45:e8492.