

POSTER PRESENTATIONS

Open Access

# Evaluation of *in vivo* Antiplasmodial Activities of extracts of *Morinda morindiodes* (Bak.) in the treatment of malaria in Ogun State

Soniran O Temidayo\*, Idowu Olufunmilayo, Idowu A Babatunde, Ajana Olusegun

From Parasite to Prevention: Advances in the understanding of malaria  
 Edinburgh, UK. 20-22 October 2010

*In vivo* study of various plant parts extracts of *Morinda morindiodes* (Bak.) was conducted to evaluate their antiplasmodial properties and effects on the liver using chloroquine sensitive *Plasmodium berghei* in mice. Water extract of the root was observed to significantly reduce parasitaemia (70%,  $P < 0.05$ ) compared to the activities of other plant parts and the untreated control. A mean survival time of 19 days observed in the root extract supported its antiplasmodial activities compared with other plant parts. The antiplasmodial activities of the plant extracts when administered twice daily were not significantly different

( $P > 0.05$ ) compared with those treated once daily. The chemosuppression produced by the extracts were significantly different compared to untreated control. Liver function tests (LFT) of uninfected mice administered with the plant extracts showed that extracts of the leaf and stem in 'fermented maize starch extract' altered the function of the liver significantly compared to normal mice. This study shows that *Morinda morindiodes* possess antimalarial properties and the root may be used as a prophylaxis where western medicine is not easily accessible and affordable. Tables 1, 2, 3, 4

**Table 1 Chemosuppression and survival time of *P. berghei* infected mice treated orally with *Morinda morindiodes* extracts at a dose of 100mg/kg body weight once a day for 5 days**

Plant part	Extract	%chemosuppression of parasitaemia at day 5 (C.P.±S.D)	%chemosuppression of parasitaemia at day 11 (C.P.±S.D)	Mean Survival time (days)
Leaf	MeOH	7.5 ± 2.8 <sup>b</sup>	82.4±5.6 <sup>e</sup>	19.5
	Water	10±2.8 <sup>bc</sup>	72.6±8.4 <sup>bc</sup>	16.5
	F.M. starch	7.5 ± 2.8 <sup>b</sup>	78.8 ± 9.9 <sup>d</sup>	16.5
Stem	MeOH	0 ± 2.8 <sup>a</sup>	60.8 ± 5.6 <sup>a</sup>	16.5
	Water	1.3 ± 1.41 <sup>ab</sup>	72.2 ± 14.8 <sup>bc</sup>	15.5
	F.M. starch	5.0 ± 2.8 <sup>b</sup>	71.6 ± 0.0 <sup>bc</sup>	16.5
Root	MeOH	20.5 ± 22.6 <sup>bc</sup>	67.3 ± 7.07 <sup>b</sup>	18.5
	Water	52.5±2.8 <sup>d</sup>	76.5 ± 0.0 <sup>cd</sup>	19.5
	F.M. starch	30±8.5 <sup>c</sup>	79.1 ± 0.0 <sup>d</sup>	19.5
Chloroquine			100 ± 0.0 <sup>e</sup>	28.5
Artesunate			100 ± 0.0 <sup>e</sup>	28.5
Control			0.0	14.5

MeOH, methanol extract; F.M. starch, "aqueous fermented maize starch ('omidun') extract".

**Table 2 Chemosuppression and survival time of *P. berghei* infected mice treated orally with *Morinda morindiodes* extracts at a dose of 100mg/kg body weight twice a day for 5 days**

Plant part	Extract	%chemosuppression of parasitaemia at day 5 (C.P.±S.D)	%chemosuppression of parasitaemia at day 11 (C.P.±S.D)	Mean Survival time (days)
Leaf	MeOH	20 ±16.9a	83.7±8.5 <sup>c</sup>	21.5
	Water	5.0 ± 5.6 <sup>ab</sup>	76.1 ± 1.4 <sup>bc</sup>	17.5
	F.M. starch	27.5 ± 2.8 <sup>abc</sup>	79.4 ± 24.1 <sup>bc</sup>	17.5
Stem	MeOH	1.3 ± 11.3 <sup>a</sup>	73.2 ± 25.4 <sup>b</sup>	16.5
	Water	2.5 ±28.2 <sup>a</sup>	71.2±2.8 <sup>b</sup>	16.5
	F.M. starch	10.0±5.6 <sup>ab</sup>	76.5 ± 9.9 <sup>bc</sup>	17.5
Root	MeOH	35 ± 5.6 <sup>bc</sup>	80.4±28.2 <sup>bc</sup>	19.5
	Water	70.0 ± 2.8 <sup>de</sup>	85.9±8.4 <sup>bc</sup>	21.5
	F.M. starch	56.2 ± 7.1 <sup>cd</sup>	85.6±8.4 <sup>c</sup>	17.5
Chloroquine			100 ± 0.0 <sup>e</sup>	28.5
Artesunate			100 ± 0.0 <sup>e</sup>	28.5
Control			0.0	14.5

MeOH, methanol extract; F.M. starch, "aqueous fermented maize starch ('omidun') extract".

**Table 3 Comparison between the liver function tests in mice treated with extracts and control group (untreated)**

Test	Leaf extracts		Stem extracts		Root extracts		Contrl	
	WL	F.M.L	WS	F.M.S	WR	F.M.R	INT.	
<b>Treated Once Daily</b>								
Total protein(g/l)	53 <sup>a</sup>	58.8 <sup>b</sup>	61.7 <sup>c</sup>	62.4 <sup>cd</sup>	63.5 <sup>cd</sup>	52.8 <sup>a</sup>	65.1d	
Cholesterol (mg/dl)	88.7 <sup>b</sup>	102.3 <sup>d</sup>	106.9 <sup>e</sup>	95.2 <sup>c</sup>	87.2 <sup>b</sup>	63.6 <sup>a</sup>	106.9e	
SGOT (iu/l)	66 <sup>c</sup>	67 <sup>c</sup>	85 <sup>d</sup>	91 <sup>e</sup>	67 <sup>c</sup>	30 <sup>a</sup>	44b	
SGPT (iu/l)	17 <sup>ab</sup>	20 <sup>b</sup>	18 <sup>b</sup>	13 <sup>a</sup>	28 <sup>c</sup>	25 <sup>c</sup>	27c	
Urea (mg/dl)	24.5 <sup>a</sup>	28.35 <sup>c</sup>	25.7 <sup>ab</sup>	26.5 <sup>bc</sup>	26.5 <sup>bc</sup>	28.35 <sup>c</sup>	28c	
Alkaline phosphatase (iu/l)	95 <sup>e</sup>	80 <sup>d</sup>	78 <sup>c</sup>	47 <sup>ab</sup>	40 <sup>a</sup>	62b		
<b>Treated twice Daily</b>								
Total protein(g/l)	60.8 <sup>c</sup>	50.4 <sup>b</sup>	76.7 <sup>e</sup>	46 <sup>a</sup>	75.5 <sup>e</sup>	65.9 <sup>d</sup>	65.1d	
Cholesterol (mg/dl)	98.2 <sup>f</sup>	75.4 <sup>b</sup>	83.6 <sup>c</sup>	92.7 <sup>e</sup>	60.9 <sup>a</sup>	87.1 <sup>d</sup>	106.9g	
SGOT (iu/l)	43 <sup>c</sup>	92 <sup>d</sup>	36 <sup>b</sup>	93 <sup>d</sup>	35 <sup>b</sup>	19 <sup>a</sup>	44c	
SGPT (iu/l)	23 <sup>c</sup>	22.5 <sup>c</sup>	15 <sup>ab</sup>	23 <sup>c</sup>	12 <sup>a</sup>	17 <sup>b</sup>	27d	
Urea (mg/dl)	22.5 <sup>a</sup>	25.05 <sup>b</sup>	27.3 <sup>c</sup>	25.1 <sup>b</sup>	24.2 <sup>ab</sup>	24.2 <sup>ab</sup>	28C	
Alkaline phosphatase (iu/l)	29 <sup>a</sup>	58 <sup>d</sup>	63 <sup>d</sup>	37 <sup>b</sup>	34 <sup>ab</sup>	60 <sup>cd</sup>	62d	

WL, water extract of leaf; F.M. L, aqueous fermented maize starch extract of leaf.

**Table 4 Phytochemical analysis of the various plant parts of *Morinda morindiodes***

Investigated Compounds	Plant Parts / Quantity of Compound		
	Leaf	Stem	Root
Alkaloid (g/100g)	1.42	1.96	1.62
Saponin (g/100g)	25.3	26.1	22.5
Tannin (mg/100g)	46.2	49.2	38.55
Flavonoid (mg/100g)	14.2	10.4	12.1
Glycosyanides (mg/100g)	<b>0.98</b>	1.06	1.12

Published: 20 October 2010

**References**

1. Abosi AO, Raseroka BH: *In vivo* antimalarial activity of *Vernonia amygdalina*. *British Journal of Biomedical Science* 2003, **60**(22):89-91.
2. Ajaiyeoba EO, Oladepo O, Fawole OI, Bolaji OM, Akinboye DO, Ogundahunsi OAT, Falade CO, Gbotosho GO, Itiola OA, Hapji TC, Ebong OO, Ononimu IM, Osowole OS, Oduola OO, Ashidi JS, Oduola AMU: Cultural categorization of febril illnesses in correlation with herbal remedies used for treatment in Southwestern Nigeria. *Journal of Ethnopharmacology* 2003, **85**:179-185.
3. Andrade-Neto VF, Brandao MG, Stehman JR, Oliveira LA, Krettli AU: Antimalarial activity of Cinchona-like plants used to treat fever and malaria in Brazil. *Journal of Ethnopharmacology* 2003, **87**:253-256.
4. Harborne JB: *Phytochemical Methods. A Guide to Modern Techniques to Plant Analysis* London, Chapman and Hall, 3rd edn 1987, 56-79.
5. Idowu OA, Soniran OT, Ajana O, Aworinde DO: *Ethnobotanical Survey of antimalarial plants used in Ogun State, Southwest Nigeria. African Journal of Pharmacy and Pharmacology* 2010, **4**:55-60.

doi:10.1186/1475-2875-9-S2-P51

**Cite this article as:** Temidayo et al.: Evaluation of *in vivo* Antiplasmodial Activities of extracts of *Morinda morindiodes* (Bak) in the treatment of malaria in Ogun State. *Malaria Journal* 2010 **9**(Suppl 2):P51.