


# The complete plastome sequence of *Monstera deliciosa* (Araceae), an ornamental foliage plant

Seona Yun<sup>a,b</sup>  and Hyeran Kim<sup>a,b,c</sup> 

<sup>a</sup>Institute of Life Sciences, Kangwon National University, Chuncheon, Republic of Korea; <sup>b</sup>Department of Biological Sciences, Kangwon National University, Chuncheon, Republic of Korea; <sup>c</sup>Interdisciplinary Graduate Program in BIT Medical Convergence, Kangwon National University, Chuncheon, Republic of Korea

## ABSTRACT

In the current study, we sequenced the complete plastome of *Monstera deliciosa* Liebm. (1849), an attractive foliage plant. The total length of the plastome of *M. deliciosa* is 163,499 bp and it consists of three distinct regions: a large single-copy (90,092 bp), a small single-copy (21,737 bp), and a pair of inverted repeats (IRs, 25,835 bp). The overall GC content is 36.2%, and the genome contains 110 functional genes, excluding pseudogenes. These functional genes encompass 77 protein-coding genes, 29 transfer RNA genes, and four ribosomal RNA genes. Notably, both the *infA* and *rpl23* genes have been identified as pseudogenes. Phylogenetic analysis based on 14 representative plastomes from seven subfamilies indicates that Monsteroideae is monophyletic and sister to Pothoideae. Furthermore, *M. deliciosa* and *M. adansonii* were shown to share a recent common ancestor, the finding for which is supported by a strong bootstrap value. The sequenced plastome of *M. deliciosa* can serve as a valuable resource for establishing phylogenetic relationships and enhancing species identification within the genus *Monstera*. In addition, it can facilitate investigations into the genetic characteristics of this plant.

## ARTICLE HISTORY

Received 21 August 2023  
Accepted 11 November 2023

## KEYWORDS

Plastome; *Monstera deliciosa*; ornamental plant; pseudogene

## Introduction

The genus *Monstera* Adans. (Araceae) is classified under the subfamily Monsteroideae and tribe Monstereae (Cabrera et al. 2008). Comprising over 60 species, the *Monstera* species is native to tropical America and grows in the regions between Mexico and Southern Brazil (Mayo et al. 1998; World Plants 2023, [www.worldplants.de](http://www.worldplants.de)). New species of *Monstera* are continuously being reported, as evidenced by studies from Zuluaga and Cameron (2018), Cedeño-Fonseca et al. (2020), and Cedeño-Fonseca et al. (2021). Despite ongoing research, the taxonomy and phylogeny of this genus remain relatively understudied. This is primarily due to the wide infraspecific variation observed within the genus, which presents challenges in species identification using traditionally diagnostic morphological features (Madison 1977). Consequently, further investigation and the development of alternative approaches to address these challenges are needed. One promising approach is the use of plastome sequences, which provides genetic insights for species delimitation and the determination of phylogenetic relationships within *Monstera*.


*Monstera deliciosa* Liebm. (1849), which originated in Central America between Mexico and Guatemala, has gained popularity as an ornamental foliage plant and has been introduced to many countries, including South Korea, Japan, and

China for the same purpose (Lee and Kim 1980; Long et al. 2017; Sakamoto et al. 2022). Recently, there has been growing interest in the chimeric forms of *M. deliciosa*, characterized by the presence of both green and albino tissues (Li et al. 2021). This intriguing phenomenon has sparked increased attention and research in horticultural genetic research (Park et al. 2023). Despite the significance of *M. deliciosa* in horticulture, there has been a lack of research utilizing the whole plastome of *M. deliciosa* as a molecular genetic resource. Therefore, the objective of this paper is to bridge this gap by providing the complete plastome sequence of *M. deliciosa*, which allows for further investigations into the genetic traits of this plant.

## Materials and methods


*Monstera deliciosa* was obtained from the Gangwon Provincial Arboretum, South Korea (37°55'22.0"N, 127°43'32.0"E, voucher number PCGE-KHR2023001, Figure 1) and grown in a growth room at Kangwon National University until needed for analysis (contact to Hyeran Kim, [ranny@kangwon.ac.kr](mailto:ranny@kangwon.ac.kr)). Genomic DNA was extracted from 100 mg of fresh leaf using the Exgene Plant SV mini kit (GenAll Biotechnology, Seoul, South Korea), following the manufacturer's instructions. The extracted DNA concentration was determined using a spectrophotometer, while

**CONTACT** Hyeran Kim  [ranny@kangwon.ac.kr](mailto:ranny@kangwon.ac.kr)  Department of Biological Science, Kangwon National University, Chuncheon 24341, Republic of Korea

 Supplemental data for this article can be accessed online at <https://doi.org/10.1080/23802359.2023.2284415>.

© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.



**Figure 1.** *Monstera deliciosa* Liebm. (A) Leaves with natural holes; (B) wrinkled geniculum at the petiole; (C) roots with aerial adventitious roots. All photos were taken by Yun at the Gangwon Provincial Arboretum, South Korea.

the integrity was confirmed through electrophoresis using a 1% agarose gel.

Sequencing was performed on the Illumina NovaSeq 6000 platform (Macrogen, Seoul, South Korea). Following this, *de novo* assembly of the plastome was conducted with NOVOplasty ver. 4.3 (Dierckx et al. 2017), using a 39 k-mer length as well as a fragment of the *rbcL* gene from *Monstera adansonii* as the seed sequence. The coverage depth of completely sequenced *M. deliciosa* was determined by performing read mapping (Figure S1). The plastome genes to be annotated were identified using CPGAVAS2 (Shi et al. 2019) and tRNAscan-SE 2.0 software with the default settings (Schattner et al. 2005). The annotation of protein-coding regions was further validated by searching for homologous genes using the NCBI Conserved Domain Database (CDD) (Marchler-Bauer et al. 2011). The circular plastome map of *M. deliciosa* was generated using the OGDRAW ver. 1.3.1 program (Greiner et al. 2019).


For phylogenetic analysis, 14 species were included, representing seven subfamilies within the Araceae family. However, the subfamily Gymnostachyoideae was excluded from the analysis due to the unavailability of plastome data. Additionally, *Alisma plantago-aquatica* from the Alismataceae family was used as an outgroup. The alignment of whole plastome sequences was carried out using MAFFT ver. 7 (Kato and Standley 2013), while gaps and poorly aligned

positions were refined using Gblocks v.0.91b (Castresana 2000). Lastly, the phylogenetic tree was constructed using the maximum-likelihood method implemented in IQ-TREE v.1.6.7 (Nguyen et al. 2015), with 1000 bootstrap replicates. The plastome sequence of *M. deliciosa* was deposited in GenBank under the accession number OR260879.

## Results

The plastome of *M. deliciosa* is 163,499 bp in size and exhibits a circular quadripartite structure consisting of a large single-copy (LSC, 90,092 bp), a small single-copy (SSC, 21,737 bp), and a pair of inverted repeat regions (IRs, each 25,835 bp) (Figure 2). The overall GC content is 36.2%, with variation observed among the different regions: 34.6% in the LSC region, 28.9% in the SSC region, and 42.0% in the IR region.

It was found that *M. deliciosa* has a total of 110 functional genes, excluding pseudogenes. These genes include 77 protein-coding genes, 29 tRNA genes, and four rRNA genes. Among these, 12 protein-coding genes and six tRNA genes possess multiple exons (Figure S2). Notably, the *infA* gene in the LSC region, as well as the *rpl23* gene in both IR regions, each exhibit one internal stop codon, implying that these genes are nonfunctional. The loss of the *infA* gene has been observed in all subfamilies other than basal group Onrntioideae.




**Figure 2.** The plastome map of *Monstera deliciosa*. Genes positioned outside the circle are transcribed in a counterclockwise direction, while genes within the inner circle are transcribed clockwise. The GC content is depicted by the dark grey inner circle.

As evidenced by the phylogenetic tree, Monsteroideae forms a monophyletic group and presents as the sister clade to Pothoideae (Figure 3). Furthermore, within the Monsteroideae clade, it can be noted that *M. deliciosa* and *M. adansonii* share a recent common ancestor. All relationships among species in the tree are strongly supported by high bootstrap values (BS = 100).

## Discussion and conclusions

This study reports the first complete plastome sequence of *M. deliciosa*. It was found that *M. deliciosa* displays the conserved quadripartite structure that has been observed in previous reports on plastome structure by Palmer (1985). In

comparison with other species within the Araceae family, *M. deliciosa* demonstrated a high level of similarity in terms of gene composition, gene order, and the number of genes (Henriquez et al. 2020a). However, in line with previous studies on Monsteroideae (Henriquez et al. 2020a, 2020b), a non-functional *infA* gene was identified as characterized by the presence of an internal stop codon in the LSC region. The loss of the *infA* gene, observed in all subfamilies other than Orontioideae, serves as a synapomorphy for the clade consisting of six subfamilies: Lemnoideae to Aroideae. Conversely, the *rpl23* gene in *M. deliciosa* was also shown to exhibit one internal stop codon, which consequently distinguishes it from other Monsteroideae species. Our phylogenetic tree is consistent with a previous study conducted by



**Figure 3.** Phylogenetic tree generated using the maximum-likelihood method implemented in IQ-TREE v.1.6.7, based on complete plastome sequences. The species sequenced in this study is indicated in red and the accession numbers of species used for phylogenetic analysis are as follows: *Alisma plantago-aquatica* (MK090659), *Orontium aquaticum* (MT226773), *Lemna minor* (DQ400350), *Spirodela polyrhiza* (MN419335), *Pothos scandens* (MN046891), *Spathiphyllum patulinervum* (MN477425), *Stenospermation multiovulatum* (MN046893), *Monstera adansonii* (MN046888), *Monstera deliciosa* (OR260879), *Lasia spinosa* (MT226772), *Zamioculcas zamiifolia* (MT226775), *Amorphophallus albus* (OP531918), *Xanthosoma sagittifolium* (MW628970), *Typhonium blumei* (MT872311), and *Arisaema heterophyllum* (ON060885).

Henriquez et al. (2014), providing further support for the monophyly of Monsteroideae and its sister relationship with Pothoideae.

Given the advantages of uniparental inheritance and the highly conserved nature of plastomes, which still provide sufficient information to capture evolutionary events (Raubeson and Jansen 2005), the complete plastome sequence of *M. deliciosa* will prove to be of considerable value in various research areas. For example, it can serve as a resource for the establishment of phylogenetic relationships within the Monsteroideae subfamily and facilitate elucidation of its evolutionary history. In addition, the plastome sequence can aid in the development of molecular markers capable of distinguishing between different *Monstera* species. In turn, these markers can contribute to more accurate species identification and subsequent classification. Ultimately, the plastome of *M. deliciosa* can facilitate investigations into its own horticultural value and genetic characteristics.

## Acknowledgements

Both authors thank the PCGE lab members, especially Yeon Hee Hyung, for their technical assistance. We thank Gangwon Provincial Arboretum for providing the sample of *Monstera deliciosa*.

## Author contributions

H.K. conceived this study and collected the sample. H.K. and S.Y. performed the experiments. S.Y. analyzed the data and wrote the manuscript. H.K. and S.Y. revised the manuscript.

## Disclosure statement

The authors declare that there is no conflict of interest regarding the publication of this article.

## Funding

This research was supported by the Basic Science Research Program, Grant Number [NRF-2022R1A2C1010869], of the National Research Foundation of Korea to H.K.

## ORCID

Seona Yun  <http://orcid.org/0000-0002-4961-7189>

Hyeran Kim  <http://orcid.org/0000-0003-4963-9510>

## Data availability statement

The genome sequence data that support the findings of this study are openly available in GenBank on the NCBI database at <https://www.ncbi.nlm.nih.gov/> under the accession no. OR260879. The associated BioProject, SRA, and Bio-Sample numbers are PRJNA992674, SRR25232409, and SAMN36379614, respectively.

## References

- Cabrera LI, Salazar GA, Chase MW, Mayo SJ, Bogner J, Dávila P. 2008. Phylogenetic relationships of aroids and duckweeds (Araceae) inferred from coding and noncoding plastid DNA. *Am J Bot.* 95(9):1153–1165. doi: [10.3732/ajb.0800073](https://doi.org/10.3732/ajb.0800073).
- Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol.* 17(4):540–552. doi: [10.1093/oxfordjournals.molbev.a026334](https://doi.org/10.1093/oxfordjournals.molbev.a026334).
- Cedeño-Fonseca M, Croat TB, Zuluaga A, Mittermeier M, Blanco MA. 2020. Two new species of *Monstera* (Araceae: Monsteroideae) from Costa Rica. *Phytotaxa.* 461(3):185–194. doi: [10.11646/phytotaxa.461.3.5](https://doi.org/10.11646/phytotaxa.461.3.5).
- Cedeño-Fonseca M, Ortiz OO, Zuluaga A, Grayum MH, Croat TB. 2021. Four new species of *Monstera* (Araceae) from Panama, including one with the largest leaves and another with the largest inflorescences in the genus. *Webbia.* 76(2):265–279. doi: [10.36253/jopt-10807](https://doi.org/10.36253/jopt-10807).
- Dierckxsens N, Mardulyn P, Smits G. 2017. NOVOPlasty: de novo assembly of organelle genomes from whole genome data. *Nucleic Acids Res.* 45(4):e18. doi: [10.1093/nar/gkw955](https://doi.org/10.1093/nar/gkw955).
- Greiner S, Lehwark P, Bock R. 2019. OrganellarGenomeDRAW (OGDRAW) version 1.3.1: expanded toolkit for the graphical visualization of organellar genomes. *Nucleic Acids Res.* 47(W1):W59–W64. doi: [10.1093/nar/gkz238](https://doi.org/10.1093/nar/gkz238).
- Henriquez CL, Arias T, Pires JC, Croat TB, Schaal BA. 2014. Phylogenomics of the plant family Araceae. *Mol Phylogenet Evol.* 75:91–102. doi: [10.1016/j.ympev.2014.02.017](https://doi.org/10.1016/j.ympev.2014.02.017).
- Henriquez CL, Abdullah , Ahmed I, Carlsen MM, Zuluaga A, Croat TB, McKain MR. 2020a. Molecular evolution of chloroplast genomes in Monsteroideae (Araceae). *Planta.* 251(3):72. doi: [10.1007/s00425-020-03365-7](https://doi.org/10.1007/s00425-020-03365-7).
- Henriquez CL, Abdullah , Ahmed I, Carlsen MM, Zuluaga A, Croat TB, McKain MR. 2020b. Evolutionary dynamics of chloroplast genomes in subfamily Aroideae (Araceae). *Genomics.* 112(3):2349–2360. doi: [10.1016/j.ygeno.2020.01.006](https://doi.org/10.1016/j.ygeno.2020.01.006).
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 30(4):772–780. doi: [10.1093/molbev/mst010](https://doi.org/10.1093/molbev/mst010).
- Lee JS, Kim MH. 1980. A study on the species of introduced landscape and cultivated plants in Jeju Island, Korea (I). *Jeju Natl Univ.* 12:97–115.
- Li F, Jin C, Zhang L, Wang J. 2021. Hyper-recombinant plants: an emerging field for plant breeding. *Crit Rev Plant Sci.* 40(5):446–458. doi: [10.1080/07352689.2021.1985819](https://doi.org/10.1080/07352689.2021.1985819).
- Long C, Fang Q, Long B, Ji Y, Shu H, Luo B, Liu B. 2017. Ornamental aroids (Araceae) in China. *Sbornik naučnyh trudov.* 145:125–131.
- Madison M. 1977. A revision of *Monstera* (Araceae). *Contrib Gray Herb Harvard Univ.* 206–208(207):3–100. doi: [10.5962/p.336443](https://doi.org/10.5962/p.336443).
- Marchler-Bauer A, Lu S, Anderson JB, Chitsaz F, Derbyshire MK, DeWeese-Scott C, Fong JH, Geer LY, Geer RC, Gonzales NR, et al. 2011. CDD: a Conserved Domain Database for the functional annotation of proteins. *Nucleic Acids Res.* 39(Database issue):D225–D229. doi: [10.1093/nar/gkq1189](https://doi.org/10.1093/nar/gkq1189).
- Mayo SJ, Bogner J, Boyce PC. 1998. Araceae. In: Kubitzki K, editor. *Flowering plants. Monocotyledons.* Berlin, Heidelberg: Springer Berlin Heidelberg; p. 26–74.
- Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol.* 32(1):268–274. doi: [10.1093/molbev/msu300](https://doi.org/10.1093/molbev/msu300).
- Palmer JD. 1985. Comparative organization of chloroplast genomes. *Annu Rev Genet.* 19(1):325–354. doi: [10.1146/annurev.ge.19.1.20185.001545](https://doi.org/10.1146/annurev.ge.19.1.20185.001545).
- Park H-S, Jeon J-H, Cho W, Lee Y, Park JY, Kim J, Park YS, Koo HJ, Kang JH, Lee TJ, et al. 2023. High-throughput discovery of plastid genes causing albino phenotypes in ornamental chimeric plants. *Hortic Res.* 10(1):uhac246. doi: [10.1093/hr/uhac246](https://doi.org/10.1093/hr/uhac246).
- Raubeson LA, Jansen RK. 2005. Chloroplast genomes of plants. In: Henry RJ, editor. *Plant diversity and evolution: genotypic and phenotypic variation in higher plants.* Wallingford (UK): CABI Publishing; p. 45–68.
- Sakamoto A, Uzuhashi S, Hoshi H, Kubota M, Horie H, Kakishima M. 2022. First report of rust caused by *Puccinia paullula* on *Monstera deliciosa* and *M. adansonii* in Japan. *Plant Dis.* 107(2):570. doi: [10.1094/PDIS-05-22-1035-PDN](https://doi.org/10.1094/PDIS-05-22-1035-PDN).
- Schattner P, Brooks AN, Lowe TM. 2005. The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res.* 33(Web Server issue):W686–W689. doi: [10.1093/nar/gki366](https://doi.org/10.1093/nar/gki366).
- Shi L, Chen H, Jiang M, Wang L, Wu X, Huang L, Liu C. 2019. CPGAVAS2, an integrated plastome sequence annotator and analyzer. *Nucleic Acids Res.* 47(W1):W65–W73. doi: [10.1093/nar/gkz345](https://doi.org/10.1093/nar/gkz345).
- World Plants. 2023. Synonymic checklist and distribution of the world flora. Version 16.1; [updated 2023 Jun 16; accessed 6 Jul 2023]. [www.worldplants.de](http://www.worldplants.de).
- Zuluaga A, Cameron KM. 2018. Two new species of *Monstera* (Araceae: Monsteroideae) with entire leaves from Panama and Costa Rica. *Phytotaxa.* 334(1):1–9. doi: [10.11646/phytotaxa.334.1.1](https://doi.org/10.11646/phytotaxa.334.1.1).