

# SCIENTIFIC REPORTS



OPEN

## Nitrogen uptake kinetics and saltmarsh plant responses to global change

Grace M. Cott<sup>1,2</sup>, Joshua S. Caplan<sup>1,3,4</sup> & Thomas J. Mozdzer<sup>1,3</sup>

Coastal wetlands are important carbon sinks globally, but their ability to store carbon hinges on their nitrogen (N) supply and N uptake dynamics of dominant plant species. In terrestrial ecosystems, uptake of nitrate ( $\text{NO}_3^-$ ) and ammonium ( $\text{NH}_4^+$ ) through roots can strongly influence N acquisition rates and their responses to environmental factors such as rising atmospheric  $\text{CO}_2$  and eutrophication. We examined the  $^{15}\text{N}$  uptake kinetics of three dominant plant species in North American coastal wetlands (*Spartina patens*,  $\text{C}_4$  grass; *Phragmites australis*,  $\text{C}_3$  grass; *Schoenoplectus americanus*,  $\text{C}_3$  sedge) under ambient and elevated  $\text{CO}_2$  conditions. We further related our results to the productivity response of these species in two long-term field experiments. *S. patens* had the greatest uptake rates for  $\text{NO}_3^-$  and  $\text{NH}_4^+$  under ambient conditions, suggesting that N uptake kinetics may underlie its strong productivity response to N in the field. Elevated  $\text{CO}_2$  increased  $\text{NH}_4^+$  and  $\text{NO}_3^-$  uptake rates for *S. patens*, but had negative effects on  $\text{NO}_3^-$  uptake rates in *P. australis* and no effects on *S. americanus*. We suggest that N uptake kinetics may explain differences in plant community composition in coastal wetlands and that  $\text{CO}_2$ -induced shifts, in combination with N proliferation, could alter ecosystem-scale productivity patterns of saltmarshes globally.

Anthropogenic activities enrich the atmosphere with  $\text{CO}_2$ , but the extent to which the biosphere can absorb this increase remains uncertain<sup>1,2</sup>. Saltmarshes play a disproportionately large role in global carbon storage, sequestering up to 87 Tg of carbon per year worldwide despite comprising less than 0.5% of the Earth's land area<sup>3</sup>. However, the capacity of a given coastal wetland to store carbon hinges primarily on its nitrogen (N) supply and the N uptake dynamics of its dominant plant species<sup>4–6</sup>. In terrestrial ecosystems including coastal wetlands, root absorption of nitrate ( $\text{NO}_3^-$ ) and ammonium ( $\text{NH}_4^+$ ) strongly influences the rate of N acquisition by plants and how this varies in response to environmental factors<sup>7–9</sup>. Nitrogen availability also constrains ecosystem productivity, with N enrichment leading to species shifts that can favor both native<sup>10</sup> and introduced species<sup>11,12</sup>. However, saltmarsh species differ in their N metabolizing activities, with rates influenced by the chemical form of N and its concentration in the substrate<sup>13–15</sup>, oxygenation of the rhizosphere<sup>16</sup>, salinity conditions<sup>17</sup>, and the presence of toxins such as sulfide<sup>18</sup>. Understanding the kinetics of root nitrogen uptake and potential differences among foundation species is therefore crucial for predicting saltmarsh ecosystems' potential for carbon storage as atmospheric  $\text{CO}_2$  levels rise.

While the effects of elevated  $\text{CO}_2$  on plant productivity are well studied in a variety of ecosystems<sup>19,20</sup> including saltmarshes<sup>21,22</sup>, our understanding of these responses is primarily based on changes in aboveground biomass,  $\text{CO}_2$  assimilation<sup>23,24</sup>, and shifts in species composition that reflect competition between plant functional groups<sup>10</sup>. Little attention has been paid to belowground mechanisms such as those associated with N acquisition or how they could be altered by global change. In an elevated  $\text{CO}_2$  environment, plants generally experience a decline in tissue N concentration<sup>25</sup>, due to either a dilution of Rubisco<sup>26–28</sup> or a reduction of transpiration-driven mass flow of N through soils<sup>29</sup>; declines have been measured despite ample N supply<sup>30,31</sup>. Recent evidence also suggests that, at least for  $\text{C}_3$  plants under  $\text{NO}_3^-$  nutrition,  $\text{CO}_2$  enrichment can decrease tissue N directly by slowing growth and inhibiting shoot  $\text{NO}_3^-$  assimilation<sup>32</sup>. However, the direct effects of elevated  $\text{CO}_2$  on N uptake

<sup>1</sup>Smithsonian Environmental Research Center, 647 Contees Wharf Road, Edgewater, Maryland, 21037, USA. <sup>2</sup>School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Fields Campus, Cork, Ireland.

<sup>3</sup>Department of Biology, Bryn Mawr College, 101 North Merion Avenue, Bryn Mawr, Pennsylvania, 19010, USA.

<sup>4</sup>Present address: Department of Landscape Architecture & Horticulture, Temple University, 580 Meetinghouse Road, Ambler, Pennsylvania, 19002, USA. Correspondence and requests for materials should be addressed to T.J.M. (email: [tmozdzer@brynmawr.edu](mailto:tmozdzer@brynmawr.edu))

have only been investigated in a small number of studies, making it difficult to generalize how different species or functional groups may adjust N uptake in response to elevated CO<sub>2</sub>.

Although photosynthetic pathways typically determine plant physiological responses to elevated CO<sub>2</sub>, the circumstances under which these physiological differences translate into a change in N uptake kinetics is not yet clear. While N acquisition is generally not affected by elevated CO<sub>2</sub> in C<sub>4</sub> plants, C<sub>3</sub> plants show variable patterns in uptake parameters under elevated CO<sub>2</sub> (e.g., V<sub>max</sub> or K<sub>m</sub>)<sup>9</sup>. For example, in studies of temperate forest trees, the effect of elevated CO<sub>2</sub> on NH<sub>4</sub><sup>+</sup> root uptake capacity was species dependent, ranging from +215% in *Acer negundo* to −40% in *Quercus macrocarpa*<sup>33</sup>. In related studies, elevated CO<sub>2</sub> increased the maximum rate of NO<sub>3</sub><sup>−</sup> uptake, specifically in *Pinus ponderosa*, *Bouteloua eriopoda* and *Pinus taeda*<sup>34–37</sup>. Other studies have found no significant effect of elevated CO<sub>2</sub> on NO<sub>3</sub><sup>−</sup> or NH<sub>4</sub><sup>+</sup> uptake rates, namely in *Pinus taeda*, *Prosopis glandulosa*, *Ceratonia siliqua*, and several herbaceous species<sup>35–40</sup>. Furthermore, effects of elevated CO<sub>2</sub> are not limited to N uptake rates. In the case of a C<sub>3</sub> tropical seagrass, *Halodule uninervis*, elevated CO<sub>2</sub> inhibited NO<sub>3</sub><sup>−</sup> assimilation and NO<sub>3</sub><sup>−</sup> nutrition alone did not enhance the CO<sub>2</sub> response<sup>41</sup>. One proposed explanation for the species-specific effects of elevated CO<sub>2</sub> on NO<sub>3</sub><sup>−</sup> assimilation in C<sub>3</sub> plants is that elevated CO<sub>2</sub> concentrations decrease photorespiration, thereby decreasing the amount of reductant (NADH) available to support NO<sub>3</sub><sup>−</sup> reduction to NO<sub>2</sub><sup>−</sup> in the first step of NO<sub>3</sub><sup>−</sup> assimilation<sup>42</sup>. In contrast, the C<sub>4</sub> carbon fixation pathway generates sufficient quantities of reductants in the cytoplasm of mesophyll cells, thus avoiding the inhibitive effect of elevated CO<sub>2</sub> on NO<sub>3</sub><sup>−</sup> assimilation<sup>43</sup>.

The physiological capacity for nitrogen uptake and assimilation may provide a key mechanistic explanation for interspecific differences in sensitivity to CO<sub>2</sub> and N addition<sup>9</sup>. For example, N uptake dynamics may explain why N addition can favor coastal wetland species that do not respond strongly to elevated CO<sub>2</sub> (i.e., *Spartina patens*), ultimately negating the enhanced productivity response at the ecosystem-level<sup>10</sup>. In the context of anthropogenically-induced changes to the carbon and nitrogen cycles in wetland ecosystems, information on physiological responses of N uptake to elevated CO<sub>2</sub> could be highly relevant for understanding these species shifts and how they influence critical ecosystem-level phenomena such as resilience to sea level rise and carbon sequestration<sup>44</sup>.

Functional taxonomic groups are often linked with suites of traits, allowing for an extrapolation of results beyond a particular ecosystem. Herbaceous-dominated systems such as grasslands, deserts, tundra, and marshes are often dominated by distinct functional groups (e.g., C<sub>3</sub> grasses), and traits associated with these functional groups may influence ecosystem-scale responses, such as shifts in net primary productivity and carbon sequestration, to interacting global change factors<sup>10</sup>. Coastal saltmarshes in North America are typically dominated by C<sub>4</sub> grasses (e.g., *Spartina patens* (Aiton) Muhl.) or C<sub>3</sub> sedges (e.g., *Schoenoplectus americanus* (Pers.) Volkart ex Schinz & R. Keller). However, an introduced lineage of the C<sub>3</sub> grass *Phragmites australis* (Cav.) Trin. ex Steud. (common reed) is invading coastal and other wetlands throughout North America<sup>45</sup>, likely altering their response to global change<sup>22</sup>.


We investigated the N uptake kinetics of three functionally distinct foundation plant species in North American coastal wetlands under ambient and elevated CO<sub>2</sub> conditions, and related these results to the growth of each species in response to global change factors with data from long-term *in situ* experiments. Specifically, we asked three questions:

- (1). Does N uptake capacity differ between *Phragmites australis*, *Schoenoplectus americanus*, and *Spartina patens*? Plants adapted to low nutrient environments typically invest considerably in belowground organs and consequently do not have a high maximum uptake capacity<sup>46</sup>. Given that *S. patens* invests in belowground organs to a lesser extent than *P. australis* and *S. americanus*, we predicted that it would have a higher maximum uptake capacity.
- (2). Does elevated CO<sub>2</sub> affect N uptake kinetics? Given the decline in tissue N status observed in many plants grown under elevated CO<sub>2</sub> despite adequate N supply, we predicted that elevated CO<sub>2</sub> would negatively affect the uptake kinetics of NO<sub>3</sub><sup>−</sup> and NH<sub>4</sub><sup>+</sup> in all three species.
- (3). Do N uptake kinetics of our species explain observations in the field? We hypothesized that patterns in N uptake kinetics would correspond to species' productivity responses to both CO<sub>2</sub> and N in long-term field experiments.

## Results

**Kinetics of NO<sub>3</sub><sup>−</sup> and NH<sub>4</sub><sup>+</sup> uptake.** We performed a series of <sup>15</sup>N uptake assays to test the hypothesis that saltmarsh species from contrasting functional groups (C<sub>4</sub> grasses, C<sub>3</sub> sedges, and C<sub>3</sub> grasses) would differ in their N uptake characteristics. In a semi-controlled outdoor setting, we presented clonally propagated plants with varying concentrations of either <sup>15</sup>NO<sub>3</sub><sup>−</sup> or <sup>15</sup>NH<sub>4</sub><sup>+</sup> and measured rates of N uptake by their root systems. All three species showed curvilinear relationships between N uptake rate (V<sub>uptake</sub>) and N concentration that closely adhered to Michaelis-Menten reaction kinetics (Fig. 1). Moreover, all three species exhibited substantially greater V<sub>uptake</sub> for NH<sub>4</sub><sup>+</sup> than NO<sub>3</sub><sup>−</sup>, with rates differing by up to a factor of 10.

There were interspecific differences in V<sub>uptake</sub> for both NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>−</sup> (Table 1). *S. patens* was primarily responsible for these differences, as it exhibited mean uptake rates up to 3 times greater than those of *P. australis* or *S. americanus* (Fig. 1a,c) and separated from both species in pairwise comparisons (Table 2). In addition, for NH<sub>4</sub><sup>+</sup>, *S. americanus* had 20–30% greater mean V<sub>uptake</sub> across the range of N concentrations than did *P. australis* (Fig. 1a). These interspecific differences also manifested in the parameter V<sub>max</sub>, the maximal uptake rate, when Michaelis-Menten curves were fit to the data; in this context, V<sub>max</sub> reflects a species' capacity for N uptake under saturating N conditions. Using bootstrapped 95% confidence intervals (CIs), we again found that *S. patens* had



**Figure 1.** Rates of  $^{15}\text{N}$  uptake by *Spartina patens*, *Phragmites australis*, and *Schoenoplectus americanus* during assays. Points are means ( $\pm$ SE) for replicate plants at the six N concentrations used; horizontal jitter has been added to reduce overlap. Shaded bands show the range of Michaelis-Menten curves corresponding to the bootstrapped 95% confidence interval for  $V_{\max}$ .

Model	Term	d.f.	F	P
$\text{NH}_4^+$ uptake	Species	2	278.38	<0.001
	$\text{CO}_2$	1	10.44	0.002
	N Conc	1	26.17	<0.001
	Species $\times$ $\text{CO}_2$	2	3.97	0.02
	Species $\times$ N Conc	2	1.74	0.18
	$\text{CO}_2 \times$ N Conc	1	0.18	0.68
	Species $\times$ $\text{CO}_2 \times$ N Conc	2	0.06	0.93
$\text{NO}_3^-$ uptake	Species	2	87.78	<0.001
	$\text{CO}_2$	1	0.05	0.83
	N Conc	1	111.94	<0.001
	Species $\times$ $\text{CO}_2$	2	15.26	<0.001
	Species $\times$ N Conc	2	8.42	<0.001
	$\text{CO}_2 \times$ N Conc	1	0.31	0.58
	Species $\times$ $\text{CO}_2 \times$ N Conc	2	0.82	0.44


**Table 1.** Results of linear modeling analysis for nitrogen uptake ( $V_{\text{uptake}}$ ).

greater  $V_{\max}$  than either of the  $\text{C}_3$  species for both  $\text{NO}_3^-$  and  $\text{NH}_4^+$  (Fig. 2a,b), and that *S. americanus* had a greater  $V_{\max}$  for  $\text{NH}_4^+$  than did *P. australis* (Fig. 2a).

We carried out an additional set of assays under elevated  $\text{CO}_2$  to determine how the N uptake kinetics of our focal species could shift under future atmospheric conditions. We found that elevated  $\text{CO}_2$  altered patterns of N uptake, though these effects were species-specific and differed by N form and concentration (Table 1, Figs 1 and 2). For  $\text{NH}_4^+$ , mean  $V_{\text{uptake}}$  increased under elevated vs. ambient  $\text{CO}_2$ , with *S. patens* showing the greatest, and statistically unequivocal, increases; the other two species exhibited non-significant trends towards increases as well (Fig. 1, Table 2). Also, *S. patens* continued to have greater  $V_{\text{uptake}}$  and  $V_{\max}$  for both  $\text{NH}_4^+$  and  $\text{NO}_3^-$  than

Species	CO <sub>2</sub> Level	NH <sub>4</sub> <sup>+</sup>			NO <sub>3</sub> <sup>-</sup>		
		Mean	SE	Group	Mean	SE	Group
<i>P. australis</i>	Ambient	9.09	2.11	a	1.60	0.24	b
	Elevated	10.29	2.04	a	0.88	0.27	a
<i>S. americanus</i>	Ambient	16.12	2.04	b	1.36	0.24	ab
	Elevated	17.76	1.98	b	1.06	0.25	ab
<i>S. patens</i>	Ambient	42.91	1.98	c	3.30	0.24	c
	Elevated	57.34	2.04	d	4.81	0.24	d


**Table 2.** Mean rates of inorganic N uptake ( $V_{\text{uptake}}$ ;  $\mu\text{mol g}^{-1} \text{h}^{-1}$ ) across N concentrations by the three focal species. Group letters that differ within an N form denote statistical separation in pairwise comparisons of means.



**Figure 2.** Median of bootstrapped estimates for the Michaelis-Menten parameters  $V_{\text{max}}$  and  $K_m$ . Error bars depict the central 95% of estimates from across all bootstrapped fits ( $n = 999$ ).

both C<sub>3</sub> species when grown under elevated CO<sub>2</sub>, and the separation between *S. americanus* and *P. australis* in these metrics was maintained under elevated CO<sub>2</sub> (Fig. 2a,b, Table 2). However, for NO<sub>3</sub><sup>-</sup>, elevated CO<sub>2</sub> induced a reduction in mean  $V_{\text{uptake}}$  for *P. australis*, such that it was not differentiable from *S. americanus* in either CO<sub>2</sub> setting (Fig. 1c vs. d, Table 2).

For NO<sub>3</sub><sup>-</sup>, interspecific differences in  $V_{\text{uptake}}$  depended on N concentrations and the CO<sub>2</sub> level (Table 2), with CO<sub>2</sub> inducing larger shifts within species at low N concentrations (Fig. 1c,d). Correspondingly, there was no evidence of CO<sub>2</sub> affecting  $V_{\text{max}}$  in any species, whereas it induced notable shifts in the Michaelis-Menten parameter  $K_m$  (Fig. 2d). In the context of this study,  $K_m$  reflects a species' affinity for an N form, with smaller values indicative of greater affinity. The shift in  $K_m$  under elevated CO<sub>2</sub> was again greatest (and statistically unequivocal) for *S. patens*; bootstrapped CIs did not overlap. *S. patens* thus had a greater affinity for NO<sub>3</sub><sup>-</sup> under elevated CO<sub>2</sub> (Fig. 2d). Although the corresponding 95% CIs for *P. australis* were partly overlapping and the three-way interaction was not significant for  $V_{\text{uptake}}$  (Table 1), our data were consistent with *P. australis* experiencing the opposite shift, namely a decrease in affinity (i.e., an increase in  $K_m$ ) for NO<sub>3</sub><sup>-</sup> under elevated CO<sub>2</sub> (Fig. 2d). The data were likewise statistically equivocal for *S. americanus* (i.e., CIs were overlapping), as were CIs for all species with respect to NH<sub>4</sub><sup>+</sup> (Fig. 2c).



**Figure 3.** Mean ( $\pm$ SE) stimulation effects (i.e., difference from ambient experimental treatments) for (a) aboveground biomass production and (b) rhizome biomass production in experimental plots Global Change Research Wetland. The three treatments were CO<sub>2</sub>, elevated atmospheric CO<sub>2</sub>; N, nitrogen fertilization; and CO<sub>2</sub> + N, both elevated CO<sub>2</sub> and N fertilization. Means are calculated using the first 5 years' worth of data from two long-term field experiments (see text).

**Growth responses to global change factors.** To determine if N uptake kinetics can explain species responses to inorganic N eutrophication in the field, we compared the results of our assays with data on biomass production (aboveground and rhizome) from two long-term field experiments in which elevated CO<sub>2</sub> and NH<sub>4</sub><sup>+</sup> were added factorially to plots in a Chesapeake Bay saltmarsh<sup>10,22</sup>. In the first five years of these experiments' lifespans, N enrichment positively stimulated aboveground biomass production by *S. patens* and *P. australis* (by 214 and 220 g m<sup>-2</sup>, respectively; Fig. 3), with stimulation defined as the absolute difference in productivity between treatment and control conditions. In contrast, *S. americanus* responded negatively to N enrichment ( $-61 \pm 27$  g m<sup>-2</sup>; mean  $\pm$  SE). Elevated CO<sub>2</sub> positively stimulated aboveground biomass production for all three species, with the inter-annual mean change being greatest for *P. australis* ( $82 \pm 26$  g m<sup>-2</sup>), intermediate for *S. americanus* ( $20 \pm 7$  g m<sup>-2</sup>), and smallest for *S. patens* ( $7 \pm 1$  g m<sup>-2</sup>; Fig. 3a). Responses to the combined treatment (elevated CO<sub>2</sub> + N) were likewise greatest for *P. australis* and *S. americanus*; they produced 202 and 161 g m<sup>-2</sup> more than they did under the control, respectively (Fig. 3a). Belowground, *P. australis* also had the strongest growth responses to all three treatments, with the largest mean stimulation to rhizome biomass production observed under elevated CO<sub>2</sub> + N ( $445 \pm 175$  g m<sup>-2</sup>, Fig. 3). N enrichment did not affect rhizome biomass stimulation of *S. americanus* ( $-10 \pm 28$  g m<sup>-2</sup>) and *S. patens* responded more strongly belowground to N enrichment ( $18 \pm 7$  g m<sup>-2</sup>) than to the other two treatments (Fig. 3).

## Discussion

Our results suggest that plant responses to interacting global change factors may be related to differences in N acquisition kinetics among plant functional groups. In a prior analysis of the native saltmarsh community's response to CO<sub>2</sub> and N at our site, C<sub>4</sub> grasses respond strongly to N addition<sup>10</sup>, demonstrating that N-induced plant community shifts can alter the ecosystem's productivity response to elevated CO<sub>2</sub>. Our data suggest that this shift may be attributable to a difference in the N uptake capacity of the dominant C<sub>3</sub> and C<sub>4</sub> species in the community (*S. americanus* and *S. patens*, respectively). A high capacity for nutrient uptake, V<sub>max</sub>, is considered to be an adaptation to nutrient rich conditions, whilst a low K<sub>m</sub> denotes a high affinity for the substrate<sup>7</sup>. Here, V<sub>max</sub> levels for NH<sub>4</sub><sup>+</sup> uptake under ambient conditions were 150% higher in *S. patens* than in *S. americanus*, indicating that it is a high-nutrient species capable of taking advantage of N enrichment (Figs 1–3). In contrast, *S. americanus* is a low nutrient specialist (evidenced by low V<sub>max</sub> and low K<sub>m</sub>), and has a limited ability to take advantage of increased soil N (Fig. 3). Furthermore, plant species with a high V<sub>max</sub> generally do not produce a high root length density and are therefore competitively inferior when nutrients in soil solution are chronically low<sup>46</sup>. Consistent with this pattern, fine root production was, on average, twice as high in stands of *S. americanus* than in stands of *S. patens* over the past 20 years<sup>23</sup>. Given this, the divergent response of these two North American wetland species to elevated N is likely attributable to differences in their N uptake kinetics, and can therefore be used in a predictive framework to project plant community shifts in response to global change.

Plasticity in N uptake physiology may explain the ability of *P. australis* to thrive in both resource-poor and resource-rich habitats. For example, our results and those of a prior study<sup>47</sup> suggest that *P. australis* is adapted to a low N environment, given its low V<sub>max</sub>. However, intermediate V<sub>max</sub> levels have been measured in *P. australis*<sup>13,48</sup>, as have levels an order of magnitude greater than we found<sup>49</sup>. As suggested by Romero *et al.*<sup>49</sup>, the ammonium uptake kinetics of *P. australis* seem to be plastic, such that they can be modified in response to nutrient availability or CO<sub>2</sub> availability. The plastic response of N uptake to varying CO<sub>2</sub> and N levels in the field study may partly

explain why introduced *P. australis* can thrive under both high and low nutrient environments<sup>50</sup>. Our results provide evidence that *P. australis* has the kinetic parameters needed to invade low nitrogen environments, whilst our long-term field study shows that the species can thrive in resource rich environments. Furthermore, our long-term study clearly shows that *P. australis* can take advantage of both CO<sub>2</sub> and N, with aboveground biomass stimulated most strongly by N addition, and belowground biomass stimulated most strongly by CO<sub>2</sub> + N (Fig. 3).

Elevated CO<sub>2</sub> affected saltmarsh functional groups differently. Both C<sub>3</sub> species (*P. australis* and *S. americanus*) exhibited a trend for lower affinity for NO<sub>3</sub><sup>-</sup> under elevated CO<sub>2</sub> conditions, evidenced by increases in K<sub>m</sub> values. The functional group-specific effect of elevated CO<sub>2</sub> on NO<sub>3</sub><sup>-</sup> uptake capacity corresponds to differences in NO<sub>3</sub><sup>-</sup> assimilation previously reported by Bloom *et al.*<sup>32,43</sup> and may be attributable to the reduction in photorespiration that C<sub>3</sub> plants experience under elevated CO<sub>2</sub> conditions, as this decreases the reductant available to power the first step of NO<sub>3</sub><sup>-</sup> assimilation<sup>42</sup>. Conversely, evidence of this repression was not observed in the C<sub>4</sub> species, for which K<sub>m</sub> values actually decreased under elevated CO<sub>2</sub>. Again, elevated CO<sub>2</sub> conditions appeared to reduce NO<sub>3</sub><sup>-</sup> assimilation in C<sub>3</sub>, but not C<sub>4</sub>, plants.

It is now well established that the active process involved in ion uptake by plant roots at relatively low nutrient levels (10–200 μM) is provided by the high affinity transport system (HATS)<sup>8,51,52</sup>. This transport system is used by plants growing in natural and semi-natural ecosystems<sup>9,53</sup> and is likely the one operating at the concentrations observed in our field experiment<sup>54</sup>. The HATS for both NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> is subject to regulation in response to changes in external N availability or in the N demand of the whole plant<sup>55</sup>. However, the mechanisms underlying the suppression of N uptake under elevated CO<sub>2</sub> remain unclear.

We found that elevated CO<sub>2</sub> enhanced the physiological capabilities of our C<sub>4</sub> species, such as increasing V<sub>uptake</sub> of NH<sub>4</sub><sup>+</sup>. This suggests that some C<sub>4</sub> plants may become more competitive for N with near-future global change. Furthermore, this taxa-specific nutrient uptake response to elevated CO<sub>2</sub> may influence differences in growth rate, as rates of N acquisition are often positively correlated with growth rates<sup>56–58</sup>. Indeed, *S. patens* had the kinetic parameters of an exploitative, fast growing species<sup>59</sup>, and data from the long-term field experiment show that its shoot biomass response to elevated N was on par with that of the invasive species *P. australis*. These physiological enhancements induced by elevated CO<sub>2</sub> may also explain the stimulation effects of CO<sub>2</sub> observed in C<sub>4</sub> species at a 30 year experiment at our field site<sup>23,60</sup>. However, rapid NH<sub>4</sub><sup>+</sup> uptake does not necessarily translate into rapid growth; Zerihun & BassiriRad<sup>33</sup> found that the relative growth rate of *Acer negundo* was unaffected by high CO<sub>2</sub> despite experiencing a two-fold increase in root NH<sub>4</sub><sup>+</sup> uptake capacity in response to high CO<sub>2</sub>.

Elevated CO<sub>2</sub> appeared to repress NH<sub>4</sub><sup>+</sup> uptake affinity in our dominant C<sub>3</sub> species; this trend may help explain long-term observations in our field experiment. The increasing K<sub>m</sub> values for NH<sub>4</sub><sup>+</sup> under elevated CO<sub>2</sub> could partly explain the reduction of tissue N levels experienced by foundational saltmarsh plants<sup>60</sup>. In some species, such as wheat, the reduction occurs despite the supply of high doses of nitrogen<sup>30</sup> indicating that the observed reduction in tissue nitrogen with elevated CO<sub>2</sub> is not due to a low nitrogen concentration in the root medium, but is related to aspects of uptake itself. Indeed, N addition did not sustain the initial positive CO<sub>2</sub> stimulation of C<sub>3</sub> biomass in one of our *in situ* experiments<sup>10</sup>. This was partially explained by competition with *S. patens*, but may also be attributable to sustained CO<sub>2</sub> enrichment having a gradual decreasing effect on NH<sub>4</sub><sup>+</sup> uptake capacity. In addition, the combined effects of N and CO<sub>2</sub> on *P. australis* shoot biomass was smaller than the effect of N alone, suggesting a potential negative effect. Another contributing factor may indeed be the reduction of transpiration-driven mass flow of N through soils due to a reduction in stomatal conductance usually experienced by plants under elevated CO<sub>2</sub> conditions<sup>29</sup>. Alternatively, the pattern may derive from reductions in root respiration, given that greater tissue N content entails greater maintenance respiration<sup>61</sup> and the fact that the energy requirements for NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> uptake and assimilation constitute a significant portion of root respiration<sup>62</sup>. Reductions in root respiration as a result of elevated CO<sub>2</sub> exposure have been reported in the literature<sup>63</sup> but no satisfactory mechanisms to explain these effects have been demonstrated<sup>64,65</sup>.

As is the case for NO<sub>3</sub><sup>-</sup> uptake, the mechanisms underlying the suppression of NH<sub>4</sub><sup>+</sup> uptake under elevated CO<sub>2</sub> remain uncertain. What is clear is that carbon metabolites such as glutamine can suppress the expression of genes associated with the HATS for both NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup><sup>66–68</sup>. Additionally, glucose supply to plant roots can inhibit the induction of some enzymatic proteins such as glutamate dehydrogenase and asparagine synthetase<sup>69</sup>, both of which are involved in N metabolism<sup>70</sup>. The fact that C and N metabolism are tightly linked is inescapable<sup>71</sup>, and it may be the case that increased carbohydrate supply to roots as a result of elevated CO<sub>2</sub> exposure may act directly or indirectly on plant nitrogen pools, ultimately causing a downregulation of genes associated with N uptake.

The extent to which N uptake is influenced by edaphic factors such as oxygenation of the rhizosphere, salinity conditions, or sulfide concentration was not investigated in this study. To evaluate uptake free of these effects, assays were conducted in aerobic solutions free from Na or hydrogen sulfide. Salinity is known to inhibit N uptake in *Spartina alterniflora* and *P. australis* by up to 40%<sup>13,17,48</sup>, especially at levels above 20 ppt. Similarly, anoxic conditions and hydrogen sulfide can inhibit N uptake<sup>18,48</sup>. How these factors influence N uptake in the field is unknown, although *S. americanus* has the ability to oxygenate its rhizosphere and can tolerate frequent flooding whilst *S. patens* inhabits higher saltier zones<sup>72</sup>. Therefore each species is specifically adapted to tolerate one of these confounding factors.

The interspecific differences in N uptake kinetics identified here provide an explanation for how individual plant-level responses to global change factors (such as CO<sub>2</sub> and N enrichment) translate into species dynamics at a community level. We suggest that the ecosystem-level response to interacting global change factors can be related to the root uptake kinetics of N acquisition by different plant functional groups. Our results further demonstrate that *P. australis* is capable of invading low nitrogen ecosystems, whilst our long-term field study shows that it can also thrive in resource rich environments. Consequently, physiological plasticity in the invasive species appears to facilitate its proliferation. Further study is required to determine if rising atmospheric CO<sub>2</sub> levels can be expected to repress N uptake in other ecosystems and to examine the specific mechanisms involved.

## Methods

**Nitrogen uptake assays.** Three wetland taxa were selected for this study: *Schoenoplectus americanus*, *Spartina patens*, and a lineage of *Phragmites australis* subsp. *australis* (haplotype M). All three are highly abundant in saltmarshes along the Atlantic Coast of North America and are representative of the plant functional groups that dominate tidal marshes, namely C<sub>3</sub> sedges, C<sub>4</sub> grasses, and C<sub>3</sub> grasses, respectively. *P. australis* subsp. *australis* is both introduced and invasive in North America<sup>73</sup>, while the two natives are dominant species in two long-term experiments situated at the Smithsonian Environmental Research Center (SERC) in Maryland, USA.

The nutrient uptake experiment was conducted in a set of six chambers (1.0 × 0.7 × 1.0 m) located at Bryn Mawr College in Pennsylvania, USA (40.0297°N, 75.3139°W). During the course of the experiment, plants experienced natural temperature fluctuations, with a mean daily high of 29.8 ± 0.8 °C and a mean daily low of 19.5 ± 0.5 °C. The chambers had closed walls constructed of Lexan polycarbonate, though they were not air-tight. Blowers continuously moved air into chambers at a rate that replaced the volume of each chamber once approximately every two minutes. Three chambers were maintained at ambient CO<sub>2</sub> and three at elevated CO<sub>2</sub> (ambient + 300 ppmv CO<sub>2</sub>). CO<sub>2</sub> concentrations in the chambers were monitored with CM-0212 CO<sub>2</sub> loggers (CO<sub>2</sub> Meter, Ormond Beach, USA) and adjusted manually on a daily basis.

Plant material was collected in the spring of 2012 from SERC, maintained for one year in the Bryn Mawr College greenhouse, and propagated from rhizome fragments or emergent shoots in May 2013. Propagules were washed clean of organic matter and dead root material, and individual shoots were placed in square pots (10 cm sides) filled with clean sand to facilitate transfer to a hydroponic medium during N uptake assays. Thirteen plants per species were placed in each chamber in June 2013 (n = 234 total plants) and fertilized weekly with a 1/10<sup>th</sup> strength Hoaglands solution. Within 10 weeks, individual plants achieved a root mass suitable for assays (>100 mg dw).

To investigate NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> uptake kinetics, we presented individual plants with a <sup>15</sup>N-labeled substrate in hydroponic solution. The protocol for assays was adapted from Epstein *et al.*<sup>74</sup> and Mozdzer *et al.*<sup>13</sup>. Briefly, plants were washed free of sand and placed in an N-free solution of 0.50 mM CaCl<sub>2</sub> overnight to maintain root epidermal cell integrity. After equilibration, each plant was exposed to one of six different N concentrations (5, 10, 25, 50, 100, and 500 μM) of either <sup>15</sup>NH<sub>4</sub>Cl or K<sup>15</sup>NO<sub>3</sub>, respectively (99% enriched; Cambridge Isotope Laboratories, Andover, USA) for 45 minutes in a well-mixed 0.50 mM CaCl<sub>2</sub> solution. To ensure that drawdown would not exceed 10% of the starting concentration, the reaction volume for assays was adjusted to 2500 ml for the lowest two concentrations and 1000 ml for the remaining concentrations. The treatment assay solution was identical to the equilibration medium but contained the labeled N dose. Each exposure series for both forms of N was applied to the three species in each chamber, such that the complete set of assays was performed in triplicate (n = 216 plants). One additional plant per species from each chamber was exposed only to the equilibration medium as a control (n = 18 plants). After 45 minutes of exposure, roots were rinsed for 2 min with 1 mM KCl to remove any excess labeled substrate from root surfaces. Each plant was then separated into root, rhizome, and stem tissue and dried at 60 °C to constant weight. Dry tissue was ground using a Retsch Mixer Mill 400 (Verder Scientific, Haan, Germany). To minimize potential effects of diurnal variation in nutrient uptake, assays were conducted at approximately the same time each day (1000–1200 h) over the course of three weeks, with the three exposure series for one plant species, one N form, and one CO<sub>2</sub> level (n = 18 plants) completed per day. Samples of root tissue were analyzed for <sup>15</sup>N using a Europa Integra continuous flow mass spectrometer (UC Davis Stable Isotope Facility).

Uptake rates of <sup>15</sup>N (V<sub>uptake</sub>) for individual plants were calculated from the mass of <sup>15</sup>N that they assimilated (m<sub>assim</sub>, in μg)<sup>13,75</sup>:

$$m_{\text{assim}} = \frac{m_1(APE_{\text{samp}} - APE_{\text{ctrl}})}{APE_{\text{treat}}} \quad (1)$$

$$V_{\text{uptake}} = \frac{(m_{\text{assim}}/MW_{\text{treat}})}{(m_2 t_{\text{exp}})} \quad (2)$$

where  $m_1$  is the mass of N in the sample (in μg), APE<sub>samp</sub> is the atom % excess <sup>15</sup>N of the root sample exposed to a labeled substrate, APE<sub>ctrl</sub> is the atom % excess <sup>15</sup>N in the control root sample, APE<sub>treat</sub> is the atom % excess of the labeled <sup>15</sup>N treatment, MW is the molecular weight of the N isotope,  $m_2$  is the dry root mass of the sample (in grams), and  $t_{\text{exp}}$  is the duration of the exposure to labeled substrate (in minutes). Several uptake rates were anomalously high, especially at low N concentrations (5–25 μM). This was probably due to carryover during mass spectrometry, so V<sub>uptake</sub> values that were greater than those at both of the next two higher N concentrations within a series were omitted (n = 19).

The <sup>15</sup>N uptake rates from each exposure series (n = 14–18 plants) were then fit to the Michaelis-Menten equation in order to derive values of maximal uptake rate (V<sub>max</sub>) and the substrate concentration at which the rate is 50% of V<sub>max</sub> (K<sub>m</sub>):

$$V_{\text{uptake}} = \frac{V_{\text{max}}[c]}{K_m + [c]} \quad (3)$$

where [c] is the concentration of NH<sub>4</sub><sup>+</sup> or NO<sub>3</sub><sup>-</sup>. V<sub>max</sub> (in μmol <sup>15</sup>N g<sup>-1</sup> h<sup>-1</sup>) provides a measure of uptake capacity under saturating N conditions, while K<sub>m</sub> (in μM of NH<sub>4</sub><sup>+</sup> or NO<sub>3</sub><sup>-</sup>) provides an estimate of the species' affinity for NH<sub>4</sub><sup>+</sup> or NO<sub>3</sub><sup>-</sup>; smaller values correspond to greater affinity. Curve fitting was carried out in R using a self-starting non-linear regression function (*SSmicmen* from the *nlstools* library). To determine if there were differences among species and/or CO<sub>2</sub> levels, we used bootstrapping to compute 95% confidence intervals for all

parameter estimates (via *nlsBoot*, also from *nlstools*;  $n = 999$  iterations). Estimates were considered different if there was no overlap between pairs of bootstrapped 95% confidence intervals<sup>76</sup>.

Linear models were used to determine how experimental factors ( $\text{CO}_2$  level, plant species, and N concentration) affected N uptake rates ( $V_{\text{uptake}}$ ), with separate models fit to data for  $\text{NO}_3^-$  and  $\text{NH}_4^+$ . Both models had the same form, with species and  $\text{CO}_2$  level included as categorical variables but N concentration included as a continuous variable. Terms for all possible two and three way interactions were also included.  $V_{\text{uptake}}$  values were square root transformed to ensure residual normality. Tukey-adjusted pairwise comparisons were subsequently made among all species- $\text{CO}_2$  level combinations; the family-wise error rate was held at 0.05. All statistical analyses were conducted in R version 3.2.3.

**Long-term field experiments.** We compared results from our *ex-situ* kinetic experiment with *in situ* data from two long-term experiments situated in a brackish tidal marsh within SERC's Global Change Research Wetland (Kirkpatrick Marsh; 38.8742° N, 76.5474° W) in Edgewater, Maryland, USA. The first experiment was established in 2006 and examines the effect of elevated  $\text{CO}_2$  and mineral N addition on the dominant native saltmarsh species *S. americanus* and *S. patens*. The second study was established in 2011 and examines the effect of identical global change manipulations on the introduced lineage of *P. australis* (haplotype M) and its encroachment into the native marsh community. For details of chamber setup and experimental design see Langley and Megonigal<sup>10</sup> and Caplan *et al.*<sup>22</sup>. Briefly, half of the plots in each experiment are fertilized with  $\text{NH}_4\text{Cl}$  at a rate of  $25 \text{ g N m}^{-2} \text{ yr}^{-1}$  and half of the plots at each N level are fumigated with sufficient  $\text{CO}_2$  to raise the atmospheric concentration within open-top chambers by approximately 300 ppm throughout the growing season (May through November).

Aboveground biomass is estimated in both experiments in late July or early August of each year. For *S. americanus* and *P. australis*, this entails combining stem density counts with measurements of stem height and width that are converted to dry mass using species-specific allometric relationships<sup>77</sup>. For *S. patens*, biomass is measured directly by clipping samples within each chamber. Rhizome productivity is estimated from annual samples collected each year using ingrowth cores, with three placed in each plot for the native marsh study and six placed in each plot for the *P. australis* study.

Biomass data from the two long-term field experiments were used to quantify productivity responses to global change factors. Specifically, we calculated stimulation effects (i.e., differences from ambient) for the elevated  $\text{CO}_2$  treatment, the N enrichment treatment, and the combination treatment for both aboveground biomass and rhizome biomass. For the native marsh study, we used biomass data from *S. americanus* and *S. patens* spanning the first five years that data were available (2006–2010 for aboveground biomass and 2007–2011 for rhizome biomass). Biomass data for *P. australis* came from the second study, but likewise spanning the first five years of its lifespan (2011–2015).

**Data availability.** The datasets used in this study are available from the corresponding author on reasonable request.

## References

- Solomon, S. *et al.* The physical science basis. *Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change*, 235–337 (2007).
- Luo, Y. Terrestrial carbon-cycle feedback to climate warming. *Annual Review of Ecology, Evolution, and Systematics* **38**, 683–712 (2007).
- Mcleod, E. *et al.* A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering  $\text{CO}_2$ . *Frontiers in Ecology and the Environment* **9**, 552–560 (2011).
- Berntson, G., Rajakaruna, N. & Bazzaz, F. Growth and nitrogen uptake in an experimental community of annuals exposed to elevated atmospheric  $\text{CO}_2$ . *Global Change Biology* **4**, 607–626 (1998).
- Oren, R. *et al.* Soil fertility limits carbon sequestration by forest ecosystems in a  $\text{CO}_2$ -enriched atmosphere. *Nature* **411**, 469–472 (2001).
- Hungate, B. A., Dukes, J. S., Shaw, M. R., Luo, Y. & Field, C. B. Nitrogen and climate change. *Science* **302**, 1512–1513 (2003).
- Chapin, F. S. III. The mineral nutrition of wild plants. *Annual Review of Ecology and Systematics* **11**, 233–260 (1980).
- Forde, B. G. & Clarkson, D. T. Nitrate and ammonium nutrition of plants: physiological and molecular perspectives. *Advances in Botanical Research* **30**, 1–90 (1999).
- Bassirirad, H. Kinetics of nutrient uptake by roots: responses to global change. *New Phytologist* **147**, 155–169 (2000).
- Langley, J. A. & Megonigal, J. P. Ecosystem response to elevated  $\text{CO}_2$  levels limited by nitrogen-induced plant species shift. *Nature* **466**, 96–99 (2010).
- Bertness, M. D., Ewanchuk, P. J. & Silliman, B. R. Anthropogenic modification of New England salt marsh landscapes. *Proceedings of the National Academy of Sciences* **99**, 1395–1398 (2002).
- King, R. S., Deluca, W. V., Whigham, D. F. & Marra, P. P. Threshold effects of coastal urbanization on *Phragmites australis* (common reed) abundance and foliar nitrogen in Chesapeake Bay. *Estuaries and Coasts* **30**, 469–481 (2007).
- Mozdzer, T. J., Zieman, J. C. & McGlathery, K. J. Nitrogen uptake by native and invasive temperate coastal macrophytes: importance of dissolved organic nitrogen. *Estuaries and Coasts* **33**, 784–797 (2010).
- Mozdzer, T., Kirwan, M., McGlathery, K. & Zieman, J. Nitrogen uptake by the shoots of smooth cordgrass *Spartina alterniflora*. *Marine Ecology Progress Series* **433**, 43–52 (2011).
- Cott, G. M., Chapman, D. V. & Jansen, M. A. Differences in nitrogen-assimilating enzyme activity in halophyte species are habitat-related. *Journal of Plant Nutrition and Soil Science* **177**, 705–713 (2014).
- Morris, J. T. & Dacey, J. W. Effects of  $\text{O}_2$  on ammonium uptake and root respiration by *Spartina alterniflora*. *American Journal of Botany*, 979–985 (1984).
- Morris, J. T. Effects of oxygen and salinity on ammonium uptake by *Spartina alterniflora* Loisel. and *Spartina patens* (Aiton) Muhl. *Journal of Experimental Marine Biology and Ecology* **78**, 87–98 (1984).
- Bradley, P. M. & Morris, J. T. Influence of oxygen and sulfide concentration on nitrogen uptake kinetics in *Spartina alterniflora*. *Ecology* **71**, 282–287 (1990).
- Ainsworth, E. A. & Long, S. P. What have we learned from 15 years of free-air  $\text{CO}_2$  enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising  $\text{CO}_2$ . *New Phytologist* **165**, 351–372 (2005).
- Talhelm, A. F. *et al.* Elevated carbon dioxide and ozone alter productivity and ecosystem carbon content in northern temperate forests. *Global Change Biology* **20**, 2492–2504 (2014).



21. Drake, B. G. Rising sea level, temperature, and precipitation impact plant and ecosystem responses to elevated CO<sub>2</sub> on a Chesapeake Bay wetland: review of a 28-year study. *Global Change Biology* **20**, 3329–3343 (2014).
22. Caplan, J. S., Hager, R. N., Megonigal, J. P. & Mozdzer, T. J. Global change accelerates carbon assimilation by a wetland ecosystem engineer. *Environmental Research Letters* **10**, 115006; <https://doi.org/10.1088/1748-9326/10/11/115006> (2015).
23. Erickson, J. E., Megonigal, J. P., Peresta, G. & Drake, B. G. Salinity and sea level mediate elevated CO<sub>2</sub> effects on C<sub>3</sub>–C<sub>4</sub> plant interactions and tissue nitrogen in a Chesapeake Bay tidal wetland. *Global Change Biology* **13**, 202–215 (2007).
24. Erickson, J. E., Peresta, G., Montovan, K. J. & Drake, B. G. Direct and indirect effects of elevated atmospheric CO<sub>2</sub> on net ecosystem production in a Chesapeake Bay tidal wetland. *Global Change Biology* **19**, 3368–3378 (2013).
25. Taub, D. R. & Wang, X. Why are nitrogen concentrations in plant tissues lower under elevated CO<sub>2</sub>? A critical examination of the hypotheses. *Journal of Integrative Plant Biology* **50**, 1365–1374 (2008).
26. Wong, S.-C. Elevated atmospheric partial pressure of CO<sub>2</sub> and plant growth: II. Non-structural carbohydrate content in cotton plants and its effect on growth parameters. *Photosynthesis Research* **23**, 171–180 (1990).
27. Kuehny, J. S., Peet, M. M., Nelson, P. V. & Willits, D. H. Nutrient dilution by starch in CO<sub>2</sub>-enriched *Chrysanthemum*. *Journal of Experimental Botany* **42**, 711–716 (1991).
28. Gifford, R. M., Barrett, D. J. & Lutz, J. L. The effects of elevated [CO<sub>2</sub>] on the C:N and C:P mass ratios of plant tissues. *Plant and Soil* **224**, 1–14 (2000).
29. McDonald, E. P., Erickson, J. E. & Kruger, E. L. Research note: Can decreased transpiration limit plant nitrogen acquisition in elevated CO<sub>2</sub>? *Functional Plant Biology* **29**, 1115–1120 (2002).
30. Hocking, P. & Meyer, C. Effects of CO<sub>2</sub> enrichment and nitrogen stress on growth, and partitioning of dry matter and nitrogen in wheat and maize. *Functional Plant Biology* **18**, 339–356 (1991).
31. Kimball, B. *et al.* Elevated CO<sub>2</sub>, drought and soil nitrogen effects on wheat grain quality. *New Phytologist* **150**, 295–303 (2001).
32. Bloom, A. J., Burger, M., Asensio, J. S. R. & Cousins, A. B. Carbon dioxide enrichment inhibits nitrate assimilation in wheat and *Arabidopsis*. *Science* **328**, 899–903 (2010).
33. Zerihun, A. & Bassirirad, H. Interspecies variation in nitrogen uptake kinetic responses of temperate forest species to elevated CO<sub>2</sub>: potential causes and consequences. *Global Change Biology* **7**, 211–222 (2001).
34. Bassirirad, H., Griffin, K. L., Strain, B. R. & Reynolds, J. F. Effects of CO<sub>2</sub> enrichment on growth and root <sup>15</sup>NH<sub>4</sub> uptake rate of loblolly pine and ponderosa pine seedlings. *Tree Physiology* **16**, 957–962 (1996).
35. Bassirirad, H., Thomas, R., Reynolds, J. & Strain, B. Differential responses of root uptake kinetics of NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> to enriched atmospheric CO<sub>2</sub> concentration in field-grown loblolly pine. *Plant, Cell & Environment* **19**, 367–371 (1996).
36. Bassirirad, H., Griffin, K. L., Reynolds, J. F. & Strain, B. R. Changes in root NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> absorption rates of loblolly and ponderosa pine in response to CO<sub>2</sub> enrichment. *Plant and Soil* **190**, 1–9 (1997).
37. Bassirirad, H., Reynolds, J., Virginia, R. & Brunelle, M. Growth and root NO<sub>3</sub><sup>-</sup> and PO<sub>4</sub><sup>3-</sup> uptake capacity of three desert species in response to atmospheric CO<sub>2</sub> enrichment. *Functional Plant Biology* **24**, 353–358 (1997).
38. Newbery, R., Wolfenden, J., Mansfield, T. & Harrison, A. Nitrogen, phosphorus and potassium uptake and demand in *Agrostis capillaris*: the influence of elevated CO<sub>2</sub> and nutrient supply. *New Phytologist* **130**, 565–574 (1995).
39. Jackson, R. & Reynolds, H. Nitrate and ammonium uptake for single- and mixed-species communities grown at elevated CO<sub>2</sub>. *Oecologia* **105**, 74–80 (1996).
40. Cruz, C., Lips, S. & Martins-Louçã, M. Changes in the morphology of roots and leaves of carob seedlings induced by nitrogen source and atmospheric carbon dioxide. *Annals of Botany* **80**, 817–823 (1997).
41. Ow, Y. X. *et al.* Nitrate fertilisation does not enhance CO<sub>2</sub> responses in two tropical seagrass species. *Scientific Reports* **6**, 23093; [10.1038/srep23093](https://doi.org/10.1038/srep23093) (2016).
42. Igamberdiev, A. U., Bykova, N. V., Lea, P. J. & Gardeström, P. The role of photorespiration in redox and energy balance of photosynthetic plant cells: a study with a barley mutant deficient in glycine decarboxylase. *Physiologia Plantarum* **111**, 427–438 (2001).
43. Bloom, A. J. *et al.* CO<sub>2</sub> enrichment inhibits shoot nitrate assimilation in C<sub>3</sub> but not C<sub>4</sub> plants and slows growth under nitrate in C<sub>3</sub> plants. *Ecology* **93**, 355–367 (2012).
44. Kirwan, M. L. & Megonigal, J. P. Tidal wetland stability in the face of human impacts and sea-level rise. *Nature* **504**, 53–60 (2013).
45. Saltonstall, K. Cryptic invasion by a non-native genotype of the common reed, *Phragmites australis*, into North America. *Proceedings of the National Academy of Sciences* **99**, 2445–2449 (2002).
46. Craine, J. M. *Resource strategies of wild plants*. (Princeton University Press, 2009).
47. Tylova-Munzarova, E., Lorenzen, B., Brix, H. & Votrubova, O. The effects of NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> on growth, resource allocation and nitrogen uptake kinetics of *Phragmites australis* and *Glyceria maxima*. *Aquatic Botany* **81**, 326–342 (2005).
48. Chambers, R. M., Mozdzer, T. J. & Ambrose, J. C. Effects of salinity and sulfide on the distribution of *Phragmites australis* and *Spartina alterniflora* in a tidal saltmarsh. *Aquatic Botany* **62**, 161–169 (1998).
49. Romero, J. A., Brix, H. & Comin, F. A. Interactive effects of N and P on growth, nutrient allocation and NH<sub>4</sub><sup>+</sup> uptake kinetics by *Phragmites australis*. *Aquatic Botany* **64**, 369–380 (1999).
50. Mozdzer, T. J. & Megonigal, J. P. Jack-and-master trait responses to elevated CO<sub>2</sub> and N: a comparison of native and introduced *Phragmites australis*. *PLoS One* **7**, e42794; <https://doi.org/10.1371/journal.pone.0042794> (2012).
51. Kronzucker, H., Glass, A., Siddiqi, M. & Kirk, G. Comparative kinetic analysis of ammonium and nitrate acquisition by tropical lowland rice: implications for rice cultivation and yield potential. *New Phytologist* **145**, 471–476 (2000).
52. Min, X., Siddiqi, M. Y., Guy, R. D., Glass, A. D. & Kronzucker, H. J. A comparative kinetic analysis of nitrate and ammonium influx in two early-successional tree species of temperate and boreal forest ecosystems. *Plant, Cell & Environment* **23**, 321–328 (2000).
53. Maire, V., Gross, N., da Silveira Pontes, L., Picon-Cochard, C. & Soussana, J. F. Trade-off between root nitrogen acquisition and shoot nitrogen utilization across 13 co-occurring pasture grass species. *Functional Ecology* **23**, 668–679 (2009).
54. Mozdzer, T. J., Langley, J. A., Mueller, P. & Megonigal, J. P. Deep rooting and global change facilitate spread of invasive grass. *Biological Invasions* **18**, 2619–2631 (2016).
55. Nacry, P., Bouguyon, E. & Gojon, A. Nitrogen acquisition by roots: physiological and developmental mechanisms ensuring plant adaptation to a fluctuating resource. *Plant and Soil* **370**, 1–29 (2013).
56. Clarkson, D., Jones, L. & Purves, J. Absorption of nitrate and ammonium ions by *Lolium perenne* from flowing solution cultures at low root temperatures. *Plant, Cell & Environment* **15**, 99–106 (1992).
57. Schenk, M. Regulation of nitrogen uptake on the whole plant level. *Plant and Soil* **181**, 131–137 (1996).
58. Caplan, J. S. *et al.* Nutrient foraging strategies are associated with productivity and population growth in forest shrubs. *Annals of Botany* **119**, 977–988 (2017).
59. Grassein, F. *et al.* Relationships between functional traits and inorganic nitrogen acquisition among eight contrasting European grass species. *Annals of Botany* **115**, 107–115 (2015).
60. Curtis, P. S., Drake, B. G. & Whigham, D. F. Nitrogen and carbon dynamics in C<sub>3</sub> and C<sub>4</sub> estuarine marsh plants grown under elevated CO<sub>2</sub> in situ. *Oecologia* **78**, 297–301 (1989).
61. Hymus, G. J., Snead, T. G., Johnson, D. P., Hungate, B. A. & Drake, B. G. Acclimation of photosynthesis and respiration to elevated atmospheric CO<sub>2</sub> in two Scrub Oaks. *Global Change Biology* **8**, 317–328 (2002).
62. Bloom, A. J., Sukrapanna, S. S. & Warner, R. L. Root respiration associated with ammonium and nitrate absorption and assimilation by barley. *Plant Physiology* **99**, 1294–1301 (1992).

63. Poorter, H., Gifford, R. M., Kriedemann, P. E. & Wong, S. C. A quantitative-analysis of dark respiration and carbon content as factors in the growth-response of plants to elevated CO<sub>2</sub>. *Australian Journal of Botany* **40**, 501–513 (1992).
64. Drake, B. G. *et al.* Does elevated atmospheric CO<sub>2</sub> concentration inhibit mitochondrial respiration in green plants? *Plant, Cell & Environment* **22**, 649–657 (1999).
65. Smith, N. G. & Dukes, J. S. Plant respiration and photosynthesis in global-scale models: incorporating acclimation to temperature and CO<sub>2</sub>. *Global Change Biology* **19**, 45–63 (2013).
66. Rawat, S. R., Silim, S. N., Kronzucker, H. J., Siddiqi, M. Y. & Glass, A. D. AtAMT1 gene expression and NH<sub>4</sub><sup>+</sup> uptake in roots of *Arabidopsis thaliana*: evidence for regulation by root glutamine levels. *The Plant Journal* **19**, 143–152 (1999).
67. Zhuo, D., Okamoto, M., Vidmar, J. J. & Glass, A. D. Regulation of a putative high-affinity nitrate transporter (Nrt2; 1At) in roots of *Arabidopsis thaliana*. *The Plant Journal* **17**, 563–568 (1999).
68. Glass, A. D. *et al.* The regulation of nitrate and ammonium transport systems in plants. *Journal of Experimental Botany* **53**, 855–864 (2002).
69. Oaks, A. & Hirel, B. Nitrogen metabolism in roots. *Annual Review of Plant Physiology* **36**, 345–365 (1985).
70. Good, A. G., Shrawat, A. K. & Muench, D. G. Can less yield more? Is reducing nutrient input into the environment compatible with maintaining crop production? *Trends in Plant Science* **9**, 597–605 (2004).
71. Coruzzi, G. & Bush, D. R. Nitrogen and carbon nutrient and metabolite signaling in plants. *Plant Physiology* **125**, 61–64 (2001).
72. Arp, W. J., Drake, B. G., Pockman, W. T., Curtis, P. S. & Whigham, D. F. Interactions between C<sub>3</sub> and C<sub>4</sub> salt marsh plant species during four years of exposure to elevated atmospheric CO<sub>2</sub>. *Vegetatio* **104**, 133–143 (1993).
73. Holm, L. G., Plucknett, D. L., Pancho, J. V. & Herberger, J. P. *The world's worst weeds: distribution and biology*. (The University Press of Hawaii, Honolulu, 1977).
74. Epstein, E., Schmid, W. E. & Rains, D. Significance and technique of short-term experiments on solute absorption by plant tissue. *Plant and Cell Physiology* **4**, 79–84 (1963).
75. Hauck, R. D. & Bremner, J. M. Use of tracers for soil and fertilizer research. *Advances in Agronomy* **28**, 219–266 (1976).
76. Christiansen, N. H., Andersen, F. Ø. & Jensen, H. S. Phosphate uptake kinetics for four species of submerged freshwater macrophytes measured by a <sup>32</sup>P phosphate radioisotope technique. *Aquatic Botany* **128**, 58–67 (2016).
77. Lu, M. *et al.* Allometry data and equations for coastal marsh plants. *Ecology* **97**, 3554–3554 (2016).

## Acknowledgements

This work was primarily supported by the Irish Research Council, Marie Curie Actions, and Bryn Mawr College. The field experiment was supported by grants from NSF LTREB (awards DEB-0950080 and DEB-1457100) and Maryland Sea Grant (award SA7528114-WW). Additional support was provided by the Smithsonian Environmental Research Center and a Bucher-Jackson Postdoctoral Fellowship to J.S.C. The authors would like to thank Patrick Megonigal, Marcel Jansen, Adam Langley, Gary Peresta, Andrew Peresta, Jim Duls, Esha Ray, and Laura Silla for their valuable contributions. The authors have no conflicts of interest with respect to this research.

## Author Contributions

T.J.M. designed the experiment and J.S.C. lead its execution. G.M.C. and J.S.C. conducted the data analysis and J.S.C. prepared the figures. G.M.C. led manuscript preparation, with J.S.C. and T.J.M. providing substantial input.

## Additional Information

**Competing Interests:** The authors declare no competing interests.

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018