

## Case Report

# Ectopia Cordis as a Lethal Neonatal Condition: A Case Report from Bahrain and a Literature Review

Shereen M. Mohamed <sup>1</sup>, Hasan M. Isa <sup>2</sup> and Amarjit K. Sandhu <sup>3</sup>

<sup>1</sup>Obstetrics and Gynaecology, Salmaniya Medical Complex, Manama, Bahrain

<sup>2</sup>Arabian Gulf University, Pediatric Department, Salmaniya Medical Complex, Manama, Bahrain

<sup>3</sup>Arabian Gulf University, Obstetrics and Gynaecology, Salmaniya Medical Complex, Manama, Bahrain

Correspondence should be addressed to Shereen M. Mohamed; [shereen\\_assiri@hotmail.com](mailto:shereen_assiri@hotmail.com)

Received 9 February 2022; Revised 19 July 2022; Accepted 27 July 2022; Published 22 August 2022

Academic Editor: Vjekoslav Krzelj

Copyright © 2022 Shereen M. Mohamed et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ectopia cordis is a rare type of malformation where the heart is not located normally. It may be partially or completely located outside the thoracic cavity and can be associated with other congenital abnormalities. It results from failure of maturation of midline mesoderm and ventral body formation during embryogenic formation. The exact etiology remains unknown. The literature review reveals the prognosis for infants with ectopia cordis is very poor. Here, we are reporting the first case of a fetus that was prenatally diagnosed with ectopia cordis that was associated with omphalocele documented in our country. Considering the poor prognosis for the fetus, conservative management during the prenatal period was chosen.

## 1. Introduction

Ectopia cordis is a rare congenital malformation with an estimated incidence of 5.5 to 7.9 per million live births [1]. Haller published the first reported case in the year 1706, and it was studied further into different types by Weese in 1818 and Todd in 1836 [2]. It accounts for only 0.1% of all congenital heart diseases. It is a malformation in which the heart is abnormally located in the extra-thoracic position and may be isolated or associated with other anomalies such as omphalocele, congenital heart disease, or integrating Cantrell syndrome. The size and location of the defect influence the prognosis. Due to the rarity and rapid postpartum mortality, limited treatment options are known. The mortality rate remains very high [1, 3, 4]. Here, we are reporting a case of ectopia cordis in a four-day-old neonate.

## 2. Case Presentation

This is a four-day-old male neonate of a 23-year-old gravida 4, para 1 plus 2 previous abortions. The mother of the patient had a healthy female full-term baby who was delivered by caesarian section for fetal distress.

During pregnancy, the mother had a routine anomaly scan at 21 weeks and 6 days of gestation which showed ectopia cordis with a small chest and lung volume. The heart was lying completely outside of the chest cavity, with a normal four-chamber view and a regular heart rate and rhythm (Video 1). The outflow tract view was difficult to obtain due to restrictions of the examination. Antenatal ultrasound scan of the anomaly is shown in Figure 1. In the abdomen, the stomach bubble was present, the diaphragm was grossly intact, the liver was difficult to visualize, and the portal vein was not clearly visualized. Normal fetal umbilical cord insertion was noted. There was some protrusion above the level of the umbilical cord insertion site that might represent the liver or small bowel. Both kidneys and urinary bladder were normal during the sonography (Figure 2). The upper and lower extremities also appeared to be normal. Due to the fetus's location, the fetal nasal bone could not be assessed as shown in the sagittal view (Figure 3). The patient was followed up prior in a different health care institute and then presented to our hospital in the late second trimester which was 26 weeks and 6 days with her all-antenatal records. Her next follow-up visit to the clinic where she





FIGURE 4: A newborn with ectopia cordis (the heart is lying outside the chest cavity with an abdominal wall defect suggestive of omphalocele).

accompanies Cantrell syndrome; (v) abdominal: where the heart passes through a defect in the diaphragm to enter the abdominal cavity [5, 6].

Ectopia cordis can be associated with chromosomal abnormalities like turner syndrome (XO), trisomy 21, and Edward syndrome (trisomy 18) [2, 7]. Intracardiac defects can also occur with ectopia cordis such as atrial septal defect (ASD), ventricular septal defect (VSD), tetralogy of Fallot (TOF), tricuspid atresia (TA), and double outlet right ventricular (DORV) [1, 8, 9].

Moreover, noncardiac malformations can occur such as pentalogy of Cantrell, omphalocele, diaphragmatic hernia, hypoplastic lung disease, skeletal dysplasia, and cleft palate [2, 3, 10].

Despite that our patient had normal chromosomal study, an associated omphalocele was noted.

Differential diagnosis of ectopia cordis includes pentalogy of Cantrell, Beckwith–Wiedemann syndrome, amniotic band syndrome, and limb-body-wall complex [3].

Ectopia cordis is associated with the presence of amniotic band syndrome. Bands may be attached to the ectopic heart or a defect in the thorax and abdominal wall. Almost all cases of amniotic band syndrome are sporadic, and the risk of recurrence is low in subsequent pregnancy [11, 12].

Prenatal diagnosis has an important role in identifying ectopia cordis and its associated abnormalities. It can be detected in the first trimester or the beginning of the second trimester [13]. Unfortunately, in our case, it was diagnosed at 21 weeks and 6 days of gestation as the patient's mother presented to the hospital during the midsecondtrimester. Nonetheless, antenatal diagnosis is possible as early as first trimester using ultrasound. Almost 90% of associated defects can be detected. However, the three-dimensional ultrasound is better for scanning to visualize fetal bones due to its greater contrast compared to two-dimensional ultrasound [2, 14–18].

Fetal cardiac magnetic resonance imaging (MRI) is an alternative imaging option to echocardiography, where it was limited by maternal or fetal factors such as maternal

obesity or abnormal fetal positions and placental calcifications [13, 16].

Chromosomal analysis is highly recommended due to its association with chromosomal abnormalities [13, 16].

Treatment of ectopia cordis requires a multidisciplinary approach consisting of a perinatologist, neonatologist, radiologist, pediatric, and pediatric cardiac surgeon. The multisurgical approach requires repair of omphalocele, sternal, diaphragmatic, pericardial defects, repair of intracardiac anomalies, and reducing the heart into the chest cavity. Since it is known that the thorax is small in ectopia cordis, the surgical approach is challenging and reducing the heart can cause kinking and compression of vessels [19]. The first trial of ectopia cordis repair was performed in 1925 by Cutler and Wilens [20], followed by the Koop trial in 1975 which achieved the first successful repair of thoracic ectopia cordis in two stages [21]. In 1995, Amato et al. achieved a successful single-stage repair of thoracic ectopia cordis [22]. Due to the rarity of ectopia cordis, there is no fixed technique used. However, the surgical approach includes closure of the chest wall and sternal defects, repair of omphalocele, placement of the heart into the thorax, and repair of intracardiac defects [14, 19, 23].

The efficacy of surgery depends on the degree of ectopia cordis and other congenital heart defects. The main aim of the surgical treatment is to provide a soft tissue cover to the heart along with reduction of the heart to the thoracic cavity and reconstruction of the chest wall [18, 24].

In the year of 2020 in Indonesia, Limanto and Soebroto did a palliative surgery on a five-day-old full-term newborn for a complete thoracic ectopia cordis with the aim of covering the exposed heart with the use of bovine pericardium material that is commonly used in reconstruction in cardiac surgery. Unfortunately, the patient expired on the third postoperative day due to heart failure [25]. In the year 2021, Yıldız et al. in Turkey reported four cases of ectopia cordis with abdominal, thoracoabdominal, partial thoracoabdominal, and middle sternum defects that were treated surgically. They concluded that the degree and level of exposure of the heart preclude initial survival, and complex repairs of intracardiac defects may be difficult in cases requiring shunts for long-term palliation [26]. In the same year in Iraq, Aboud et al. reported a case of complete thoracic ectopia cordis, but due to poor general condition, the newborn expired on 36 hours of life before any intervention [27]. Another case of thoracoabdominal ectopia cordis from Tanzania was reported by Lodhia et al. but in view of the lack of a cardiac center and waiting for relocating the patient to a cardiac institute, the newborn had a cardiac arrest on the fifth day of life and surgery of relocating the heart was not carried out [28]. Lastly, in the same year, Kebalo et al. in the Togolese Republic reported a case of midsternal defect of ectopia cordis and the newborn expired 22 hours after birth which also made them unable to perform any procedure [29]. The most recent case reported in the year 2022 from Saudi Arabia by Alshamiri et al. was a case of incomplete ectopia cordis in a 23-year-old male, who expired due to COVID-19 infection [30]. This last patient might indicate

that in patients with incomplete or partial ectopia cordis, survival may be possible.

Unfortunately, no surgical intervention could be carried out in our case because the pediatric surgeon believed that the thoracic cavity volume was very small.

The overall prognosis of ectopia cordis is poor as it is a lethal anomaly, and despite advances in neonatal cardiac surgery, this remains a surgical challenge with only a few long-term survivals.

In conclusion, the management of ectopia cordis is a challenge for any obstetrician and pediatric surgeon due to its rarity. It causes mental agony and anxiety to the parents. With early diagnosis, termination of pregnancy can be one of the options. Once it is diagnosed late, it needs a multi-disciplinary approach, and the services of a very experienced pediatric surgeon to improve the prognosis.

### Data Availability

The data supporting the findings of this study are available within the article. Raw data that supports the findings of this study are available from the corresponding author, upon reasonable request.

### Ethical Approval

Ethical approval from the secondary health care research subcommittee (SCHRC) (serial number: 62130622) was obtained.

### Consent

Written and informed consent was taken from the child's mother.

### Conflicts of Interest

The authors declare that there are no conflicts of interest.

### Authors' Contributions

SM was involved in conceptualization, data curation, writing, editing, and the final revision of the manuscript. HM was involved in conceptualization, writing, editing, and final revision. AS contributed to conceptualization and supervision.

### Supplementary Materials

There is an ultrasound video before birth and a video of the newborn right after birth showing the dynamics of the heartbeat outside the thoracic cavity uploaded as a zip file in supplementary files. (*Supplementary Materials*)

### References

- [1] J. M. Morales, S. G. Patel, J. A. Duff, R. L. Villareal, and J. W. Simpson, "Ectopia cordis and other midline defects," *The Annals of Thoracic Surgery*, vol. 70, no. 1, pp. 111–114, 2000.
- [2] A. Gabriel, J. Donnelly, A. Kuc et al., "Ectopia cordis: a rare congenital anomaly: ectopia cordis," *Clinical Anatomy*, vol. 27, no. 8, pp. 1193–1199, 2014.
- [3] P. Sadlecki, M. Krekora, G. Krasomski et al., "Prenatally evolving ectopia cordis with successful surgical treatment," *Fetal Diagnosis and Therapy*, vol. 30, no. 1, pp. 70–72, 2011.
- [4] A. M. Taksande and K. Y. Vilhekar, "A case report of ectopia cordis and omphalocele," *Indian Journal of Human Genetics*, vol. 19, no. 4, pp. 491–493, 2013.
- [5] K. A. Kim, W. R. Vincent, S. K. Muenchow, W. J. Wells, and S. E. Downey, "Successful repair of ectopia cordis using alloplastic materials," *Annals of Plastic Surgery*, vol. 38, no. 5, pp. 518–522, 1997.
- [6] A. R. Dobell, H. B. Williams, and R. W. Long, "Staged repair of ectopia cordis," *Journal of Pediatric Surgery*, vol. 17, no. 4, pp. 353–358, 1982.
- [7] Y. Çelik, O. Halloğlu, N. Basut, H. Demetgül, and A. E. Kibar, "A rare case of cardiac anomaly: prenatally diagnosed ectopia cordis," *Turk Pediatri Ars*, vol. 50, no. 2, pp. 129–131, 2015.
- [8] N. Alphonso, P. S. Venugopal, R. Deshpande, and D. Anderson, "Complete thoracic ectopia cordis," *European Journal of Cardio-Thoracic Surgery*, vol. 23, no. 3, pp. 426–428, 2003.
- [9] R. Achiron, M. Shimmel, B. Farber, and J. Glaser, "Prenatal sonographic diagnosis, and perinatal management of ectopia cordis: ectopia cordis," *Ultrasound in Obstetrics and Gynecology*, vol. 1, no. 6, pp. 431–434, 1991.
- [10] P. Russ and J. Drose, *Cardiac Malposition*, Saunders Elsevier, St. Louis, MO, USA, 2nd edition, 2010.
- [11] Y. Doi, H. Kawamata, K. Asano, and Y. Imai, "A case of amniotic band syndrome with cleft lip and palate," *Journal of Maxillofacial and Oral Surgery*, vol. 10, no. 4, pp. 354–356, 2011.
- [12] D. Das, G. Das, S. Gayen, and A. Konar, "Median facial cleft in amniotic band syndrome," *Middle East African Journal of Ophthalmology*, vol. 18, no. 2, pp. 192–194, 2011.
- [13] W. Sepulveda, A. E. Wong, L. Simonetti, E. Gomez, V. Dezerega, and J. Gutierrez, "Ectopia cordis in a first-trimester sonographic screening program for aneuploidy," *Journal of Ultrasound in Medicine*, vol. 32, no. 5, pp. 865–871, 2013.
- [14] J. Shad, K. Budhwani, and R. Biswas, "Thoracic ectopia cordis," *BMJ Case Reports*, vol. 2012, no. sep26 1, Article ID bcr1120115241, 2012.
- [15] D. Chelli, K. Dimassi, S. Jallouli-Bouzuenda et al., "Prenatal diagnosis of ectopia cordis: case report," *Tunisie Medicale*, vol. 86, no. 2, pp. 171–173, 2008.
- [16] M. A. Ergenoğlu, A. Ö. Yenieli, N. Peker, M. Kazandı, F. Akercan, and S. Sağol, "Prenatal diagnosis of Cantrell pentalogy in first trimester screening: case report and review of literature," *Journal of the Turkish-German Gynecological Association*, vol. 13, no. 2, pp. 145–148, 2012.
- [17] Y. Chang, M. J. Yang, P. H. Wang, and C. Y. Chen, "Three-dimensional HDlive image of ectopia cordis in a twin fetus at 9 gestational weeks," *Taiwanese Journal of Obstetrics & Gynecology*, vol. 54, no. 4, pp. 463–464, 2015.
- [18] D. Ramasauskaite, V. Snieckuviene, V. Zitkute, R. Vankeviciene, D. Lauzikiene, and G. Drasutiene, "A rare case report of thoracic ectopia cordis: an obstetrician's point of view in multidisciplinary approach," *Case Reports in Pediatrics*, vol. 2016, Article ID 5097059, 3 pages, 2016.
- [19] R. Carmi and J. A. Boughman, "Pentalogy of Cantrell and associated midline anomalies: a possible ventral midline

- developmental field,” *American Journal of Medical Genetics*, vol. 42, no. 1, pp. 90–95, 1992.
- [20] G. D. Cutler and G. Wilens, “Ectopia cordis: report of a case,” *American Journal of Diseases of Children*, vol. 30, no. 1, p. 76, 1925.
- [21] A. Saxena, “Pectus excavatum, pectus carinatum and other forms of thoracic deformities,” *Journal of Indian Association of Pediatric Surgeons*, vol. 10, no. 3, p. 147, 2005.
- [22] J. J. Amato, J. Zelen, and N. G. Talwalkar, “Single-stage repair of thoracic ectopia cordis,” *The Annals of Thoracic Surgery*, vol. 59, no. 2, pp. 518–520, 1995.
- [23] M. R. Harrison, R. A. Filly, P. Stanger, and A. A. de Lorimier, “Prenatal diagnosis and management of omphalocele and ectopia cordis,” *Journal of Pediatric Surgery*, vol. 17, no. 1, pp. 64–66, 1982.
- [24] S. G. Serrano Gómez, M. T. Bermlúez Sosa, E. Palma Hernández et al., “Ectopia cordis: multidisciplinary approach with successful result (in Spanish),” *Ginecología y Obstetricia de México*, vol. 74, pp. 439–443, 2006.
- [25] D. H. Limanto and H. Soebroto, “Thoracic ectopia cordis: a case report,” *Jurnal Kedokteran dan Kesehatan Indonesia*, vol. 11, no. 1, pp. 99–104, 2020.
- [26] K. Yıldız, M. Kır, ŞB. Uğurlu, H. Z. Genç, and N. Ünal, “Four ectopia cordis cases surgically managed with different strategies,” *The Turkish Journal of Thoracic and Cardiovascular Surgery*, vol. 29, no. 1, pp. 114–118, 2021.
- [27] M. J. Aboud, M. J. Aboud, N. Abudi, H. Joudi, and Z. Joudi, “Complete thoracic ectopia cordis, dilemma of the outcome: a case report,” *Surgical Case Reports*, pp. 1–4, 2021.
- [28] J. Lodhia, H. Chipongo, K. Chilonga, D. Mchaile, R. Mbwasii, and R. Philemon, “Ectopia cordis: a case report of pre-surgical care in resource-limited setting,” *International Journal of Surgery Case Reports*, vol. 83, Article ID 105965, 2021.
- [29] S. P. Kebalo, K. O. Sibabi-Akpo, D. V. Teko, I. Nengha, J. S. Gbelesso, and K. J.-P. Gnassingbe, “Ectopia cordis: about a case in Togo,” *Cirugía Cardiovascular*, vol. 29, no. 1, pp. 49–51, 2022.
- [30] K. M. Alshamiri, A. Z. Albriek, T. W. Farrag, and M. Q. Alshamiri, “Ectopia cordis in an adult patient with COVID-19: a case report and literature review,” *Clinical Case Reports*, vol. 10, no. 2, Article ID e05389, 2022.