

Draft Genome Sequence of *Frankia* sp. Strain BCU110501, a Nitrogen-Fixing Actinobacterium Isolated from Nodules of *Discaria trinevis*

Luis G. Wall,^a Nicholas Beauchemin,^b Michael N. Cantor,^c Eugenia Chaia,^d Amy Chen,^e J. Chris Detter,^c Teal Furnholm,^b Faten Ghodhbane-Gtari,^{b,f} Lynne Goodwin,^c Maher Gtari,^{b,f} Cliff Han,^c James Han,^e Marcel Huntemann,^e Susan Xinyu Hua,^e Natalia Ivanova,^e Nikos Kyrpides,^e Victor Markowitz,^e Kostas Mavrommatis,^e Natalia Mikhailova,^e Henrik P. Nordberg,^c Imen Nouioui,^f Galina Ovchinnikova,^e Ioanna Pagani,^e Amrita Pati,^e Arnab Sen,^g Saubashya Sur,^g Ernest Szeto,^e Subarna Thakur,^g Chia-Lin Wei,^c Tanja Woyke,^e Louis S. Tisa^b

University of Quilmes, Bernal, Argentina^a; University of New Hampshire, Durham, New Hampshire, USA^b; Los Alamos National Laboratory, Los Alamos, New Mexico, USA^c; Centro Regional Universitario, Bariloche, Argentina^d; Department of Energy (DOE) Joint Genome Institute, Walnut Creek, California, USA^e; University of Tunis-El Manar, Tunis, Tunisia^f; University of North Bengal, Siliguri, India^g

***Frankia* forms a nitrogen-fixing symbiosis with actinorhizal plants. We report a draft genome sequence for *Frankia* sp. strain BCU110501, a nitrogen-fixing actinobacterium isolated from nodules of *Discaria trinevis* grown in the Patagonia region of Argentina.**

Received 7 June 2013 Accepted 7 June 2013 Published 11 July 2013

Citation Wall LG, Beauchemin N, Cantor MN, Chaia E, Chen A, Detter JC, Furnholm T, Ghodhbane-Gtari F, Goodwin L, Gtari M, Han C, Han J, Huntemann M, Hua SX, Ivanova N, Kyrpides N, Markowitz V, Mavrommatis K, Mikhailova N, Nordberg HP, Nouioui I, Ovchinnikova G, Pagani I, Pati A, Sen A, Sur S, Szeto E, Thakur S, Wei C-L, Woyke T, Tisa LS. 2013. Draft genome sequence of *Frankia* sp. strain BCU110501, a nitrogen-fixing actinobacterium isolated from nodules of *Discaria trinevis*. *Genome Announc.* 1(4):e00503-13. doi: 10.1128/genomeA.00503-13.

Copyright © 2013 Wall et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Louis S. Tisa, louis.tisa@unh.edu.

The genus *Frankia* forms a symbiotic nitrogen-fixing association with woody trees and shrubs, resulting in a root nodule formation (1–3). *Frankia* exists either in a free-living state in the soil or in symbiosis with actinorhizal plants. These actinorhizal plants comprise over 200 different plant species belonging to 8 distinct plant families. Because of symbiosis, these globally distributed actinorhizal plants are able to colonize harsh environmental terrains under diverse ecological conditions. Phylogenetic studies have shown that *Frankia* has four major lineages or clusters that can also be defined by physiology, host specificity, and mode of infection of the host plant (4–7). Genomes for representatives from each of these clusters have been sequenced (8–13) and have provided essential baseline information for genomic approaches toward understanding these novel bacteria.

Frankia strains from cluster III have the greatest metabolic versatility, are known to associate with five plant families (*Betulaceae*, *Myricaceae*, *Elaeagnaceae*, *Rhamnaceae*, and *Casuarinaceae*), and are considered broad-host-range symbionts. Many of these strains have adapted to harsh environmental conditions. *Frankia* sp. strain BCU110501 (also called Dtl1 and T1) was isolated from a field nodule of *Discaria trinevis* growing wild in a semiarid steppe (40°41'S, 71°10'W) of the Argentinian Reserve and National Park “Parque Nacional Nahuel Huapi,” located in the northwest of Patagonia (14). The physiology and infectivity of this strain have been well studied (15–20), and it uses an intercellular route of plant infection (18). *Frankia* sp. strain BCU110501 was chosen for sequencing as another cluster III representative with broad-host-range properties and an intercellular infection route. Strain BCU110501 was sequenced to provide information about the potential ecological roles of the *Frankia* strains and their interactions with actinorhizal plants.

The draft genome of *Frankia* sp. strain BCU110501 was generated at the Department of Energy (DOE) Joint Genome Institute (JGI) using Illumina technology (21). An Illumina standard shotgun library was constructed and sequenced using the Illumina HiSeq 2000 platform, which generated 12,553,550 reads totaling 1,883.0 Mbp. All techniques for DNA isolation, library construction, and sequencing were performed according to JGI standards and protocols (<http://www.jgi.doe.gov>). The Illumina sequence data were assembled using Velvet (version 1.1.04) (22) and Allpaths-LG (version r41043) (23). The final draft assembly contained 207 contigs in 207 scaffolds. The total size of the genome is 7.9 Mbp and the final assembly is based on 950.8 Mbp of Illumina data, which provides an average 120.4× coverage of the genome.

The draft genome of *Frankia* BCU110501 was resolved to 207 scaffolds consisting of 7,891,711 bp with a G+C content of 72.39%, 6,742 candidate protein-encoding genes, 47 tRNA genes, and 2 rRNA regions.

Nucleotide sequence accession numbers. The *Frankia* sp. strain BCU110501 genome sequence has been deposited at DDBJ/EMBL/GenBank under the accession number [ARDT00000000](https://www.ncbi.nlm.nih.gov/nuclink/ARDT00000000). The version described in this paper is the first version, [ARDT01000000](https://www.ncbi.nlm.nih.gov/nuclink/ARDT01000000).

ACKNOWLEDGMENTS

The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under contract no. DE-AC02-05CH11231. This project (L.S.T.) was supported in part by Agriculture and Food Research Initiative grant 2010-65108-20581 from the USDA National Institute of Food and Agriculture, Hatch grant NH530, and the College of Life Sciences and Agriculture at the University of New Hampshire, Durham, NH. M.G. and

F.G.-G. were supported in part by a Visiting Scientist and Postdoctoral Scientist Program administered by the NH AES at the University of New Hampshire.

REFERENCES

- Benson DR, Silvester WB. 1993. Biology of *Frankia* strains, actinomycete symbionts of actinorhizal plants. *Microbiol. Rev.* 57:293–319.
- Schwencke J, Caru M. 2001. Advances in actinorhizal symbiosis: host plant-*Frankia* interactions, biology, and applications in arid land reclamation. A review. *Arid Land Res. Manag.* 15:285–327.
- Chaia EE, Wall LG, Huss-Danell K. 2010. Life in soil by actinorhizal root nodule endophyte *Frankia*. A review. *Symbiosis* 51:201–226.
- Benson DR, Clawson ML. 2000. Evolution of the actinorhizal plant symbioses, p 207–224. In Triplett EW (ed), *Prokaryotic nitrogen fixation: a model system for analysis of biological process*. Horizon Scientific Press, Wymondham, United Kingdom.
- Normand P, Orso S, Cournoyer B, Jeannin P, Chapelon C, Dawson JO, Evtushenko L, Mirsra AK. 1996. Molecular phylogeny of the genus *Frankia* and related genera and emendation of the family *Frankiaceae*. *Int. J. Syst. Bacteriol.* 46:1–9.
- Nouioui I, Ghodhbane-Gtari F, Beauchemin NJ, Tisa LS, Gtari M. 2011. Phylogeny of members of the *Frankia* genus based on *gyrB*, *nifH* and *glnII* sequences. *Antonie Van Leeuwenhoek* 100:579–587.
- Ghodhbane-Gtari F, Nouioui I, Chair M, Boudabous A, Gtari M. 2010. 16S-23S rRNA intergenic spacer region variability in the genus *Frankia*. *Microb. Ecol.* 60:487–495.
- Normand P, Lapiere P, Tisa LS, Gogarten JP, Alloisio N, Bagnarol E, Bassi CA, Berry AM, Bickhart DM, Choisine N, Couloux A, Cournoyer B, Cruveiller S, Daubin V, Demange N, Francino MP, Goltsman E, Huang Y, Kopp OR, Labarre L, Lapidus A, Lavire C, Marechal J, Martinez M, Mastrorunzio JE, Mullin BC, Niemann J, Pujic P, Rawnsley T, Rouy Z, Schenowitz C, Sellstedt A, Tavares F, Tomkins JP, Vallenet D, Valverde C, Wall LG, Wang Y, Medigue C, Benson DR. 2007. Genome characteristics of facultatively symbiotic *Frankia* sp. strains reflect host range and host plant biogeography. *Genome Res.* 17:7–15.
- Normand P, Queirox C, Tisa LS, Benson DR, Rouy Z, Cruveiller S, Medigue C. 2007. Exploring the genomes of *Frankia*. *Physiol. Plant.* 130:331–343.
- Persson T, Benson DR, Normand P, Vanden Heuvel B, Pujic P, Chertkov O, Teshima H, Bruce DC, Detter C, Tapia R, Han S, Han J, Woyke, Pitlock S, Pennacchio L, Nolan M, Ivanova N, Pati A, Land ML, Pawlowski K, Berry AM. 2011. Genome sequence of “*Candidatus Frankia datiscacae*” Dg1, the uncultured microsymbiont from nitrogen-fixing root nodules of the dicot *Datisca glomerata*. *J. Bacteriol.* 193:7017–7018.
- Ghodhbane-Gtari F, Beauchemin N, Bruce D, Chain P, Chen A, Walston Davenport K, Deshpande S, Detter C, Furnholm T, Goodwin L, Gtari M, Han C, Han J, Huntemann M, Ivanova N, Kyrpides N, Land ML, Markowitz V, Mavrommatis K, Nolan M, Nouioui I, Pagani I, Pati A, Pitluck S, Santos CL, Sen A, Sur S, Szeto E, Tavares F, Teshima H, Thakur S, Wall LG, Woyke T, Tisa LS. 2013. Draft genome sequence of *Frankia* sp. strain CN3, an atypical, noninfective (Nod⁻) ineffective (Fix⁻) isolate from *Coriaria nepalensis*. *Genome Announc.* 1(2):e00085-13. doi: [10.1128/genomeA.00085-13](https://doi.org/10.1128/genomeA.00085-13).
- Sen A, Beauchemin N, Bruce D, Chain P, Chen A, Walston Davenport K, Deshpande S, Detter C, Furnholm T, Ghodhbane-Gtari F, Goodwin L, Gtari M, Han C, Han J, Huntemann M, Ivanova N, Kyrpides N, Land ML, Markowitz V, Mavrommatis K, Nolan M, Nouioui I, Pagani I, Pati A, Pitluck S, Santos CL, Sur S, Szeto E, Tavares F, Teshima H, Thakur S, Wall L, Woyke T, Wishart J, Tisa LS. 2013. Draft genome sequence of *Frankia* sp. strain QA3, a nitrogen-fixing actinobacterium isolated from the root nodule of *Alnus nitida*. *Genome Announc.* 1(2):e00103-13. doi: [10.1128/genomeA.00103-13](https://doi.org/10.1128/genomeA.00103-13).
- Nouioui I, Beauchemin N, Cantor MN, Chen A, Detter JC, Furnholm T, Ghodhbane-Gtari F, Goodwin L, Gtari M, Han C, Han J, Huntemann M, Hua SX, Ivanova N, Kyrpides N, Markowitz V, Mavrommatis K, Mikhailova N, Nordberg HP, Ovchinnikova G, Pagani I, Pati A, Sen A, Sur S, Szeto E, Thakur S, Wall L, Wei C-L, Woyke T, Tisa LS. 2013. Draft genome sequence of *Frankia* sp. strain BMG5.12, a nitrogen-fixing actinobacterium isolated from Tunisian soils. *Genome Announc.* 1(4):e00468-13. doi: [10.1128/genomeA.00468-13](https://doi.org/10.1128/genomeA.00468-13).
- Chaia E. 1998. Isolation of an effective strain of *Frankia* from nodules of *Discaria trinervis* (*Rhamnaceae*). *Plant Soil* 205:99–102.
- Chaia EE, Solans M, Vobis G, Wall LG. 2007. Infectivity variation of *Discaria trinervis*-nodulating *Frankia* in Patagonian soil according to season and storage conditions. *Physiol. Plant.* 130:357–360.
- Gabbarni LA, Wall LG. 2008. Analysis of nodulation kinetics in *Frankia-Discaria trinervis* symbiosis reveals different factors involved in the nodulation process. *Physiol. Plant.* 133:776–778.
- Valverde C, Ferrari A, Wall LG. 2002. Phosphorus and the regulation of nodulation in the actinorhizal symbiosis, between *Discaria trinervis* (*Rhamnaceae*) and *Frankia* BCU11050. *New Phytol.* 153:43–52.
- Valverde C, Wall LG. 1999. Time course of nodule development in *Discaria trinervis* (*Rhamnaceae*)-*Frankia* symbiosis. *New Phytol.* 141:345–354.
- Valverde C, Wall LG. 1999. Regulation of nodulation in *Discaria trinervis* (*Rhamnaceae*)-*Frankia* symbiosis. *Can. J. Bot.* 77:1302–1310.
- Valverde C, Ferrari A, Wall LG. 2009. Effect of calcium in the nitrogen-fixing symbiosis between actinorhizal *Discaria trinervis* (*Rhamnaceae*) and *Frankia*. *Symbiosis* 49:151–155.
- Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438.
- Zerbino D, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.
- Gnerre S, MacCallum I. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc. Natl. Acad. Sci. U. S. A.* 108:1513–1518.