



OPEN

Physical activity and gender buffer the association of retirement with functional impairment in Ghana

Razak M. Gyasi^{1,2}✉, Padmore Adusei Amoah³, Seth Agyemang⁴, Lawrence Pokua Siaw⁴, Foster Frempong⁴, Ritu Rani⁵ & David R. Phillips⁶

Females on average live longer but with higher rates of functional impairment and lower physical and economic activities than men. However, research linking retirement to functional impairment and the modifying role of gender and physical activity (PA) is limited especially in low- and middle-income countries. This paper examines the association between retirement and functional impairment in Ghana and evaluates the effect modification of the association by gender and PA. The sample included 1201 adults aged ≥ 50 years from a population-based study. Functional impairment was assessed with the activities of daily living scale. Ordinary least squares regression models adjusted for confounding variables and estimated gender-wise and PA heterogeneity effect of retirement on functional impairment. Regressions showed that retirement predicted an increase in functional impairment score in the full sample ($\beta = .76, p < .001$) and in men ($\beta = 1.96, p < .001$), but not in women. Interestingly, retirement significantly increased functional impairment in ≥ 65 age cohort (full sample: $\beta = .71, p < .005$; men: $\beta = 1.86, p < .001$) although not in women. However, the effect was significantly moderated by PA such that retirement \times PA predicted a decrease in functional impairment in the full sample ($\beta = -.81, p < .005$) and the ≥ 65 age group ($\beta = -.43, p < .005$). Functional impairment risk of retirement is gender-specific, but PA buffers the relationship. Retirement is generally commonplace, but these findings imply that promoting PA may hold promise for addressing functional impairment in old age. Attending to the physical health needs of men during retirement should be a social policy priority.

Abbreviations

ADL	Activities of daily living
CHRPE	Committee on Human Research Publication and Ethics
CI	Confidence intervals
GSS	Ghana Statistical Service
HRS	Health and Retirement Study
IPAQ-SF	International Physical Activity Questionnaire short form
LMICs	Low- and middle-income countries
MET	Metabolic equivalent
OLS	Ordinary least squares
PA	Physical activity
SRH	Self-rated health
SSA	Sub-Saharan Africa

¹Aging and Development Unit, African Population and Health Research Center, Manga Close, Off-Kirawa Road, P. O. Box 10787-00100, Nairobi, Kenya. ²National Centre for Naturopathic Medicine (NCNM), Faculty of Health, Southern Cross University, Lismore, NSW, Australia. ³School of Graduate Studies, Department of Applied Psychology, Asia Pacific Institute of Aging Studies, Lingnan University, Tuen Mun, Hong Kong. ⁴Department of Geography and Rural Development, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana. ⁵International Institute for Population Sciences, Mumbai, India. ⁶Department of Sociology and Social Policy, Lingnan University, Tuen Mun, Hong Kong. ✉email: RGYasi.Research@gmail.com

UN United Nations
VIF Variance Inflation Factor

Functional impairment, defined as the inability of persons to carry out certain functions in their daily lives due to illnesses and diseases¹ and the tendency for reduced physical activities on retirement, in combination pose significant health and contemporary social policy challenge. In sub-Saharan Africa (SSA), an estimated 43% of persons 60 years or over report disabilities^{2,3}, and difficulty undertaking ADL, mainly due to contextual risk factors^{4,5}. Functional impairment is a leading cause of disability, long-term severe care needs, and medical costs⁶. However, the right for older people to live and participate socially is emphasized by the United Nations Convention on the Right of Persons with Disabilities⁷ and for successful aging⁸. In this context, the concern about apparently rising functional impairment in rapidly aging adults in SSA stimulates a renewed interest in identifying the risk factors associated with functional impairment.

Research in Western and Asian societies has suggested effects of retirement on subsequent health outcomes, including cognitive decline^{9,10}, depression¹¹, psychological state¹², and functional impairment. Findings are not clear cut and a systematic review found inconsistencies in the association of retirement with functional impairment¹³. Dave et al.¹⁴ relate retirement to a 5–16% increase in difficulty in ADL. Some studies suggest that the transition into retirement alleviates physical limitations and contributes to better health^{15,16} while others found no associations between the variables^{17,18}. Retirement may be linked to functional impairment via several mechanisms. For example, retirement may invoke changes in lifestyle behavior, including physical activity (PA) and social participation¹⁹ which can potentially stimulate body-mind interactions for healthy living. Further, the anticipation of disengagement and uncertainty in retirement-induced economic hardships can lead to stress responses, contributing to depression and anxiety²⁰ and subsequently amplifying functional limitations²¹.

Several authors have identified heterogeneity in the aging population in terms of physical health outcomes and survival and have noted marked gender differences^{22,23}. The prevalence of functional impairment, including ADL and mobility limitations, is generally higher in women, partly due to women's greater socioeconomic disadvantage (especially in low- and middle-income countries), their chances of surviving longer, and other biological disparities^{6,24}. With aging populations generally including more females than males and with often established gender differences in the nature of work and retirement, it is essential to understand whether any functional impairment effect of retirement varies between gender subgroups. Studies have indicated that females have an increased risk of retiring earlier than males²⁵. Others such as Radl²⁶, on the other hand, note that in countries with low female labor participation rates, females retire later than males. Yet other studies suggest no apparent differences in retirement between the gender subgroups as males and females have markedly different socioeconomic positions and retirement patterns^{27–29}. It has also been shown that although physical activity (PA) is a salient psychosocial resource for health through coping with stressful life events³⁰, PA declines after retirement⁹. As a social determinant of health, PA may relate to life course transitions and improve functional well-being³¹. Cross-sectional and prospective studies report various levels of PA associated with better functional abilities^{32–35}. For example, longitudinal data from the Frailty, Dynapenia, and Sarcopenia study among Mexican adults found that moderate to vigorous PA intensity significantly reduced difficulties in ADL and physical performance³⁶. However, we are unaware of any study exploring the effect modification of the association between retirement and functional impairment by PA, particularly in low- and middle-income countries. In SSA in particular, whether the risk of functional impairment after retirement differs across gender subgroups remains unclear.

This study employs exploratory survey data to examine the gender-specific associations between retirement and functional impairment in Ghana and evaluates the potential modifying effect of PA in these associations. Despite the rising and gendered patterns of life expectancy, the average retirement age is declining locally due to increasing trends in voluntary retirement³⁷. For example, the Ghana Statistical Service³⁸ has shown a reduction in the average retirement age from 60 years to about 58 years for males and 55 years for females. This is related to the factors that motivate premature retirement, including the perception of worsening health status, rising burden of disease, increasing caregiving burden, and other socioeconomic events^{11,39}. Our study is important to improve understanding of gender-based retirement and functional status with relevant implications for public health policy. We hypothesized: (1) that the risks of functional impairment increase after retirement; (2) that the impact of retirement on functional impairment would differ by gender; (3) that PA would modify the association between retirement and functional impairment.

Results

After exclusion criteria were applied, 1201 respondents were included in the final analytical cohort (Fig. 1). About 63% were women, and the mean age was 66.2 years (SD = 11.9). Most participants were not married (65%), lived in urban areas (55%), and had up to primary education (86%). Income levels were low with an average of GH¢308 (SD = 338.89) (US\$64). Almost 50% of the sample perceived their overall health as poor/fair, while the mean psychological distress, functional impairment, and physical activity scores were 15.91 (SD = 4.66), 13.70 (SD = 5.09), and 6.10 (SD = 2.68) respectively. About 16% of the sample were retired and the mean social connectedness score was 6.09 (SD = 2.6). The characteristics of the sample are summarized in Table 1.

Among those in the retirement cohort, more women were retired (53.2% [95CI = 45.8–60.5]) compared with men (46.8%, 95CI = 39.5–54.2) ($p < 0.005$). The retirees were more likely to be functionally impaired (44.7%, 95%CI 37.4–52.1) compared with those who were not retired (27.5%, 95%CI 24.7–30.3) ($p < 0.001$). Women were more likely to be functionally impaired (34.4%, 95%CI 31.0–37.9) than men (22.9%, 95%CI 19.1–27.1) ($p < 0.001$) (Table 2). Correlation analysis (Table 3) revealed that retirement ($r = 0.136$, $p < 0.001$), income ($r = -0.143$, $p < 0.001$), marital status ($r = 0.126$, $p < 0.001$) and psychological distress ($r = 0.182$, $p < 0.001$) were significantly associated with functional impairment following a Bonferonni correction for multiple correlations.




Figure 1. Flow chart of the selection of study participants.

Variable	N= 1201	(%)	Mean	(±SD)	Range
Retired	188	(15.7)			
PA score			6.10	(± 2.68)	0–21
Mean age (in years)			66.15	(± 11.85)	50–111
Women	759	(63.3)			
Rural residence	540	(45.0)			
Marital status: Married	521	(44.3)			
Marital status: Widow	473	(40.3)			
Marital status: Divorced/separated	181	(15.4)			
Primary/no education	1034	(86.2)			
Secondary education	104	(8.7)			
Higher education	62	(5.2)			
Income level (in Ghana Cedis)			308.18	(± 338.89)	100–4000
Social connectedness			6.09	(± 2.679)	0–20
SRH: Excellent/Very Good	239	(19.9)			
SRH: Good	369	(30.8)			
SRH: Fair	348	(29.0)			
SRH: Poor	244	(20.3)			
Psychological distress score			15.91	(± 4.66)	0–40
FI score			13.70	(± 5.09)	0–18

Table 1. Characteristics of the sample. FI—functional impairment; AP—physical activity; SRH—self-rated health; ^aExchange rate: \$1 ~ ₵4.8 as of the time of data collection.

Moreover, PA, age, gender, education, social connectedness, and SRH significantly related to both functional impairment and retirement.

Table 4 shows the results of OLS regressions. Adjusting for potential confounders, retirement was significantly associated with increasing functional impairment in the total sample ($\beta = 0.76$, $p < 0.001$) and in men ($\beta = 1.96$, $p < 0.001$). In contrast, retirement was not associated with functional impairment in women ($p = 0.075$). Decreasing PA, worsening SRH, psychological distress, socially connected, increasing age, and being widowed were

	Gender				Rao Scott χ^2 <i>p</i> -value	Retirement status				Rao Scott χ^2 <i>p</i> -value
	Men		Women			Not retired		Retired		
	%	(95%CI)	%	(95%CI)		%	(95%CI)	%	(95%CI)	
Number	441		759			1012		188		
Total	36.8	(34.0–39.5)	63.2	(60.5–66.0)	–	84.3	(82.2–86.3)	15.7	(13.7–17.8)	–
Age (in years)										
50–59	34.9	(30.5–39.6)	35.8	(32.4–39.4)	0.478	40.7	(37.7–43.8)	7.4	(4.1–12.2)	<0.001
60–69	30.6	(26.3–35.1)	27.4	(24.3–30.7)		27.1	(24.4–29.9)	36.7	(29.8–44.0)	
70+	34.5	(30.0–39.1)	36.8	(33.3–40.3)		32.2	(29.3–35.2)	55.9	(48.4–63.1)	
Retirement status										
Not retired	86.8	(84.2–89.2)	80.0	(76.0–83.7)	0.002	–	–	–	–	
Retired	20.0	(10.8–15.8)	13.2	(16.3–24.0)						
Gender										
Female	–	–	–	–	–	65.1	(62.1–68.1)	53.2	(45.8–60.5)	0.002
Male	–	–	–	–		34.9	(31.9–37.9)	46.8	(39.5–54.2)	
PA										
Inactive	38.8	(34.2–43.5)	55.7	(52.1–59.3)	<0.001	47.5	(44.4–50.7)	60.1	(52.7–67.2)	0.002
Active	61.2	(56.5–65.8)	44.3	(40.7–47.9)		52.5	(49.3–55.6)	33.9	(32.8–47.3)	
Residential status										
Rural	47.8	(43.1–52.6)	43.3	(39.8–47.0)	0.131	45.4	(42.3–48.5)	43.1	(35.9–50.5)	0.565
Urban	52.2	(47.4–56.9)	56.7	(53.0–60.2)		54.6	(51.5–57.7)	56.9	(49.5–64.1)	
Marital status										
Married	71.1	(66.6–75.4)	29.1	(25.9–32.5)	<0.001	44.1	(41.0–47.3)	45.7	(38.3–53.1)	0.075
Widow	12.7	(9.7–16.2)	55.9	(52.3–59.5)		41.4	(38.3–44.5)	34.2	(27.4–41.6)	
Divorced/separated	16.2	(12.8–20.0)	15.0	(12.5–17.7)		14.5	(12.4–16.9)	20.1	(14.6–26.6)	
Level of education										
Primary/Never	76.4	(72.2–80.3)	91.8	(89.7–93.7)	<0.001	88.9	(86.8–90.8)	71.3	(64.2–77.6)	<0.001
Secondary	13.6	(10.5–17.2)	5.8	(4.2–7.7)		7.9	(6.3–9.7)	12.8	(8.4–18.4)	
Higher	10.0	(7.3–13.2)	2.4	(1.4–3.7)		3.2	(2.2–4.4)	1.6	(1.1–2.2)	
Income level (In Ghana Cedis)										
Low	64.6	(59.7–69.3)	81.0	(77.7–83.9)	<0.001	75.1	(72.1–77.9)	72.7	(64.8–79.6)	0.533
High	35.4	(30.7–40.3)	19.0	(16.1–22.3)		24.9	(22.1–27.9)	27.3	(20.4–35.2)	
Social connectedness										
Low	39.9	(35.3–44.6)	38.9	(35.4–42.4)	0.721	40.0	(37.0–43.1)	35.1	(29.3–42.4)	0.205
High	60.1	(55.4–64.7)	61.1	(57.6–64.6)		60.0	(56.9–63.0)	64.9	(57.6–71.7)	
Self-rated health (SRH)										
Excellent/Very Good	28.6	(24.4–33.0)	14.9	(12.4–17.6)	<0.001	20.9	(18.5–23.6)	14.4	(9.7–20.2)	0.027
Good	30.6	(26.3–35.1)	30.8	(27.6–34.3)		31.2	(28.4–34.2)	28.2	(21.9–35.2)	
Fair	26.1	(22.0–30.4)	30.7	(27.4–34.1)		27.5	(24.7–30.3)	37.2	(30.3–44.6)	
Poor	14.7	(11.6–18.4)	23.6	(20.6–26.8)		20.4	(17.9–23.0)	20.2	(14.7–26.7)	
Psychological Distress										
Low	62.4	(57.7–66.9)	50.2	(46.6–53.8)	<0.001	53.7	(50.5–56.8)	60.1	(52.7–67.2)	0.103
High	37.6	(33.1–42.3)	49.8	(46.2–53.4)		46.3	(43.2–49.5)	39.9	(32.8–47.3)	
FI										
Not impaired	77.1	(72.9–80.9)	65.6	(62.1–69.0)	<0.001	72.5	(69.7–75.3)	55.3	(47.9–62.6)	<0.001
Impaired	22.9	(19.1–27.1)	34.4	(31.0–37.9)		27.5	(24.7–30.3)	44.7	(37.4–52.1)	

Table 2. Bivariate distribution of demographic and health-related characteristics by gender and retirement status ($N = 1201$). FI—functional impairment; AP—physical activity; SRH—self-rated health.

associated with an increased functional impairment. The second aim of this study tested the effect modification of the retirement-functional impairment association by PA. The main model was extended by adding the interaction term, retirement \times PA. We found that PA moderated and dampened the relationship of retirement with functional impairment: retirement \times PA predicted a decrease in functional impairment ($\beta = -0.81$, $p < 0.005$) (Fig. 2). In sensitivity analysis, we extracted age group ≥ 65 and estimated the effect of retirement on functional impairment (Table 5). Results showed slightly reduced estimates in terms of effect size: retirement increased functional impairment in the total sample ($\beta = 0.71$, $p < 0.005$), and in men only ($\beta = 1.86$, $p < 0.001$), but the effect was significantly moderated by PA ($\beta = -0.43$, $p < 0.005$) (Fig. 3). In a supplementary analysis, employment status

Variable	FI	Retired
FI	1	.136***
Retired	.136***	1
PA	-.399***	-.091***
Age (in years)	.338***	.233***
Women	-.121***	.090***
Rural residence	.007	.017
Marital status: Not married	.126***	.020
Higher education	-.060**	.218***
Income level (in Ghana Cedis)	-.143***	-.017
Social connectedness	.101***	.115**
Suboptimal SRH	.390***	.057**
Psychological distress	.182***	-.047

Table 3. Correlations between the independent and dependent variables of interest. Pearson product-moment correlations were used to calculate the association between continuous variables, point-biserial correlations were used to assess the relationship between continuous and dichotomous variables, and phi correlations were used to assess the relationship between dichotomous variables. FI—functional impairment; AP—physical activity; SRH—self-rated health.

Variables	(1) Pooled Sample		(2) Male		(3) Female		(4) Interaction: Retirement × PA	
	β	(SE)	β	(SE)	β	(SE)	β	(SE)
Retired (Not retired) ^a	0.764	(0.254)***	1.955	(0.474)***	0.233	(0.328)	0.956	(0.251)***
PA score	-0.190	(0.023)***	-0.150	(0.045)***	-0.216	(0.028)***	-1.264	(0.229)***
Age (in years)	0.046	(0.009)***	0.076	(0.018)***	0.039	(0.010)***	0.045	(0.008)***
Gender (Male) ^a	1		1		1		1	
Female	0.281	(0.217)	-	-	-	-	0.182	(0.208)
Residential status (Rural) ^a	1		1		1		1	
Urban	0.065	(0.189)	-0.267	(0.384)	0.077	(0.227)	0.006	(0.184)
Marital status (Married) ^a			1		1		1	
Widow	0.453	(0.221)*	0.980	(0.448)*	0.271	(0.264)	0.502	(0.215)*
Divorced/separated	-0.492	(0.299)	-0.194	(0.535)*	-0.714	(0.388)	-0.619	(0.283)*
Education (Primary or none) ^a	1		1		1		1	
Secondary	-0.644	(0.387)	-1.426	(0.698)*	-0.459	(0.515)	-0.780	(0.386)*
Tertiary	-0.020	(0.415)	-0.565	(0.576)	-0.011	(0.711)	0.009	(0.396)
Income level (In Ghana Cedis)	-0.031	(0.242)	0.864	(0.470)	-0.181	(0.303)	-0.019	(0.237)
Social connectedness score	0.135	(0.067)*	-0.160	(0.132)	0.222	(0.082)**	0.250	(0.188)
SRH (Excellent/Very Good) ^a	1		1		1		1	
Good	0.036	(0.315)	-0.075	(0.573)	0.093	(0.389)	0.126	(0.309)
Fair	0.808	(0.298)**	1.342	(0.558)*	0.663	(0.370)	0.979	(0.295)**
Poor	2.004	(0.316)***	2.146	(0.567)***	1.934	(0.392)***	2.208	(0.314)***
Psychological Distress Score (10–50)	0.463	(0.187)*	1.094	(0.407)**	0.410	(0.224)	0.437	(0.180)*
Interaction term: Retirement × PA							-0.805	(0.457)**
Adjusted Pseudo R ²	0.477		0.515		0.468		0.479	

Table 4. Predicting FI Score with Retirement and PA Score: OLS Regression Models. β —Coefficients; SE—robust standard errors are presented in parentheses; SRH—Self-rated Health; ^aReference Group. *** $p < 0.001$; ** $p < 0.005$; * $p < 0.05$.

was stratified into employed, unemployed, and retired. Retirement was positively associated with functional impairment with larger effect sizes in the total sample and men.

Discussion

Main findings. This study from Ghana, an SSA country context, found a retirement rate of 15.7%, and 53.2% of the retirees were women. The prevalence of functional impairment was more common in women and retirees than in their counterparts. Independent of potential confounders, retirement was associated with increased




Figure 2. Physical activity (PA, the moderator) dampens the association of retirement with functional impairment (FI) in the overall sample.

Variables	(1) Pooled Sample		(2) Male		(3) Female		(4) Interaction: Retirement × PA	
	β	(SE)	β	(SE)	β	(SE)	β	(SE)
Retired (Not retired) ^a	0.707	(0.268)**	1.864	(0.478)***	0.019	(0.358)	0.773	(0.273)**
PA score	-1.114	(0.208)***	-1.024	(0.390)**	-1.268	(0.266)***	-1.004	(0.240)***
Age (in years)	0.044	(0.011)***	0.049	(0.022)*	0.042	(0.014)**	0.044	(0.011)***
Gender (Male) ^a	1		1		1		1	
Female	-0.022	(0.241)	-	-	-	-	-0.029	(0.242)
Residential status (Rural) ^a	1		1		1		1	
Urban	-0.058	(0.204)	-0.380	(0.407)	-0.123	(0.259)	-0.050	(0.205)
Marital status (Married) ^a			1		1		1	
Widow	0.769	(0.248)**	1.558	(0.517)**	0.451	(0.307)	0.451	(0.307)**
Divorced/separated	-0.610	(0.330)	0.047	(0.560)	-1.094	(0.462)*	-1.094	(0.462)
Education (Primary or none) ^a	1		1		1		1	
Secondary	-0.683	(0.438)	-1.706	(0.768)*	-0.239	(0.637)	-0.718	(0.441)
Tertiary	0.353	(0.469)	-0.569	(0.611)	0.989	(0.887)	0.312	(0.463)
Income level (In Ghana Cedis)	-0.060	(0.278)	1.181	(0.527)*	-0.375	(0.373)	-0.041	(0.279)
Social connectedness score	0.257	(0.215)	-0.302	(0.415)	0.427	(0.273)	0.259	(0.215)
SRH (Excellent/Very Good) ^a	1		1		1		1	
Good	-0.012	(0.349)	-0.322	(0.609)	-0.001	(0.448)	-0.019	(0.349)
Fair	0.894	(0.333)**	1.640	(0.592)**	0.527	(0.429)	0.908	(0.333)**
Poor	2.122	(0.357)***	2.240	(0.601)***	1.995	(0.458)***	2.149	(0.359)***
Psychological Distress Score (10–50)	0.513	(0.202)*	1.293	(0.432)**	0.277	(0.248)	0.518	(0.203)*
Interaction term: Retirement × PA							-0.431	(0.485)**
Adjusted Pseudo R ²	0.400		0.510		0.371		0.401	

Table 5. Predicting FI Score with Retirement and PA Score among ≥ 65 age group: OLS Regression Models. β—Coefficients; SE—robust standard errors are presented in parentheses; FI—functional impairment; AP—physical activity; SRH—self-rated health; ^aReference Group. ****p* < 0.001; ***p* < 0.005; **p* < 0.05.




Figure 3. Physical activity (PA, the moderator) dampens the association of retirement with functional impairment (FI) in the ≥ 65 age group.

functional impairment in a dose–response manner in the overall sample and in the men sub-group but not in women. The stratified analysis for the ≥ 65 -year group observed a relatively reduced estimate. Additionally, PA moderated the relationship between retirement and functional impairment: the risk of impairment after retirement grew weaker for those engaged in PA. This moderation suggests that PA potentially attenuates the links between retirement and functional impairment. To our knowledge, this is the first study to estimate the effect modification of the association between retirement and functional impairment by PA and gender, particularly in the LMICs context. Although retirement remains inevitable, our findings emphasize the potential of PA in ensuring functional ability during retirement.

Previous research and interpretation of the findings. As noted, research testing the association of retirement with functional impairment is limited in the SSA context, but our findings are aligned with prior studies reporting a positive retirement–functional impairment relationship in Western and Asian countries. For example, among 21,608 Australians from the 45 and Up Study, Byles et al.⁴⁰ found a strong association between retirement and physical dysfunction over time. Analyzing data from the Health and Retirement Study (HRS) ($n = 17,844$), Stenholm et al.¹⁹ intimated that physical function declines faster in retirement than in full-time work. Dave et al.¹⁴ utilized a sample of 77,194 adults and found that a 6-year average post-retirement caused a 5–16% increase in mobility and ADL difficulties and a 5–6% increase in illness conditions. Calvo et al.⁴¹ reported that early retirement dampens subjective physical health among Americans. Xue et al.⁴² found that verbal memory function declined after retirement among 3,433 individuals from the Whitehall II Cohort Study. According to Hale et al.⁴³, postponed retirement benefits cognitive function across diverse populations. However, our findings are inconsistent with previous studies suggesting better health outcomes during retirement^{44,45}. In addition, Mein et al.¹⁸ in the Whitehall II longitudinal study, found no association of retirement at age 60 with physical health functioning.

Indeed, the physical health effects of retirement remains far from certain¹³. While some studies included small sample sizes⁴⁶, others involved only specific adult population groups⁴⁷. Crucially, the evidence from previous studies may not fully apply to the SSA context given the diversity of sociocultural, and demographic conditions, including aging dynamics. The current study builds on the previously published literature by providing new evidence of the effect modification of the association of retirement with functional impairment by PA and gender in the SSA population. Our findings may present important implications for relevant social policy and public health interventions aimed at improving functional independence and successful aging.

Retirement may be an important contributor to functional impairment through several plausible pathways. First, retirement may lead to disengagement and dissociation from an economic activity which, in turn, causes a feeling of alienation, powerlessness, social anxiety, and somatic disorders^{48,49}. These mental stressors have been identified as key risk factors of functional limitations²¹. Second, having a satisfying job can be central to self-concept and personal identity⁵⁰ and the exit from such jobs can make one feel lost through negative alterations in mood. This can potentially lead to a decline in perceived physical health⁵¹. Third, many people in SSA do not plan adequately for their retirement, largely due to unemployment, low-income levels, and the predominant

contribution of the informal economic sector³⁷. The transition to retirement could, therefore, spell shocks and plunge the retirees into potentially ongoing hopelessness and a feeling of uselessness. Retirement-induced sadness and emotional hyperarousal symptoms could neurologically damage physical functioning capacities, given that prior research has strongly linked mental distress in old age to functional impairment⁵².

Fourth, the paltry sum paid in many middle and lower-income and some high-income countries as pensions due to inadequate government/state pension schemes may contribute to the worsening economic well-being of retirees. Poor economic well-being has also been related to declining health status, including physical health^{53,54}. Moreover, the retirement-induced role loss can dampen the hitherto, social networks and social interactions. Retirement, to a larger extent, shuts mobility and social connectedness which may lead to solitude, loneliness, and the likelihood of drug use. These circumstances, in turn, may contribute to poor physical health outcomes in old age⁵⁵. Interestingly, as observed in our own study population, the magnitude of the effect of retirement on functional impairment appears to attenuate with advancing age. This suggests that older people might have developed coping strategies and become more resilient against the momentous retirement shocks over time.

Crucially, our analysis found effect modification of the association of retirement with functional impairment by PA. Thus, the engagement in PA after retirement was found to retard the risk of functional impairment. This observation is distinctive given that previous studies have not examined the interactive effect of PA and retirement on functional impairment. Prior research has shown that moderate-to-vigorous PA, which is likely less common after retirement, is associated with a lower risk of functional impairment³⁵. As a health promotion mechanism, PA may potentially stimulate body and mind coordination to control physical functioning via neurotransmitters with a likely pathway that PA is a potent stimulus of immune function⁵⁶. PA may also engender resourceful social interaction crucial to improving physical health outcomes, particularly in old age³⁰.

Our study indicates that the majority of those who retired were women. This finding is similar to a recent report from the Longitudinal Social Protection Survey in which women were found to leave the labor market earlier than men among Latin American populations²⁵. In addition, the risk of functional impairment after retirement differed by gender in specific subgroups. The association between retirement and functional impairment among women did was not robust. These observations are similar to results from a study in the USA that found differences in the retirement-cognitive decline association between men and women⁴³. These findings might be, at least in part, due to women's commonly low-status jobs and weaker attachment to the job market, particularly in SSA. Women are more likely to have employment fragmented by longer career breaks, unemployment spells, part-time jobs, and have as a result fewer pensions than men^{12,28,57}. In the African socio-cultural and traditional system, men are considered the breadwinners while women mainly perform family roles, including caregiving responsibilities. Women in many cases do not stay long in the job market, hence, retirement can have a little or no impact on their physical health outcomes.

Our study utilized a large sample size and high-quality data from a representative survey. This allowed the unique opportunity to analyze the effect modification of the association of retirement with functional impairment by PA and gender in Ghana for the first time. Despite these strengths, the study has several limitations. Although we hypothesized retirement to influence functional impairment, the present study cannot determine causality, and it is important to note that the associations of retirement and functional impairment are most likely bidirectional. Future studies might, usefully consider longitudinal analysis to replicate or validate these relationships. Although rigorous adjustments in regression models were undertaken, we were unable to rule out potential confounding incidence by unobserved factors. It should also be noted that the measurement of retirement, functional impairment, and PA was based on self-report which had not been validated in the study context. Recall bias is also inevitable.

Public health and policy implications. The findings from the present study contribute to the published literature on the physical health impacts of retirement. Our study also provides important implications for policy and public health interventions targeted at improving physical health outcomes during retirement, and old age in general. Policymakers should make concerted efforts to enact policies to prolong the retirement age and also offer near-retirement work incentives to encourage working beyond a certain limit, particularly among men who may have the capacity to work. In addition, measures to promote physical health in later life should be implemented during retirement and often planned and initiated well in advance of retirement. Creating the enabling environment for regular effective PA and exercise modules for older people may be desirable in retarding or delaying the onset of functional impairments after retirement³⁰. Policies that encourage PA would also be potent in buffering the retirement-induced sedentary lifestyle, immobility, loneliness, and their concomitant functional health effects. Promoting functional health and well-being in retirees may be of utmost importance, and this approach may consist of public education programs, clinical therapy, and behavioral change interventions.

Conclusions

This representative study from the SSA context has shown that functional impairment in old age, particularly among men increases after retirement. Although retirement might be one promising risk factor for physical health overall, the association non-existed in women. The findings provide additional evidence that the adverse health effects of retirement are mitigated if older persons continue to engage in PA post-retirement. This indicates that work may be underscored as a fundamental form of PA for many older people. Future research would benefit from replicating the current study using more longitudinally robust data in other settings and the potential mediating variables in the association.

Methods

Participants. Data for this analysis came from the AgeHeaPsyWel-HeaSeeB, a nationally representative sample of community-dwelling adults aged ≥ 50 years. The original study investigated the specific impacts of the sociopolitical, informal social support, and socio-demographic inequalities on the health, well-being, and health-seeking options of older people in Ghana⁵⁸. The design was based upon a multi-stage stratified cluster sampling of households within representative statistical areas delineated by geographical and socio-demographic criteria. Briefly, the area was demarcated into three sub-regional sectors based on geographic and locational uniqueness, and two districts were randomly selected from each sub-regional zone. The selected districts were demarcated into rural and urban areas based on the GSS's³⁷ classification; nine urban and 15 rural neighborhoods were randomly selected.

We estimated the sample size (N) following Lwanga et al.⁵⁹: $N = \text{design effect} \times \left[\left(Z_{\alpha/2} \right)^2 \times P(1 - P) \right] / \varepsilon^2$. We assumed a 5% margin of error, 95% confidence interval, design effect of 1.5, 5%, and 15% of type 1 and type 2 errors respectively, and a conservative prevalence of 50% given that the actual proportion of ≥ 50 olds was unknown. This sample achieved a statistical power of 85% and 5% (two-sided) significance level to detect an odds ratio of ≥ 2 and the required minimum sample size was 901. We oversampled by 28% to cater for potential non-responses and to improve generalizability. Therefore, 1247 older persons were recruited via systematic random sampling technique. We excluded those who were not available during data collection ($n = 17$) and those who declined to participate ($n = 11$). Final analyses were performed for 1201 older adults after excluding participants whose essential data were missing ($n = 15$) or contained outliers ($n = 3$) (Fig. 1). Data were collected by trained interviewers and informed written or oral consent was obtained from all participants. The study protocol received ethics approval from the Committee on Human Research Publication and Ethics, School of Medical Sciences, Kwame Nkrumah University of Science and Technology, and Komfo Anokye Teaching Hospital, Kumasi, Ghana (Ref: CHRPE/AP/507/16). In addition, the research ethics committee of Lignan University approved the study and all the procedures involved in this study. The methods were performed taking into account the relevant guidelines, standards, and regulations.

Measures. *Gender and potential confounders.* Gender differences were self-reported as male or female. Other demographic covariates included age (in years), residence (rural, urban), and marital status (currently married, widowed, divorced/separated). Socioeconomic covariates included education reflected the highest obtained education (primary or none, secondary, tertiary), income (in Ghana Cedis), social connectedness/participation reflected past 30-day involvement, and social participation in the community (never, less frequently, frequently, very frequently, every day). The overall score ranged from 0 to 20 with a higher score reflecting higher social connectedness. Self-rated health was assessed with a single-item measure of the Short Form Health Survey 36 Questionnaire (very good, good, fair, and poor)⁶⁰. Mental health was measured by the Kessler Psychological Distress Scale (K-10) (none of the time, a little of the time, some of the time, most of the time, all of the time)²⁰ with a sum score ranging from 0 to 40 with higher scores reflecting higher psychological distress outcomes.

Functional impairment. Self-reported difficulty in conducting ADL in the preceding 30 days commonly used to gauge older people's daily performance⁶¹ with good psychometric properties⁶² was used as proxies to assess functional impairment. These were based on the six items on eating, bathing or washing the whole body, dressing up or putting on clothes, getting in or out of bed, reaching and using the toilet, and moving around inside your house. These responses were measured on a four-point scale ranging from (0) not limited at all (1) less limited (2) somewhat limited and (3) much limited. The total score ranged from 0 to 18 with a higher score indicating greater functional impairment. The physical function scale has strong validity and internal consistency ($\alpha = 0.83$).

Retirement. Retirement is defined as being out of the labor force with no intent to seek employment in the future. Based on this definition, the study sample was considered as retired when they had not been working for pay for at least one year⁶³, and they self-reported that they retired at the time of the interview⁹. This reflected a self-reported retirement status or self-reported employment status. The analysis focuses on retirees rather than non-workers as many published literature refers exclusively to the former. Prior research used a similar approach with the rationale that being out of the labor force does not instantaneously affect health status⁶³. The retirement status variable was, therefore, constructed to indicate whether individuals were retirees (those who had not worked for at least a year and considered themselves as retired) (coded 1) or non-retirees (those working for pay, had not worked for less than a year, and/or did not consider themselves as retired) (coded 0).

Physical activity (PA). PA was assessed using the International Physical Activity Questionnaire short form (IPAQ-SF). The IPAQ-SF is a validated screening tool that measures three dimensions of sitting time and PA intensity (low, moderate, and vigorous) over the past seven days as part of the daily activities of adults. The IPAQ-SF is used to calculate total physical activity in metabolic equivalent (MET) energy expenditure-min per week⁶⁴. Respondents were asked: "During the last 7 days, on how many days (1) ...did you walk for at least 10 min at a time including walking at work, at home, and to travel from place to place? (2) ...did you do moderate physical activities like gardening, cleaning, bicycling at a regular pace, swimming, or other fitness activities? (3) ...did you do vigorous physical activities like heavy lifting, digging, heavier garden, or construction work, chopping woods, aerobics, jogging/running, or fast bicycling?" The responses were taken on a continuous scale. The higher the score the higher the level of PA. The IPAQ-SF has been validated in the older African population with good reliability/validity⁶⁵. The physical activity scale has strong validity and reliability ($\alpha = 0.89$).

Statistical analysis. Descriptive analyses were first performed to describe the sample and reported as means and standard deviations for continuous variables, or counts and percentages for categorical variables. We performed Pearson's zeroth-order correlations of relevant exposure variables with functional impairment outcome. The p -value was adjusted for the multiple correlations, which can increase the risk of a type I error, i.e., to erroneously conclude the presence of a significant correlation. Rao Scott contingency table χ^2 statistics were used for simple comparisons of categorical variables and a student t -test was performed for continuous variables between gender subgroups and retirement status. The 95% confidence intervals (CI) for each variable category were estimated.

A three-stage OLS model was built in which the ADL score was regressed on the study variables to test the adjusted effect of retirement on functional impairment outcome. First, the functional impairment score was regressed on the retirement status in the full sample adjusting for potential covariates. Next, stratified analysis by gender was conducted to investigate whether the effect of retirement on functional impairment could be tempered by gender differences. Finally, we included the interaction term, retirement \times PA, in the overall sample to evaluate the effect modification of the association of retirement with functional impairment by PA. In addition, a simple slope test was performed by testing the conditional effects at one SD above and below the mean when the interaction effect reached significance. The auxiliary analysis considered a within-employment status variability modeling. Thus, we stratified the sample into three employment cohorts: employed (44.4%), unemployed (39.9%), and retired (15.7%) to understand any employment-related differences in the effect of retirement on functional impairment. Cluster-robust standard errors were calculated. We checked for multicollinearity by computing the Variance Inflation Factor (VIF) but the VIF scores ranged ≤ 1.3 , indicating no multicollinearity. All analyses were conducted using IBM-SPSS V.25 Software and the level of significance was $p < 0.05$ (two-tailed).

Ethical approval. The study protocol received ethics approval from the Committee on Human Research Publication and Ethics, School of Medical Sciences, Kwame Nkrumah University of Science and Technology, and Komfo Anokye Teaching Hospital, Kumasi, Ghana (Ref: CHRPE/AP/507/16).

Informed consent. Written informed consent was obtained from all individual participants included in the study.

Data availability

Those interested in the data and materials in this paper should contact the corresponding authors, Razak M. Gyasi at the email address RYyasi@aphrc.org.

Code availability

Data reported in this study is not freely available online but would be made available upon request.

Received: 22 January 2022; Accepted: 21 July 2022

Published online: 27 July 2022

References

1. World Health Organization. *International classification of functioning, disability and health (ICF)* Geneva: World Health Organization (2001).
2. Mlinac, M. E. & Feng, M. C. Assessment of activities of daily living, self-care, and independence. *Arch. Clin. Neuropsychol.* **31**(6), 506–516 (2016).
3. WHO. Disability and health. WHO, Geneva (2020). <https://www.who.int/news-room/factsheets/detail/disability-and-health>.
4. Hosseinpoor, A. R. *et al.* Socio-demographic patterns of disability among older adult populations of low-income and middle-income countries: Results from World Health Survey. *Int J Public Health* **61**(3), 337–345 (2015).
5. Visagie, S. *et al.* Factors related to environmental barriers experienced by persons with and without disabilities in diverse African settings. *PLoS ONE* **12**(10), e0186342 (2017).
6. Gyasi, R. M. & Phillips, D. R. Gender, self-rated health and functional decline among community-dwelling older persons. *Arch. Geront. Geriatrics* **77**, 174–183 (2018).
7. United Nations. Convention on the rights of persons with disabilities and optional protocol. New York (2006). <https://www.un.org/development/desa/disabilities/convention-on-the-rights-of-persons-with-disabilities/convention-on-the-rights-of-persons-with-disabilities-2.html>.
8. Rowe, J. W. & Kahn, R. L. Successful aging. *The Gerontologist* **37**(4), 433–440 (1997).
9. Meng, A., Nexø, M. A. & Borg, V. The impact of retirement on age related cognitive decline—A systematic review. *BMC Geriatr.* **17**, 160 (2017).
10. Oi, K. Does gender differentiate the effects of retirement on cognitive health?. *Res. Aging* **41**(6), 575–601 (2019).
11. Nexø, M. A. *et al.* Depressive symptoms and early retirement intentions among Danish eldercare workers: Cross-sectional and longitudinal analyses. *BMC Public Health* **15**, 677 (2015).
12. Kim, J. E. & Moen, P. Retirement transitions, gender, and psychological wellbeing: A life-course, ecological model. *J. Gerontol B Psychol. Sci. Soc. Sci.* **57**, 212–222 (2002).
13. van der Heide, I. *et al.* Is retirement good for your health? A systematic review of longitudinal studies. *BMC Public Health* **13**, 1180 (2013).
14. Dave, D., Rashad, I. & Spasojevic, J. The effects of retirement on physical and mental health outcomes. *South. Econ. J.* **75**, 497–523 (2008).
15. Gayman, M. D., Pai, M., Kail, B. L. & Taylor, G. Limitations pre- and postretirement: exploring racial differences reciprocity between depressive symptoms and physical. *J. Aging Health* **25**, 555 (2013).
16. Tuomi, K., Järvinen, E., Eskelinen, L., Ilmarinen, J. & Klockars, M. Effect of retirement on health and work ability among municipal employees. *Scand. J. Work Environ. Health* **17**(suppl 1), 75–81 (1991).
17. Ekerdt, D. J., Baden, L., Bossé, R. & Dibbs, E. The effect of retirement on physical health. *Am. J. Public Health* **73**(7), 779–783 (1983).

18. Mein, G., Martikainen, P., Hemingway, H., Stansfeld, S. & Marmot, M. Is retirement good or bad for mental and physical health functioning? Whitehall II longitudinal study of civil servants. *J. Epidemiol. Commun. Health* **57**(1), 46–49. <https://doi.org/10.1136/jech.57.1.46> (2003).
19. Stenholm, S. *et al.* Age-related trajectories of physical functioning in work and retirement: the role of socio-demographic factors, lifestyle and disease. *J. Epidemiol. Community Health* **68**, 503–509 (2014).
20. Kessler, R. C. The effects of stressful life events on depression. *Annu. Rev. Psychol.* **48**, 191–214. <https://doi.org/10.1146/annurev.psych.48.1.191> (1997).
21. Tanner, J. *et al.* Psychiatric symptom profiles predict functional impairment. *Front. Psychiatry* **10**, 37 (2019).
22. Gyasi, R. M., Phillips, D. R. & David, R. Explaining the gender gap in health services use in Ghanaian community-dwelling older cohorts. *Women Health* **59**(10), 1089–1104 (2019).
23. Oksuzyan, A. *et al.* Sex differences in health and mortality in Moscow and Denmark. *Eur. J. Epidemiol.* **29**(4), 243–252 (2014).
24. Tareque, M. I., Tiedt, A. D., Islam, T. M., Begum, S. & Saito, Y. Gender differences in functional disability and self-care among seniors in Bangladesh. *BMC Geriatr.* **17**, 177 (2017).
25. Canales, A., Salinas, V. & Biehl, A. Gender Differences in Retirement Behavior: How Family Work, and Pension Regime Explain Retirement in Chile. *Work Aging Retirement* **7**(2), 107–128 (2021).
26. Radl, J. Labor market exit and social stratification in Western Europe: The effects of social class and gender on the timing of retirement. *Eur. Sociol. Rev.* **29**(3), 654–668. <https://doi.org/10.1093/esr/jcs045> (2013).
27. Aaron, H. J., & Callan, J. M. Who retires early? Boston College Center for Retirement (Research Working Paper No. 2011–10) (2011). <https://ssrn.com/abstract=1857145>.
28. Finch, N. Why are women more likely than men to extend paid work? The impact of work–family life history. *Eur. J. Ageing* **11**, 31–39 (2014).
29. König, S. Career histories as determinants of gendered retirement timing in the Danish and Swedish pension systems. *Eur. J. Ageing* **14**, 1–10. <https://doi.org/10.1007/s10433-017-0424-5> (2017).
30. Gyasi, R. M. Social support, physical activity and psychological distress among community-dwelling older Ghanaians. *Arch Geront Geriatrics* **81**, 142–148 (2019).
31. WHO World Report on Ageing and Health Geneva: WHO (2015). <http://www.who.int/ageing/events/world-report-2015-launch/en/>. 18/10/2016.
32. Balogun, S., Scott, D., Cicutinni, F., Jones, G. & Aitken, D. Longitudinal study of the relationship between physical activity and knee pain and functional limitation in community-dwelling older adults. *Arch. Gerontol. Geriatr.* **90**, 104101. <https://doi.org/10.1016/j.archger.2020.104101> (2020).
33. Roberts, C. E., Phillips, L. H., Cooper, C. L., Gray, S. & Allan, J. L. Effect of different types of physical activity on activities of daily living in older adults: Systematic review and meta-analysis. *J. Aging Phys. Act.* **25**(4), 653–670. <https://doi.org/10.1123/japa.2016-0201> (2017).
34. Garcia Meneguci, C. A., Meneguci, J., Sasaki, J. E., Tribess, S. & Júnior, J. S. V. Physical activity, sedentary behavior and functionality in older adults: A cross-sectional path analysis. *PLoS ONE* **16**(1), e0246275. <https://doi.org/10.1371/journal.pone.0246275> (2021).
35. Paterson, D. H. & Warburton, D. E. Physical activity and functional limitations in older adults: A systematic review related to Canada's Physical Activity Guidelines. *Int. J. Behav. Nutr. Phys. Act.* **7**, 38. <https://doi.org/10.1186/1479-5868-7-38> (2010).
36. Martínez-Hernández, B. M. *et al.* Association between physical activity and physical and functional performance in non-institutionalized Mexican older adults: a cohort study. *BMC Geriatr.* **22**, 388. <https://doi.org/10.1186/s12877-022-03083-7> (2022).
37. GSS. Ghana Living Standards Survey (GLSS) 7. Government of Ghana (2019). https://statsghana.gov.gh/gssmain/fileUpload/press%20release/GLSS7%20MAIN%20REPORT_FINAL.pdf.
38. Kpessah, M. W. Retirement income security under Ghana's three-tier pension model: Assessment of risks and options for reform. *Pensions* **16**(2), 127–136 (2011).
39. Jensen, L. D., Ryom, P. K., Christensen, M. V. & Andersen, J. H. Differences in risk factors for voluntary early retirement and disability pension: A 15-year follow-up in a cohort of nurses' aides. *BMJ Open* **2**, e000991 (2012).
40. Byles, J. E. *et al.* Gender, mental health, physical health and retirement: A prospective study of 21,608 Australians aged 55–69 years. *Maturitas* **87**, 40–48 (2016).
41. Calvo, E., Sarkisian, N. & Tamborini, C. R. Causal effects of retirement timing on subjective physical and emotional health. *J. Gerontol. B Psychol. Sci. Soc. Sci.* **68**(1), 73–84 (2013).
42. Xue, B. *et al.* Effect of retirement on cognitive function: The Whitehall II cohort study. *Eur. J. Epidemiol.* **33**(10), 989–1001 (2018).
43. Hale, J. M., Bijlsma, M. J. & Lorenti, A. Does postponing retirement affect cognitive function? A counterfactual experiment to disentangle life course risk factors. *SSM Popul. Health* **26**(15), 100855. <https://doi.org/10.1016/j.ssmph.2021.100855> (2021).
44. Eibich, P. Understanding the effect of retirement on health: Mechanisms and heterogeneity. *J. Health Econ.* **43**, 1–12 (2015).
45. Jokela, M. *et al.* From midlife to early old age: health trajectories associated with retirement. *Epidemiology* **21**(3), 284–290. <https://doi.org/10.1097/EDE.0b013e3181d61f53> (2010).
46. LeMasters, G., Bhattacharya, A., Borton, E. & Mayfield, L. Functional impairment and quality of life in retired workers of the construction trades. *Exp. Aging Res.* **32**(2), 227–242. <https://doi.org/10.1080/03610730600554065> (2006).
47. Hallberg, D., Johansson, P. & Josephson, M. Is an early retirement offer good for your health? Quasi-experimental evidence from the army. *J. Health Econ.* **44**, 274–285 (2015).
48. Farré, L., Fasani, F. & Mueller, H. Feeling useless: the effect of unemployment on mental health in the Great Recession. *IZA J. Labor. Econ.* **7**, 8 (2018).
49. Gallo, W. T., Bradley, E. H., Siegel, M. & Kasl, S. V. Health effects of involuntary job loss among older workers: Findings from the Health and Retirement Survey. *J. Gerontol.: Ser. B* **55**(3), S131–S140 (2000).
50. Szabó, A., Allen, J., Stephens, C. & Alpass, F. Is retirement associated with physical health benefits? A longitudinal investigation with older New Zealanders. *Age Ageing* **48**(2), 267–272 (2019).
51. Sahlgren, G. H. Work longer, live healthier: The relationship between economic activity, health and government policy. The Institute of Economic Affairs, IEA Discussion Paper No. 46 (2013). <https://www.iea.org.uk/wp-content/uploads/2016/07/Work%20Longer%20Live%20Healthier.pdf>
52. Stoeckel, K. J. & Litwin, H. The impact of social networks on the relationship between functional impairment and depressive symptoms in older adults. *Int. Psychogeriatr.* **28**(1), 39–47 (2016).
53. Macintyre, A., Ferris, D., Gonçalves, B. & Quinn, N. What has economics got to do with it? The impact of socioeconomic factors on mental health and the case for collective action. *Palgrave Commun.* **4**, 10 (2018).
54. Wang, J. & Geng, L. Effects of socioeconomic status on physical and psychological health: Lifestyle as a mediator. *Int. J. Environ. Res. Public Health* **16**(2), 281. <https://doi.org/10.3390/ijerph16020281> (2019).
55. Philip, K. E. J. *et al.* Social isolation, loneliness and physical performance in older-adults: fixed effects analyses of a cohort study. *Sci. Rep.* **10**, 13908 (2020).
56. WHO. Global action plan on physical activity 2018–2030: more active people for a healthier world. Geneva: World Health Organization; 2018. Licence: CC BY-NC-SA 3.0 IGO (2018). <https://www.who.int/publications/i/item/9789241514187>.
57. Riekhoff, A. J., Järnefelt, N. & Laaksonen, M. Workforce composition and the risk of labor market exit among older workers in Finnish companies. *Work Aging Retirement.* **6**(2), 88–100. <https://doi.org/10.1093/workar/waz023> (2020).

58. Gyasi RM (2018) Ageing, health and health-seeking behaviour in Ghana (Doctor's thesis, Lingnan University, Hong Kong). <https://commons.ln.edu.hk/otd/41/>.
59. Lwanga, S. K., Lemeshow, S., & World Health Organization. Sample size determination in health studies: A practical manual/S. K. Lwanga and S. Lemeshow. World Health Organization (1991). <https://apps.who.int/iris/handle/10665/40062>.
60. Ware, J. E. IQOLA: The SF-36 Health Survey: Development and Use in Mental Health Research and the IQOLA Project. *Int. J. Ment. Health* **23**(2), 49–73 (1994).
61. WHO. *World Health Day 2012: Ageing and Health: Toolkit for Event Organizers 2012* (World Health Organization, 2012).
62. Ustun, T. B., Kostanjsek, N., Chatterji, S., Rehm, J., & WHO Measuring health and disability: manual for WHO Disability Assessment Schedule (WHODAS 2.0) / edited by T.B. Üstün, N. Kostanjsek, S. Chatterji, J.Rehm. World Health Organization (2010). <https://apps.who.int/iris/handle/10665/43974>.
63. Bonsang, E., Adam, S. & Perelman, S. Does retirement affect cognitive functioning?. *J. Health Econ.* **31**, 490–501 (2012).
64. Craig, C. L. *et al.* International physical activity questionnaire: 12-country reliability and validity. *Med. Sci Sports Exerc.* **35**(8), 1381–1395. <https://doi.org/10.1249/01.MSS.0000078924.61453.FB> (2003).
65. Kolbe-Alexander, T. L., Lambert, E. V., Harkins, J. B. & Ekelund, U. Comparison of two methods of measuring physical activity in South African older adults. *J. Aging Phys. Act.* **14**(1), 98–114 (2006).

Author contributions

All authors contributed to the conceptualization and the design of the study. Material preparation, data collection, and analysis were performed by R.M.G., P.A.A., S.A., L.P.S., F.F., and R.R. The first draft of the manuscript was written by R.M.G. and all authors commented on previous versions of the manuscript. The supervision of data acquisition, data analysis, and interpretation was done by D.R.P. All authors reviewed, revised, read, and approved the final manuscript.

Funding

This work was supported by Lingnan University, Hong Kong [Grant Number: RPG1129310]. The sponsor played no role in the design, execution, analysis, and interpretation of data and preparation of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to R.M.G.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022