

RESEARCH

Open Access

Phosphoproteomic analysis of the non-seed vascular plant model *Selaginella moellendorffii*

Xi Chen^{1,2†}, Wai Lung Chan^{1†}, Fu-Yuan Zhu¹ and Clive Lo^{1*}

Abstract

Background: *Selaginella* (*Selaginella moellendorffii*) is a lycophyte which diverged from other vascular plants approximately 410 million years ago. As the first reported non-seed vascular plant genome, *Selaginella* genome data allow comparative analysis of genetic changes that may be associated with land plant evolution. Proteomics investigations on this lycophyte model have not been extensively reported. Phosphorylation represents the most common post-translational modifications and it is a ubiquitous regulatory mechanism controlling the functional expression of proteins inside living organisms.

Results: In this study, polyethylene glycol fractionation and immobilized metal ion affinity chromatography were employed to isolate phosphopeptides from wild-growing *Selaginella*. Using liquid chromatography-tandem mass spectrometry analysis, 1593 unique phosphopeptides spanning 1104 non-redundant phosphosites with confirmed localization on 716 phosphoproteins were identified. Analysis of the *Selaginella* dataset revealed features that are consistent with other plant phosphoproteomes, such as the relative proportions of phosphorylated Ser, Thr, and Tyr residues, the highest occurrence of phosphosites in the C-terminal regions of proteins, and the localization of phosphorylation events outside protein domains. In addition, a total of 97 highly conserved phosphosites in evolutionary conserved proteins were identified, indicating the conservation of phosphorylation-dependent regulatory mechanisms in phylogenetically distinct plant species. On the other hand, close examination of proteins involved in photosynthesis revealed phosphorylation events which may be unique to *Selaginella* evolution. Furthermore, phosphorylation motif analyses identified Pro-directed, acidic, and basic signatures which are recognized by typical protein kinases in plants. A group of *Selaginella*-specific phosphoproteins were found to be enriched in the Pro-directed motif class.

Conclusions: Our work provides the first large-scale atlas of phosphoproteins in *Selaginella* which occupies a unique position in the evolution of terrestrial plants. Future research into the functional roles of *Selaginella*-specific phosphorylation events in photosynthesis and other processes may offer insight into the molecular mechanisms leading to the distinct evolution of lycophytes.

Keywords: *Selaginella*, IMAC, PEG fractionation, Phosphosites, Phosphorylation motif

Background

Selaginella (*Selaginella moellendorffii*) is a lycophyte believed to be originated from the earliest vascular plants approximately 410 million years ago [1]. Although lycophytes have existed twice as long as angiosperms, they have not evolved flowers and seeds since their divergence from other plant lineages. For this reason, *Selaginella* has

been selected as a model plant to understand the early evolution of developmental and metabolic processes that are unique to vascular plants [2]. After a bacterial artificial chromosome library was constructed from clonally propagated plants [3], the complete *Selaginella* genome sequence was released in 2007 [4]. Subsequently, a number of investigations on *Selaginella* were launched in different areas including gene evolution [5-10], pathway conservation [11-16], genomic DNA composition and methylation [17-19], sRNA functions and RNA editing [20,21], and transposons [22]. Interestingly, *Selaginella* was found to utilize genes significantly different from flowering plants

* Correspondence: clivelo@hku.hk

†Equal contributors

¹School of Biological Sciences, The University of Hong Kong, Pokfulam Hong Kong, China

Full list of author information is available at the end of the article

to generate secondary metabolites with potentials for pharmaceutical applications [23-26]. Meanwhile, proteomic investigations on this non-seed vascular plant model have not been extensively reported. A two-dimensional electrophoresis-based approach was recently employed to explore the desiccation tolerance mechanism in the resurrection plant *Selaginella tamariscina* [27].

Post-translational modifications (PTMs) play important roles in the regulation of protein functions and they occur at distinct amino acid side chains or peptide linkages. It has been estimated that more than 200 types of PTMs exist in proteins [28]. Protein phosphorylation, principally on serine, threonine or tyrosine residues, is one of the most important and well-investigated PTMs. It represents a reversible molecular switch controlled by protein kinases and protein phosphatases, either activating or inactivating the target proteins [29]. Approximately one-third of all proteins in eukaryotic cells were estimated to be phosphorylated at any given time [30]. In plants, protein phosphorylation plays a central role in virtually all cellular processes, including carbon and nitrogen metabolism, growth and development, transcription and translation, responses to abiotic and biotic stresses, cell cycle, and apoptosis [31]. Therefore, the identification of protein kinases and phosphatases, their substrates, and the phosphorylation sites involved is crucial for the understanding of many fundamental processes in plants. Interestingly, *Arabidopsis* contains over 1000 protein kinases [32], which is twice as many as those in human, while the two genomes share similar number of genes [33]. Hence, protein phosphorylation events in plants appear to be very different and more complicated than those in mammals. In fact, a number of plant protein kinases implicated in early events of signal transduction are unique with no mammalian orthologs [34].

Phosphoproteomic investigations in plants were initiated in recent years following the completion of different genome sequencing projects. The highly abundant ribulose-1,5-bisphosphate carboxylase/oxygenase (RUBISCO) protein, which accounts of about 50% of total soluble proteins [35], hindered the detection of low-abundant proteins including many phosphoproteins [36]. Polyethylene glycol (PEG) fractionation has been used as a cost-effective and contaminant-free procedure to remove RUBISCO for improved detection of low-abundant proteins [37-39]. In addition, phosphopeptide enrichment procedures, such as immobilized metal ion affinity chromatography (IMAC), are necessary to reduce the complexity of proteolyzed lysates for mass spectrometry analysis. IMAC is based on affinity purification through metal complexation with the phosphate group in phosphopeptides [40] and it has been adopted in Phosphoproteomic analysis in different plant systems [41-46].

In the present study, we used the PEG fractionation approach followed by the IMAC procedure to prepare *Selaginella* samples for phosphoproteome profiling and identified 1588 unique phosphorylation sites. Our dataset revealed features that are consistent with the *Arabidopsis* phosphoproteome. We further identified phosphorylation events that are conserved between *Selaginella* and angiosperm orthologous sequences. Novel and unique phosphosites were detected in several photosynthesis-related proteins in *Selaginella*. Phosphorylation motifs recognized by known protein kinase classes were revealed for both evolutionarily conserved and *Selaginella*-specific proteins.

Results and discussion

General features of the *Selaginella* phosphoproteome dataset

We employed the procedures of PEG fractionation (Additional file 1: Figure S1) and IMAC enrichment to isolate phosphopeptides from wild-growing *Selaginella* (*Selaginella moellendorffii*) for LC-MS/MS analysis. A total of 1593 unique phosphopeptides containing 1588 non-redundant phosphosites were discovered in our study (Additional file 2: Table S1). Among them, 1104 were identified with high confidence of localization (localization probability $\geq 95\%$), 116 with median confidence of localization ($80\% \leq$ localization probability $< 95\%$), and 368 with low confidence of localization (localization probability $< 80\%$). Phosphosites with high confidence of localization were categorized into pSer (86.2%), pThr (13.3%), and pTyr (0.5%). The relative distribution of the three phosphorylated residues is consistent with previous reports for different flowering plant species [45,46]. As Ser/Thr kinases are commonly encoded in plant genomes, more frequent Ser and Thr phosphorylation events are expected. On the other hand, while typical Tyr-specific kinases are absent in plant genomes, a few plant kinases with dual specificity are believed to phosphorylate Tyr residues in proteins [47].

The 1104 confirmed phosphosites correspond to a total of 716 *Selaginella* proteins, 665 of them can be assigned to orthologous protein groups (Additional file 3: Table S2) using the OrthoMCL algorithm with a cut-off of E-5 e-value and 50% sequence match [48,49]. Forty two proteins are considered *Selaginella*-specific proteins since they could not be assigned to any OrthoMCL groups or do not have any matching sequences in the OrthoMCL database. These proteins may have evolved in lycophytes after their separation from other vascular plants including ferns and seed plants.



Analysis of phosphosite locations in *Selaginella* proteins

To analyze the locations of the identified phosphorylation sites, protein sequences were divided into 5% fractions and the number of phosphorylation events was counted

within each fraction. As shown in Figure 1A, the highest number of phosphosites is found in the last fraction, i.e. the C-termini of proteins. We performed parallel analysis using an Arabidopsis phosphoproteome dataset (retrieved from P³DB) and found a very similar distribution pattern for the phosphosites (Figure 1B). Such phenomenon was also described in a phosphoproteomic study of mouse liver [50]. Hence, the more frequent C-terminal phosphorylation in proteins appears to be a common feature in different organisms, including plants and animals. The C-terminal region was suggested to be more exposed and flexible for protein phosphorylation [50].

Functional categorization of the identified Selaginella phosphoproteins

To understand the functional distribution of the unique Selaginella phosphoproteins identified in this study, their cellular localization, molecular function, and biological processes were analyzed and compared with those of 2400 Selaginella proteins identified after LC-MS/MS analysis of PEG-fractionated samples without the IMAC enrichment procedure. Based on the comparison of Gene Ontology (GO) term annotations (Figure 2), the 3 most over-represented categories for the identified phosphoproteins in each GO vocabulary are: nucleus, plasma membrane and cytosol for “cellular component”; DNA/RNA binding, kinase activity, and transferase activity for “molecular function”; protein modification, phosphorus metabolic process, and transcription for “biological process”.



kinase activity, and transferase activity for “molecular function”; protein modification, phosphorus metabolic process, and transcription for “biological process”.

Location of phosphosites in characterized protein domains

To determine whether the Selaginella phosphosites are located in known structural and/or functional protein domains, Pfam database search (Wellcome Trust Sanger Institute) was performed to extract domain information

of our identified phosphoproteins. A total of 594 proteins with domain information were retrieved. Among the 893 phosphosites in these proteins, only 201 (22.5%) were located inside protein domains (Table 1). Our findings are consistent with the observations from Arabidopsis phosphoproteome analysis suggesting that phosphorylation events may not have significant impact on domain-associated functions [51,52].

Phosphorylation motif analysis

A phosphorylation motif search was performed on our phosphopeptide dataset (localization probability $\geq 95\%$) using the Motif-X algorithm [53]. Peptide sequences are aligned with their length adjusted to ± 7 residues from the central phosphosite for data submission. Over-represented patterns of amino acid sequences were generated with a minimum occurrence of 20 and a significance value of 10^{-6} . All together, we obtained a total of 11 phosphorylation (9 Ser and 2 Thr) motifs containing at least one fixed amino acid aside from the central phosphorylated residue (Figure 3A). Both the Thr-motifs are Pro-targeted (TP and PXTTP) and there are 3 Ser Pro-targeted motifs (SP, PXSP, SPXR). All these motifs are possible substrates of glycogen synthase kinase 3, cyclin-dependent kinase, and mitogen-activated protein kinase. In addition, 3 basophilic motifs (LXRXXS, RXXS, KXXS) likely to be associated with the activities of Ca^{2+} -dependent protein kinase (CPK), Ca^{2+} /calmodulin-dependent protein kinase, or protein kinase A were identified. Furthermore, 3 acidic motifs (SDXE, SXD, and SE) potentially recognized by casein kinase II were generated. We also performed parallel Motif-X analysis using Arabidopsis phosphopeptides retrieved from P³DB and those obtained by Wang et al. (2013). One of the Selaginella motifs, KXXS, was not generated from the Arabidopsis analysis. Thirty two (out of 38) occurrences of such motif correspond to proteins assigned with OrthoMCL group with e-value $< E^{-50}$ (Additional file 4: Table S3), indicating that this basophilic motif is primarily associated with evolutionarily conserved proteins in Selaginella. Analysis of the 107 phosphosites in the Selaginella-specific proteins

(those without any assigned OrthoMCL groups) revealed that they are more enriched in Pro-directed motifs when compared to all the identified phosphosites (49% vs 35%) (Figure 3B). On the other hand, the basophilic motifs are under-represented in the Selaginella-specific proteins when compared to all proteins identified (23% vs 38%). Consistently, a single SP motif with 36 occurrences was generated by Motif-X analysis for the 107 phosphosites found in the Selaginella-specific proteins. Taken together, most of the Selaginella phosphorylation events identified in this study are likely to be catalyzed by known classes of protein kinase classes in plants.

Phosphorylation events in evolutionary conserved proteins

To identify phosphorylation events highly conserved between Selaginella and flowering plants, our identified phosphopeptides with confirmed phosphosites were clustered with phosphopeptides of Arabidopsis, rice, rapeseed, soybean and *Medicago truncatula* (retrieved from P³DB) by CD-HIT [54] using a sequence identity cutoff of 0.6 and an alignment bandwidth of 5 [44]. A total of 107 Selaginella phosphopeptides harboring 115 Ser/Thr phosphosites were found to cluster with phosphopeptides from the other plants. More than 80% (97/123) of those Selaginella phosphosites were found to have equivalent phosphosites in at least one other species. The majority (90/106) of the Selaginella proteins harboring the conserved phosphosites are evolutionarily conserved proteins belonging to Orthogroups identified with e values $< E^{-50}$ (Additional file 5: Table S4). Many of these proteins are involved in primary metabolism (e.g. Calvin cycle, glycolysis, TCA cycle, lipid biosynthesis), RNA processing, transcriptional regulation, cell cycle, protein phosphorylation (kinases), and signaling (e.g. G proteins, 14-3-3 protein, LRR-containing kinases). On the other hand, 17 of these highly conserved phosphorylation events are found in proteins with unknown functions. Selected phosphopeptide alignments containing highly conserved phosphosites in multiple plant species are shown in Table 2.

Furthermore, we performed a close examination on the phosphorylation events in Selaginella photosynthesis-related proteins. The molecular machinery of photosynthesis has been highly conserved during plant evolution. Among our identified phosphoproteins with confirmed phosphosites, seven are involved in photosystem II (PSII) and two are involved in photosystem I (PSI) (Figure 4A). To reveal possible evolutionary significance, sequences were aligned with orthologs from Arabidopsis, rice, and *Physcomitrella patens* (moss), representing diverse lineages of dicot, monocot, and bryophytes, respectively (Figure 4B and Additional file 6: Figure S2). In all cases, phosphorylation information is only available for the Arabidopsis proteins. Sequences of rice and moss are

Table 1 Location of phosphorylation sites on characterized protein domains

Number of proteins possessing Pfam domain	Number of phosphorylation sites ⁽¹⁾			Total (%)
	Pfam domain ⁽²⁾		Total (%)	
	Inside (%)	Outside (%)		
pS	538	163 (21.0%)	612 (79.0%)	775 (100%)
pT	102	36 (31.6%)	78 (68.4%)	114 (100%)
pY	4	2 (50%)	2 (50%)	4 (100%)
All	594	201 (22.5%)	692 (77.5%)	893 (100%)

⁽¹⁾All analyzed phosphorylation sites were confirmed with the localization probability above 95% deduced from Ascore.

⁽²⁾Location of phosphorylation sites relative to the conserved domains annotated in the Pfam database.

A

Serine/Threonine Motifs ⁽¹⁾	No. of Occurrences	Kinase Reported	Motifs Reported
.....P.sP.....	79	Extracellular signal-regulated kinase, MAPK	A.K. Pullikuth <i>et al.</i> ⁽²⁾
.....sP.R....	53	Growth associated histone H1 kinase, CDK1, 2, 4, 6, CDC2	A. Errico <i>et al.</i> ⁽³⁾
.....sP.....	280	Proline-directed kinase, MAPK, CDC2	P3DB ⁽⁴⁾ / X. Wang <i>et al.</i> ⁽⁵⁾
..L.R..s.....	51	OST1	P3DB
...R..s.....	111	SPK, CDPK	P3DB / X. Wang <i>et al.</i>
.....sD.E....	27	Casein kinase II	P3DB / X. Wang <i>et al.</i>
.....s.D.....	50	Casein kinase II	P3DB
.....sE.....	37		X. Wang <i>et al.</i>
...K..s.....	38		
....P.tP.....	26	MAPK, GSK3, CDK5	X. Wang <i>et al.</i>
.....tP.....	55	Pro-directed kinase	P3DB / X. Wang <i>et al.</i>

(1) Phosphorylation of serine or threonine as indicated as 's' or 't'.

(2) A.K. Pullikuth *et al.*, *Mol. Cell Biol.* 30: 3233-3248.

(3) A. Errico *et al.*, *Advances in Enzyme Regulation* 50: 375-399.

(4) Non-redundant phosphopeptides provided from Plant Protein Phosphorylation Database (P3DB).

(5) Phosphopeptides data obtained from X. Wang *et al.*, *J. of Proteomics* 78: 486-498.




Figure 3 Phosphorylation motifs in Selaginella phosphoproteins. (A) Motif-x analysis of phosphorylation peptides with localization probability 95%. **(B)** Motif classes distribution of Selaginella(Sm) specific phosphorylation site (P-site).

included for examination of phosphorylatable residues at equivalent sites.

As the first link in the chain of light-dependent reactions, PSII captures photons and uses the energy to extract electrons from water molecules. The light-harvesting chlorophyll a/b protein complex LHCII in PSII is composed of three proteins, namely Lhcb1, Lhcb2, and Lhcb3 [55]. Non-phosphorylated LHCII functions as an antenna for PSII, but it migrates to come in contact with PSI following light-dependent phosphorylation of Lhcb1 and Lhcb2 [56] which is likely to occur at N-terminal Thr residues [57]. Although no explicit phosphosites were identified [55-57], STN7 kinase was strongly suggested to be required for phosphorylation of Lhcb1 and Lhcb2 to achieve state transitions between PSII and PSI [56]. In Selaginella, we detected N-terminal Thr phosphosites in D8QN27 (Lhcb1: Thr-44) and D8SUF1 (Lhcb2: Thr-42) (Figure 4B). Both phosphorylation events are conserved in Arabidopsis Lhcb1 and Lhcb2 [58,59]. Equivalent phosphorylatable residues are also found in rice Lhcb1 and Lhcb2 (Figure 4B). As these Thr residues are preceded by

a basic residue (Lys or Arg), they represent potential signatures recognized by STN7 [60]. In fact, the Thr-40 in Arabidopsis Lhcb2 is phosphorylated in wild-type but not in the *stn7* mutant [59], further suggesting that it is a target of STN7. In D8QN27 (Lhcb1), we also identified the Ser-54 Pro-directed phosphosite which is conserved in Arabidopsis Lhcb1 [43] and an equivalent Ser residue in rice Lhcb1 (Figure 4B). Interestingly, all the N-terminal Ser/Thr residues mentioned above are not conserved in *P. patens* Lhcb1 and Lhcb2 and they were probably only evolved after the emergency of vascular plants. On the other hand, the Ser-48 and Ser-49 phosphosites in D8QN27 (Lhcb1) are located in a region not conserved with the Arabidopsis and rice sequences, but equivalent Ser residues are identified in moss Lhcb1. They may represent phosphorylation events that are lost in the angiosperm lineage.

Lhcb4, a minor chlorophyll-binding protein, was found to be phosphorylated in maize upon exposure to high light intensity for protection against cold stress [61]. The phosphosite Thr-112, a potential casein kinase II target,

Table 2 Selected conserved phosphorylation sites within the orthologous proteins

Accession	Phosphopeptides alignment ⁽¹⁾	Protein annotation	MapmanBin name	OrthoMCL group	E-value
D8QZX4	VLEGGsDDEGASTEAHGR	ILITYHIA	Not assigned.no ontology	OG5_128125	1E-181
AT1G64790.1	ALLEGGsDDEGASTEAAQGR				
Os03g51140.1	AILEGGsDDEGASTEAHGR				
D8RKK4	SNsFVGTTEEYIAPEIIK	KCBP-interacting protein kinase	Protein. stranslational modification	OG5_129514	1E-181
AT3G52890.2	SNsFVGTHEYLAPEIIK				
Os02g43740.1	SMsFVGTHEYLAPEIIK				
D8T2E8	ASGAFILTAshNPPGGPHEDFGIK	Phosphoglucomutase/phosphomannomutase family protein	Glycolysis.PGM	OG5_127226	1E-181
AT1G23190.1	ATGAFILTAshNPPGGPTEDFGIK				
Os03g50480.1	ATGAFILTAshNPPGGPTEDFGIK				
D8SQG3	sQAGTPEWMAPEVLR	Protein kinase superfamily protein	Signalling.MAP kinases	OG5_130251	1E-181
AT1G08720.1	sTAGTPEWMAPEVLR				
Os02g50970.2	sTAGTAEWMAPEVLR				
D8SCJ0	ALVANYNQtPR	Cell division cycle 5	RNA.regulation of transcription.MYB-related transcription factor family	OG5_128000	1E-181
AT1G09770.1	ALLANYSQtPR				
Os04g28090.1	ALLSSYSQtPR				
D8RHD1	GILAMDEsNATCGK	Aldolase superfamily protein	Glycolysis.aldolase	OG5_127143	1E-179
AT3G52930.1	GILAADEsTGTIGK				
Os05g33380.1	GILAADEsTGTIGK				
D8R365	GFVPILPGsPGSS	Na ⁺ /H ⁺ + exchanger 1	Transport.unspecified cations	OG5_126729	1E-181
AT5G27150.1	GFVPFVPGsPTER				
Os11g42790.1	GFVPFVPGsPTER				
D8QX62	PsGSPPVPMHsPPRPVTVK	Chromatin protein family	DNA.synthesis/chromatin structure	OG5_128357	1E-181
AT1G77180.1	AsGSPPVPMHsPPRPVTVK				
Os02g52250.1	AsGSPPVPMHsPPRPVTVK				
D8QU29	YHGHsMSDPGSTYR	Pyruvate dehydrogenase complex E1 alpha subunit	TCA/org.transformation.TCA. pyruvate DH.E1	OG5_127216	1E-176
AT1G59900.1	YHGHsMSDPGSTYR				
Medtr5g037700.1	YHGHsMSDPGSTYR				
D8SRX1	WGTQAPVQLGsLR	GDP-D-mannose 3',5'-epimerase	Redox. ascorbate and glutathione. ascorbate. GME	OG5_135736	1E-181
AT5G28840.1	WGTQAPVQLGsLR				
Os10g28200.1	WSTQAPVQLGsLR				
D8RPK5	AHGPAVGLPTEDDMGNsEVGHNALGAGR	Phosphoglycerate mutase, 2,3-bisphosphoglycerate-independent	Glycolysis.phosphoglycerate mutase	OG5_129051	1E-181
AT1G09780.1	AHGTAVGLPSEDDMGNsEVGHNALGAGR				
Os05g40420.3	AHGTAVGLPSDDDMGNsEVGHNALGAGR				

Table 2 Selected conserved phosphorylation sites within the orthologous proteins (Continued)


D8SC69	VQsSSAIWVHPR	Receptor protein kinase TMK1 precursor, putative, expressed	-	OG5_141123	1E-181
Os03g50810.1	VQsPHAMVVHPR				
Glyma02g40980.1	VQsPNALVIHPR				
D8S0V4	QLsIDQFENEGR	BRI1 suppressor 1 (BSU1)-like 2	Protein,postranslational modification	OG5_132764	1E-181
AT1G08420.1	QLsIDQFENEGR				
Os12g42310.1	QLsIDQFENEGR				
D8QXA8	NFRPDsLLGEGGFGSVFK	Protein kinase superfamily protein	Protein,postranslational modification. kinase.receptor like cytoplasmatic kinase VII	OG5_147118	1E-163
AT1G07570.1	NFRPDsVLGEGGFGCVFK				
Glyma01g05160.1	NFRPDsLLGEGGFGYVYK				
D8RMG5	ALsPDRNDAFAMGDK	Splicing factor, putative	RNA,processing.splicing	OG5_127822	1E-181
AT5G64270.1	VLsPDRVDAFAMGDK				
Medtr2g009180.1	ILsPDRHDAFAAGEK				
D8RZ45	LTsFEALQSATK	ADP-glucose pyrophosphorylase family protein	Major CHO metabolism.synthesis. starch.AGPase	OG5_129964	1E-181
AT1G74910.1	RVsSFEALQPATR				
Os03g11050.3	RVsSFEALHSATK				
D8QZR7	AMKsPDPLEEQR	Protein kinase superfamily protein	Protein,postranslational modification.kinase	OG5_129183	1E-149
AT1G67580.1	MVKsPDPLEEQR				
Os02g39010.3	HMKsPDPLEEQR				
D8RF63	EIsDDEEEEEK	HEAT SHOCK PROTEIN 81.4	Stress.abiotic.heat	OG5_126623	1E-181
AT5G56000.1	EIsDDEEEEEK				
Os08g39140.1	EIsDDEEEEEK				
D8QX94	tSCGSPNYAAPEVISGK	SNF1-related protein kinase 1.3	Protein,postranslational modification	OG5_126655	1E-181
AT5G39440.1	tSCGSPNYAAPEVISGK				
Os03g17980.1	tSCGSPNYAAPEVISGK				
D8SX29	GEPNISyICSR	SHAGGY-related protein kinase dZeta	Protein,postranslational modification	OG5_126888	1E-181
AT2G30980.1	GEANISyICSR				
Os02g14130.1	GEANISyICSR				
D8R4T7	GGMTsHAAWAR	Pyruvate orthophosphate dikinase	Glycolysis.PPFK	OG5_127082	1E-181
AT4G15530.3	GGMTsHAAWAR				
Os05g33570.3	GGMTsHAAWAR				
D8T8I0	GLDIDTIQQHYtV	H(+)-ATPase 1	transport,p- and v-ATPases.H + -exporting ATPase	OG5_127253	1E-181
AT2G18960.1	GLDIDTAGHHYtV				
Os04g56160.1	GLDIDTIQQNYtV				

Table 2 Selected conserved phosphorylation sites within the orthologous proteins (Continued)

D8R651	VHACVGGtDVR	Eukaryotic translation initiation factor 4A1	Protein.synthesis.initiation	OG5_126984	1E-181
AT3G13920.1	VHACVGGtSVR				
Medtr2g120800.1	VHACVGGtSVR				
D8QVP0	TIQFVDWCPtGFK	Tubulin alpha-3	Cell.organisation	OG5_126605	1E-181
AT5G19770.1	TVQFVDWCPtGFK				
Os03g51600.3	TIQFVDWCPtGFK				

⁽¹⁾Phosphorylation sites are indicated as small letters.

Arabidopsis (AT); Rice (Os); Soybean (Glyma); Medicago trunculata (Medtr).



was identified in maize Lhcb4 [62]. This residue is not conserved in *Selaginella* Lhcb4 (D8RTB9) but present in *Arabidopsis* (pThr-109), rice (Thr-111), and moss (Thr-119) (Figure 4B). On the other hand, 2 consecutive phosphosites (Ser-57, 58) were detected in D8RTB9 and the equivalent Ser residues are only found in moss but not in *Arabidopsis* or rice. While both of them are located in basic motifs, Ser-58 may also represent a target for acidic casein kinase II.

The PSII core proteins PsbA, PsbD and PsbC are also known to undergo a strong and dynamic redox-regulated phosphorylation cycle [63-65]. STN8-dependent phosphorylation of PSII proteins is required for rapid turnover of photo-damaged PSII complexes and it is highly important during prolonged exposure of the photosynthetic apparatus to excess light [66]. As determined by its structure, STN8 kinase was reported to have a peculiar substrate specificity restricted to the very N-terminal Thr residue of PsbA, PsbD and PsbC [60]. For example, the phosphosite Thr-2 in *Arabidopsis* PsbD is phosphorylated by STN8 [42,58]. The same phosphorylation event is detected in *Selaginella* C7B2K2 (PsbD) while an equivalent Thr residue is found in rice (Figure 4B). On the other hand, while no N-terminal Thr phosphosites were identified in *Selaginella* C7B2K3 (PsbC), its Thr-346 phosphorylation is conserved in *Arabidopsis* PsbC [42,52] and equivalent Thr residues are found in rice and moss. This site may represent a substrate of acidic or basic motif recognizing kinases, indicating the possibility of cross-talk between kinases as suggested previously [60].

The oxygen-evolving complex (OEC) is consisted of PsbO, PsbP and PsbQ. PsbO stabilizes the manganese cluster which is the primary site of water splitting. Besides, PsbO regulates dephosphorylation and turnover of the PSII reaction center PsbA [67,68]. However, no phosphorylation events in PsbO have been reported previously in any plants. In *Selaginella* D8TBN9 (PsbO), we identified a unique Pro-directed Ser-219 phosphosite. The equivalent residues in other PsbO sequences examined are all Lys which is non-phosphorylatable (Figure 4B). PsbQ is required for PSII assembly, stability, and photoautotrophic growth under low light conditions [69]. The *Selaginella* PsbQ (D8S1M9) was found to be phosphorylated at the Ser-61 residue, which is a potential target of Pro-directed kinase. Equivalent pSer and Ser residues are found in *Arabidopsis* and moss PsbQ sequences, respectively (Figure 4B).

PsaC and PsaF are components of PSI which performs the light-induced electron transfer from plastocyanin or cytochrome c6 (Cyt_c) to ferredoxin. As a chloroplast-encoded PSI subunit, PsaC binds the two terminal electron acceptors (F_A and F_B). No phosphorylation was reported in PsaC previously in any plants. PsaC is extremely conserved among the four plant species examined

here with most of the residues identical (Additional file 6: Figure S2). Intriguingly, the phosphorylation event occurs at a unique residue (Ser-71) in *Selaginella* PsaC (C7B2J3). The equivalent residues in the other plant sequences are all non-phosphorylatable. The nuclear subunit PsaF provides a docking site for plastocyanin and Cyt_c on the luminal side of PSI. In *Arabidopsis*, PsaF was reported to be phosphorylated at Ser-94, Ser-95, Tyr-99, and Ser-103 [42,52]. Most of the equivalent residues in *Selaginella* PsaF (D8QPQ3) are conserved except for Ser-95. On the other hand, the Ser-184 phosphosite in D8QPQ3 is located in the very C-terminal region which is absent in *Arabidopsis* and rice. The same residue was identified in the moss PsaF sequence, suggesting that the Ser-184 phosphorylation event might have been lost during the evolution of flowering plants.

Overall, several phosphorylated residues in the *Selaginella* photosynthesis proteins are conserved with equivalent phosphorylation in *Arabidopsis* and/or phosphorylatable residues in most of the plants examined, including Lhcb1: Thr-44, Lhcb2: Thr-42 and 46, PsbD: Thr-2, PsbC: Thr-346, and psbQ: Ser-61. The phosphorylation of Thr-46 in Lhcb2 is first identified in *Selaginella* and the equivalent residues in other plant sequences are likely to be phosphorylated. We also identified unique phosphorylated residues within highly conserved regions in *Selaginella* PsbO (Ser-219) and PsaC (Ser-71). On the other hand, phosphorylation events with equivalent residues only in moss were detected in *Selaginella* Lhcb1, Lhcb4 and PsaF. These phosphosites are located in low-homology regions when compared with the *Arabidopsis* and rice sequences, implicating that they were lost in the flowering plants during evolution. It will be very interesting to investigate how the different unique phosphorylation events are involved in light reactions in *Selaginella*.

Conclusions

Our work generates the first large-scale atlas of phosphoproteins in *Selaginella* which occupies a unique position in the evolution of terrestrial plants. Combining PEG fractionation with IMAC enrichment, a total of 1593 unique phosphopeptides (1588 individual phosphosites) representing 851 unique phosphoproteins were retrieved. An overview of the *Selaginella* phosphoproteomics data revealed general features which are largely consistent with the dicot model *Arabidopsis*. Known plant phosphorylation Ser/Thr motifs were extracted from total and *Selaginella*-specific phosphopeptides, implicating the conservation of phosphorylation machineries during vascular plant evolution. In fact, 97 highly conserved phosphorylation events were identified among *Selaginella* and flowering plant homologs. In PSI proteins, we identified conserved residues which are potential targets of STN7 and STN8 kinases. On the other hand, several phosphosites unique to

Selaginella were detected in the highly conserved PSI and PSII proteins. Future research into functional roles of Selaginella-specific phosphorylation events in photosynthesis and other processes may offer insight into the molecular mechanisms leading to the distinct evolution of lycophytes.

Methods

Protein extraction and PEG fractionation

Two-gram aerial tissues of wild-growing *Selaginella moellendorffii* collected from the Victoria Peak in Hong Kong were ground to fine powder in liquid nitrogen. The powder was homogenized in 10 mL of ice-cold Mg/NP-40 extraction buffer containing 0.5 M Tris-HCl (pH 8.3), 20 mM MgCl₂, 2% v/v NP-40, 2% v/v β-mercaptoethanol, 1 mM phenylmethylsulfonyl fluoride and 1% w/v polyvinylpyrrolidone using the Tissue-Tearor (BioSpec) operated at maximum speed for 1 min on ice [70]. After centrifugation at 12000 × g for 15 min at 4°C, the supernatant was treated with 15% PEG-4000 and incubated on ice for 30 min, followed by centrifugation at 1500 × g for 10 min at 4°C. The pellet was washed sequentially with ice-cold 10% trichloroacetic acid/acetone, ice-cold 100% methanol containing 0.1 M ammonium acetate, and ice-cold 100% acetone. The supernatant was precipitated by adding four volumes of ice-cold acetone and then incubated at -20°C for 2 h. After centrifugation at 12000 × g for 5 min at 4°C, the pellet was rinsed as described above. For the plant debris left after the initial Mg/NP-40 extraction, residual protein was extracted by 4% SDS. After centrifugation, the supernatant was precipitated with ice-cold acetone, followed by sequential rinsing of the pellet.

Protein digestion and phosphopeptide enrichment

The pellets obtained from each of the above step were re-suspended in solution containing 0.2 M Tris-HCl (pH 8.0), 8 M urea and 4 mM CaCl₂. Dissolved protein samples were reduced with 10 mM dithiothreitol for 30 min at 56°C, and the alkylated with 40 mM iodoacetamide for 30 min at room temperature in the dark. Protein concentration was measured by the Bio-Rad Protein Assay kit. Afterwards, trypsin (Worthington) was added in a 1:50 (enzyme: protein) w/w ratio and the mixture incubated overnight at 37°C. Trypsinized peptides were loaded onto a 1 g Sep-Pak C18 column (Waters), washed twice with 10 mL 1% acetic acid, eluted with 7 mL 80% acetonitrile containing 0.1% acetic acid, dried under speed-vacuum, re-suspended in 400 μL 1% acetic acid, and then loaded onto a mini-column of 40 μL IMAC resin prepared as described previously [71]. The IMAC mini-column was rinsed twice with 40 μL wash buffer containing 25% v/v acetonitrile, 100 mM NaCl and 0.1% v/v acetic acid, then washed once each with 40 μL 1% v/v acetic acid and 20 μL double-distilled water, eluted with 120 μL 6% w/v NH₃.

H₂O, and dried under speed-vacuum. IMAC-enriched phosphopeptides derived from different PEG fractionated samples (Additional file 1: Figure S1) were subject to LC-MS/MS analysis.

RPLC-ESI-MS/MS detection

The Triple TOF 5600 mass spectrometer (AB SCIEX), a hybrid quadrupole TOF platform [72], was coupled with an Nano-LC system (Agilent) utilizing Nanospray III ion-source (AB SCIEX). Mobile phase A (2% ACN, 0.1% formic acid) and mobile phase B (98% ACN, 0.1% formic acid) were used to establish a 120 min gradient comprised of 80 min (5-30% B), 12 min (30-60% B), 6 min (60-90% B), 10 min (90% B), and 12 min (90-5% B). The flow rate was 300 nL/min. Peptides were separated on a fused silica capillary emitter (New Objective) packed in-house with 5 μm C18 resin (New Objective), and analyzed in positive ion mode by electrospray ionization. For information dependent acquisition, each survey scan was acquired in 250 ms followed by 20 product ion scans collected in 50 ms/per scan.

Database searching of MS/MS spectra

For proteome analysis, raw data from Triple TOF 5600 were searched with ProteinPilot software (version 4.0, AB SCIEX) against the Uniprot *Selaginella moellendorffii* complete proteome database (downloaded in April 2011, 33195 sequences) using following parameters: Sample Type (Identification), Cys Alkylation (Iodoacetamide), Digestion (Trypsin), Search Effort (Rapid). The false discovery rate (FDR) analysis was done by using the tool integrated in ProteinPilot. All data were filtered at 1% FDR.

For phosphoproteome analysis, raw data MS/MS (wiff files) were converted to .mgf files and searched with the Mascot (version 2.2, Matrix Science) software [73] against the Selaginella proteome database using following parameters: fixed modifications was set to carbamidomethylation on cysteine, variable modifications was set to oxidation of methionine and phosphorylation at serine, threonine and tyrosine, peptide and MS/MS fragment tolerances were set to 20 ppm and 0.2 Da respectively, trypsin was selected as digestion enzyme, and up to two missed cleavages were allowed. All .mgf files were merged into one file followed by database searching.

Post-search data processing and phosphosite localization

The Mascot search result was first loaded into Scaffold (version 3.0, Proteome Software) for further analysis. In order to screen phosphopeptides with high confidence, "Min Protein" (protein identification probability), "Min # Peptide" (the number of unique peptides on which a protein identification is based) and "Min Peptide" (peptide identification probability) were adjusted to 20%, 1 and 95% respectively [74,75]. Afterwards, the mzIdentML file

generated by Scaffold was loaded into Scaffold PTM (version 1.1, Proteome Software) to determine the localization probability of phosphosites [76].

Gene ontology annotations

Gene ontology (GO) annotations of all identified proteins and phosphoproteins in *Selaginella* categorized into 3 classifications (Cellular Component, Molecular Function and Biological Process) were batch-retrieved from the Protein Information Resource (<http://pir.georgetown.edu/pirwww/search/batch.shtml>).

Analysis of phosphorylation site conservation

The *Selaginella* phosphopeptides were clustered with different plant phosphopeptides retrieved from the Plant Protein Phosphorylation Database (P³DB; <http://www.p3db.org/>) using the CD-HIT web server [54] (<http://www.bioinformatics.org/cd-hit/>). All phosphopeptide sequences were combined into a single Fasta file for data upload. Default parameters were adopted together with a 60% similarity cutoff and a bandwidth of 5. Conservation of phosphorylation sites among different plant species were then identified by manual inspection of the sequence alignment in each cluster.

Phosphorylation motif analysis

Sequence was centered on each phosphosite and extended to 15 amino acids (± 7 residues). Phosphosites, which could not be extended because of N- or C-termini, were excluded from motif analysis. Only phosphosites with localization probability above 95% were used. General phosphorylation motif classes were assigned as defined previously [50]: P at +1 (Pro-directed); D/E at +1/+2 or +3 (Acidic), 5 or more D/E at +1 to +6 (Acidic); K/R at -3 (Basic), 2 or more K/R at -6 to -1 (Basic); otherwise (Others). Specific motifs were extracted from the data set by using motif-x algorithm (<http://motif-x.med.harvard.edu/motif-x.html>) [53]. The *Selaginella* proteome database in fasta format was retrieved (<http://www.phytozome.com/>) and uploaded as background. The significance threshold was set to 10^{-6} and the minimum number of motif occurrences was 20.

Additional files

Additional file 1: Figure S1. SDS-PAGE analysis of protein samples after PEG fractionation. Total protein (T) was obtained from the supernatant after extraction of tissues in Mg/NP-buffer. Residual protein (R) was obtained by extracting the plant debris in 4% SDS. The supernatant was subject to 15% PEG precipitation and the pellet (P) was resuspended in urea-containing buffer. Protein in the final supernatant (S) was precipitated by acetone and resuspended in urea-containing buffer. Each lane was loaded with 50 μ g of protein. Following Coomassie blue staining, the large subunit of RUBISCO (RBCL) can be visualized as a discrete band corresponding to a MW of ~55 kDa. Note the substantially reduced abundance of RBCL in the sample (S) after PEG precipitation.

The fractionated samples (S, P, and R) were trypsin-digested, followed by IMAC enrichment of phosphopeptides.

Additional file 2: Table S1. List of all identified *Selaginella* phosphopeptides.

Additional file 3: Table S2. List of 716 *Selaginella* phosphoproteins (localization probability $\geq 95\%$) with OrthoMCL Group information.

Additional file 4: Table S3. List of *Selaginella* phosphopeptides with the KXXS motif.

Additional file 5: Table S4. List of *Selaginella* phosphopeptides with conserved phosphosites in other plant phosphopeptides.

Additional file 6: Figure S2. Complete ClustalW sequence alignment of the nine identified *Selaginella* PSI proteins with Arabidopsis, rice and moss orthologous sequences. Phosphosites identified in this study and in Arabidopsis are highlighted in red and yellow, respectively. The rice and moss sequences are included as references although no phosphorylation information is available for these proteins.

Abbreviations

FDR: False discovery rate; GO: Gene ontology; IMAC: Immobilized metal affinity chromatography; P³DB: Plant protein phosphorylation database; PEG: Polyethylene glycol; PSI: Photosystem I; PSII: Photosystem II; PTMs: Post-translational modifications; RUBISCO: Ribulose-1,5-bisphosphate carboxylase/oxygenase.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

XC, WLC, and CL initially conceived of and designed the proteomics experiments. XC and WLC participated in all experimental procedures, data analysis, and manuscript preparation. FYZ was involved in data analysis. CL finalized the manuscript for submission. All authors read and approved the final manuscript.

Acknowledgements

This work is supported by the HKU Seed Funding Programme for Basic Research (201011159033) and HKU Small Project Funding (201209176039).

Author details

¹School of Biological Sciences, The University of Hong Kong, Pokfulam Hong Kong, China. ²Wuhan Institute of Biotechnology, Wuhan, Hubei, China.

Received: 31 December 2013 Accepted: 6 March 2014

Published: 17 March 2014

References

1. Kenrick P, Crane PR: The origin and early evolution of plants on land. *Nature* 1997, **389**:33–39.
2. Banks JA, Nishiyama T, Hasebe M, Bowman JL, Gribskov M, dePamphilis C, Albert VA, Aono N, Aoyama T, Ambrose BA, Ashton NW, Axtell MJ, Barker E, Barker MS, Bennetzen JL, Bonawitz ND, Chapple C, Cheng C, Correa LG, Dacre M, DeBarry J, Dreyer I, Elias M, Engstrom EM, Estelle M, Feng L, Finet C, Floyd SK, Frommer WB, Fujita T, et al: The *Selaginella* genome identifies genetic changes associated with the evolution of vascular plants. *Science* 2011, **332**:960–963.
3. Wang W, Tanurdzic M, Luo M, Sisneros N, Kim HR, Weng JK, Kudrna D, Mueller C, Arumuganathan K, Carlson J, Chapple C, de Pamphilis C, Mandoli D, Tomkins J, Wing RA, Banks JA: Construction of a bacterial artificial chromosome library from the spikemoss *Selaginella moellendorffii*: a new resource for plant comparative genomics. *BMC Plant Biol* 2005, **5**:10.
4. Banks JA: *Selaginella* and 400 million years of separation. *Annu Rev Plant Biol* 2009, **60**:223–238.
5. Yin C, Richter U, Borner T, Weihe A: Evolution of phage-type RNA polymerases in higher plants: characterization of the single phage-type RNA polymerase gene from *Selaginella moellendorffii*. *J Mol Evol* 2009, **68**:528–538.
6. Hedman H, Kallman T, Lagercrantz U: Early evolution of the MFT-like gene family in plants. *Plant Mol Biol* 2009, **70**:359–369.
7. Alvarez-Venegas R, Avramova Z: Evolution of the PWWP-domain encoding genes in the plant and animal lineages. *BMC Evol Biol* 2012, **12**:101.

8. Tegeder M, Ward JM: **Molecular evolution of plant AAP and LHT amino acid transporters.** *Front Plant Sci* 2012, **3**:21.
9. Gomez-Porras JL, Riano-Pachon DM, Benito B, Haro R, Sklodowski K, Rodriguez-Navarro A, Dreyer I: **Phylogenetic analysis of k(+) transporters in bryophytes, lycophytes, and flowering plants indicates a specialization of vascular plants.** *Front Plant Sci* 2012, **3**:167.
10. Pedersen CN, Axelsen KB, Harper JF, Palmgren MG: **Evolution of plant p-type ATPases.** *Front Plant Sci* 2012, **3**:31.
11. Kopriva S, Wiedemann G, Reski R: **Sulfate assimilation in basal land plants - what does genomic sequencing tell us?** *Plant Biol (Stuttg)* 2007, **9**:556-564.
12. Hirano K, Nakajima M, Asano K, Nishiyama T, Sakakibara H, Kojima M, Katoh E, Xiang H, Tanahashi T, Hasebe M, Banks JA, Ashikari M, Kitano H, Ueguchi-Tanaka M, Matsuo M: **The GID1-mediated gibberellin perception mechanism is conserved in the lycophyte Selaginella moellendorffii but not in the bryophyte Physcomitrella patens.** *Plant Cell* 2007, **19**:3058-3079.
13. Paponov IA, Teale W, Lang D, Paponov M, Reski R, Rensing SA, Palme K: **The evolution of nuclear auxin signalling.** *BMC Evol Biol* 2009, **9**:126.
14. Rychel AL, Peterson KM, Torii KU: **Plant twitter: ligands under 140 amino acids enforcing stomatal patterning.** *J Plant Res* 2010, **123**:275-280.
15. Eklund DM, Svensson EM, Kost B: **The Physcomitrella patens: a model to investigate the role of RAC/ROP GTPase signalling in tip growth.** *J Exp Bot* 2010, **61**:1917-1937.
16. Hanada K, Hase T, Toyoda T, Shinozaki K, Okamoto M: **Origin and evolution of genes related to ABA metabolism and its signaling pathways.** *J Plant Res* 2011, **124**:455-465.
17. Smith DR: **Unparalleled GC content in the plastid DNA of Selaginella.** *Plant Mol Biol* 2009, **71**:627-639.
18. Shakirov EV, Shippen AB: **Selaginella moellendorffii telomeres: conserved and unique features in an ancient land plant lineage.** *Front Plant Sci* 2012, **3**:161.
19. Chan AP, Melake-Berhan A, O'Brien K, Buckley S, Quan H, Chen D, Lewis M, Banks JA, Rabinowicz PD: **The highest-copy repeats are methylated in the small genome of the early divergent vascular plant Selaginella moellendorffii.** *BMC Genomics* 2008, **9**:282.
20. Axtell MJ, Snyder JA, Bartel DP: **Common functions for diverse small RNAs of land plants.** *Plant Cell* 2007, **19**:1750-1769.
21. Hecht J, Grewe F, Knoop V: **Extreme RNA editing in coding islands and abundant microsatellites in repeat sequences of Selaginella moellendorffii mitochondria: the root of frequent plant mtDNA recombination in early tracheophytes.** *Genome Biol Evol* 2011, **3**:344-358.
22. Novikova O, Smyshlyayev G, Blinov A: **Evolutionary genomics revealed interkingdom distribution of Tcn1-like chromodomain-containing Gypsy LTR retrotransposons among fungi and plants.** *BMC Genomics* 2010, **11**:231.
23. Wang YH, Long CL, Yang FM, Wang X, Sun QY, Wang HS, Shi YN, Tang GH: **Pyrrrolidinoindoline alkaloids from Selaginella moellendorffii.** *J Nat Prod* 2009, **72**:1151-1154.
24. Wu B, Wang J: **Phenolic compounds from Selaginella moellendorffii.** *Chem Biodivers* 2011, **8**:1735-1747.
25. Cao Y, Tan NH, Chen JJ, Zeng GZ, Ma YB, Wu YP, Yan H, Yang J, Lu LF, Wang Q: **Bioactive flavones and biflavones from Selaginella moellendorffii Hieron.** *Fitoterapia* 2010, **81**:253-258.
26. Wang HS, Sun L, Wang YH, Shi YN, Tang GH, Zhao FW, Niu HM, Long CL, Li L: **Carboxymethyl flavonoids and a monoterpene glucoside from Selaginella moellendorffii.** *Arch Pharm Res* 2011, **34**:1283-1288.
27. Wang X, Chen S, Zhang H, Shi L, Cao F, Guo L, Xie Y, Wang T, Yan X, Dai S: **Desiccation tolerance mechanism in resurrection fern-ally Selaginella tamariscina revealed by physiological and proteomic analysis.** *J Proteome Res* 2010, **9**:6561-6577.
28. Muir T: **Posttranslational modification of proteins: expanding nature's inventory: by Christopher T Walsh.** *Chem Bio Chem* 2006, **7**:1623-1624.
29. Hubbard MJ, Cohen P: **On target with a new mechanism for the regulation of protein phosphorylation.** *Trends Biochem Sci* 1993, **18**:172-177.
30. Zolnierowicz S, Bollen M: **Protein phosphorylation and protein phosphatases: De Panne, Belgium, September 19-24, 1999.** *EMBO J* 2000, **19**:483-488.
31. De la Fuente van Bentem S, Hirt H: **Using phosphoproteomics to reveal signalling dynamics in plants.** *Trends Plant Sci* 2007, **12**:404-411.
32. Initiative TAG: **Analysis of the genome sequence of the flowering plant Arabidopsis thaliana.** *Nature* 2000, **408**:796-815.
33. Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S: **The protein kinase complement of the human genome.** *Science* 2002, **298**:1912-1934.
34. Stone JM, Walker JC: **Plant protein kinase families and signal transduction.** *Plant Physiol* 1995, **108**:451-457.
35. Des Francs CC, Thiellement H, De Vienne D: **Analysis of leaf proteins by two-dimensional gel electrophoresis: protease action as exemplified by ribulose bisphosphate carboxylase/oxygenase degradation and procedure to avoid proteolysis during extraction.** *Plant Physiol* 1985, **78**:178-182.
36. Aryal UK, Krochko JE, Ross AR: **Identification of phosphoproteins in Arabidopsis thaliana leaves using polyethylene glycol fractionation, immobilized metal-ion affinity chromatography, two-dimensional gel electrophoresis and mass spectrometry.** *J Proteome Res* 2012, **11**:425-437.
37. Xi J, Wang X, Li S, Zhou X, Yue L, Fan J, Hao D: **Polyethylene glycol fractionation improved detection of low-abundant proteins by two-dimensional electrophoresis analysis of plant proteome.** *Phytochemistry* 2006, **67**:2341-2348.
38. Lee DG, Ahsan N, Lee SH, Kang KY, Lee JJ, Lee BH: **An approach to identify cold-induced low-abundant proteins in rice leaf.** *C R Biol* 2007, **330**:215-225.
39. Acquadro A, Falvo S, Mila S, Giuliano Albo A, Comino C, Moglia A, Lanteri S: **Proteomics in globe artichoke: protein extraction and sample complexity reduction by PEG fractionation.** *Electrophoresis* 2009, **30**:1594-1602.
40. Scanff P, Yvon M, Pelissier JP: **Immobilized Fe3+ affinity chromatographic isolation of phosphopeptides.** *J Chromatogr* 1991, **539**:425-432.
41. Benschop JJ, Mohammed S, O'Flaherty M, Heck AJ, Slijper M, Menke FL: **Quantitative phosphoproteomics of early elicitor signaling in Arabidopsis.** *Mol Cell Proteomics* 2007, **6**:1198-1214.
42. Reiland S, Messerli G, Baerenfaller K, Gerrits B, Endler A, Grossmann J, Gruissem W, Baginsky S: **Large-scale Arabidopsis phosphoproteome profiling reveals novel chloroplast kinase substrates and phosphorylation networks.** *Plant Physiol* 2009, **150**:889-903.
43. Nakagami H, Sugiyama N, Mochida K, Daudi A, Yoshida Y, Toyoda T, Tomita M, Ishihama Y, Shirasu K: **Large-scale comparative phosphoproteomics identifies conserved phosphorylation sites in plants.** *Plant Physiol* 2010, **153**:1161-1174.
44. Meyer LJ, Gao J, Xu D, Thelen JJ: **Phosphoproteomic analysis of seed maturation in Arabidopsis, rapeseed, and soybean.** *Plant Physiol* 2012, **159**:517-528.
45. Grimsrud PA, den Os D, Wenger CD, Swaney DL, Schwartz D, Sussman MR, Ane JM, Coon JJ: **Large-scale phosphoprotein analysis in Medicago truncatula roots provides insight into in vivo kinase activity in legumes.** *Plant Physiol* 2010, **152**:19-28.
46. Bi YD, Wang HX, Lu TC, Li XH, Shen Z, Chen YB, Wang BC: **Large-scale analysis of phosphorylated proteins in maize leaf.** *Planta* 2011, **233**:383-392.
47. Rudrabhatla P, Reddy MM, Rajasekharan R: **Genome-wide analysis and experimentation of plant serine/threonine/tyrosine-specific protein kinases.** *Plant Mol Biol* 2006, **60**:293-319.
48. Li L, Stoeckert CJ Jr, Roos DS: **OrthoMCL: identification of ortholog groups for eukaryotic genomes.** *Genome Res* 2003, **13**:2178-2189.
49. Chen F, Mackey AJ, Vermunt JK, Roos DS: **Assessing performance of orthology detection strategies applied to eukaryotic genomes.** *PLoS One* 2007, **2**:e383.
50. Villen J, Beausoleil SA, Gerber SA, Gygi SP: **Large-scale phosphorylation analysis of mouse liver.** *Proc Natl Acad Sci U S A* 2007, **104**:1488-1493.
51. Nuhse TS, Stensballe A, Jensen ON, Peck SC: **Phosphoproteomics of the Arabidopsis plasma membrane and a new phosphorylation site database.** *Plant Cell* 2004, **16**:2394-2405.
52. Sugiyama N, Nakagami H, Mochida K, Daudi A, Tomita M, Shirasu K, Ishihama Y: **Large-scale phosphorylation mapping reveals the extent of tyrosine phosphorylation in Arabidopsis.** *Mol Syst Biol* 2008, **4**:193.
53. Schwartz D, Gygi SP: **An iterative statistical approach to the identification of protein phosphorylation motifs from large-scale data sets.** *Nat Biotechnol* 2005, **23**:1391-1398.
54. Li W, Godzik A: **Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences.** *Bioinformatics* 2006, **22**:1658-1659.

55. Jansson S: **The light-harvesting chlorophyll a/b-binding proteins.** *Biochim Biophys Acta* 1994, **1184**:1–19.
56. Bellaïflore S, Barneche F, Peltier G, Rochaix JD: **State transitions and light adaptation require chloroplast thylakoid protein kinase STN7.** *Nature* 2005, **433**:892–895.
57. Bennett J: **Chloroplast phosphoproteins: phosphorylation of polypeptides of the light-harvesting chlorophyll protein complex.** *Eur J Biochem* 1979, **99**:133–137.
58. Vener AV, Harms A, Sussman MR, Vierstra RD: **Mass spectrometric resolution of reversible protein phosphorylation in photosynthetic membranes of *Arabidopsis thaliana*.** *J Biol Chem* 2001, **276**:6959–6966.
59. Tikkanen M, Piippo M, Suorsa M, Sirpio S, Mulo P, Vainonen J, Vener AV, Allahverdiyeva Y, Aro EM: **State transitions revisited—a buffering system for dynamic low light acclimation of *Arabidopsis*.** *Plant Mol Biol* 2006, **62**:779–793.
60. Vainonen JP, Hansson M, Vener AV: **STN8 protein kinase in *Arabidopsis thaliana* is specific in phosphorylation of photosystem II core proteins.** *J Biol Chem* 2005, **280**:33679–33686.
61. Bergantino E, Dainese P, Cerovic Z, Sechi S, Bassi R: **A post-translational modification of the photosystem II subunit CP29 protects maize from cold stress.** *J Biol Chem* 1995, **270**:8474–8481.
62. Bergantino E, Sandona D, Cugini D, Bassi R: **The photosystem II subunit CP29 can be phosphorylated in both C3 and C4 plants as suggested by sequence analysis.** *Plant Mol Biol* 1998, **36**:11–22.
63. Aro EM, Ohad I: **Redox regulation of thylakoid protein phosphorylation.** *Antioxid Redox Signal* 2003, **5**:55–67.
64. Vener AV: **Environmentally modulated phosphorylation and dynamics of proteins in photosynthetic membranes.** *Biochim Biophys Acta* 2007, **1767**:449–457.
65. Rochaix JD: **Role of thylakoid protein kinases in photosynthetic acclimation.** *FEBS Lett* 2007, **581**:2768–2775.
66. Tikkanen M, Nurmi M, Kangasjarvi S, Aro EM: **Core protein phosphorylation facilitates the repair of photodamaged photosystem II at high light.** *Biochim Biophys Acta* 2008, **1777**:1432–1437.
67. Yi X, McChargue M, Laborde S, Frankel LK, Bricker TM: **The manganese-stabilizing protein is required for photosystem II assembly/stability and photoautotrophy in higher plants.** *J Biol Chem* 2005, **280**:16170–16174.
68. Lundin B, Hansson M, Schoefs B, Vener AV, Spetea C: **The *Arabidopsis* PsbO2 protein regulates dephosphorylation and turnover of the photosystem II reaction centre D1 protein.** *Plant J* 2007, **49**:528–539.
69. Yi X, Hargett SR, Frankel LK, Bricker TM: **The PsbQ protein is required in *Arabidopsis* for photosystem II assembly/stability and photoautotrophy under low light conditions.** *J Biol Chem* 2006, **281**:26260–26267.
70. Kim ST, Cho KS, Jang YS, Kang KY: **Two-dimensional electrophoretic analysis of rice proteins by polyethylene glycol fractionation for protein arrays.** *Electrophoresis* 2001, **22**:2103–2109.
71. Chen X, Wu D, Zhao Y, Wong BH, Guo L: **Increasing phosphoproteome coverage and identification of phosphorylation motifs through combination of different HPLC fractionation methods.** *J Chromatogr B Analyt Technol Biomed Life Sci* 2011, **879**:25–34.
72. Andrews GL, Simons BL, Young JB, Hawkridge AM, Muddiman DC: **Performance characteristics of a new hybrid quadrupole time-of-flight tandem mass spectrometer (TripleTOF 5600).** *Anal Chem* 2011, **83**:5442–5446.
73. Perkins DN, Pappin DJ, Creasy DM, Cottrell JS: **Probability-based protein identification by searching sequence databases using mass spectrometry data.** *Electrophoresis* 1999, **20**:3551–3567.
74. Keller A, Nesvizhskii AI, Kolker E, Aebersold R: **Empirical statistical model to estimate the accuracy of peptide identifications made by MS/MS and database search.** *Anal Chem* 2002, **74**:5383–5392.
75. Nesvizhskii AI, Keller A, Kolker E, Aebersold R: **A statistical model for identifying proteins by tandem mass spectrometry.** *Anal Chem* 2003, **75**:4646–4658.
76. Beausoleil SA, Villen J, Gerber SA, Rush J, Gygi SP: **A probability-based approach for high-throughput protein phosphorylation analysis and site localization.** *Nat Biotechnol* 2006, **24**:1285–1292.

doi:10.1186/1477-5956-12-16

Cite this article as: Chen et al.: Phosphoproteomic analysis of the non-seed vascular plant model *Selaginella moellendorffii*. *Proteome Science* 2014 **12**:16.

Submit your next manuscript to BioMed Central and take full advantage of:

- **Convenient online submission**
- **Thorough peer review**
- **No space constraints or color figure charges**
- **Immediate publication on acceptance**
- **Inclusion in PubMed, CAS, Scopus and Google Scholar**
- **Research which is freely available for redistribution**

Submit your manuscript at
www.biomedcentral.com/submit

