





3. National Institute of Health. Osteoporosis prevention, diagnosis, and therapy. NIH Consensus Statement 2000. Available from: <http://consensus.nih.gov/2000/20000steoporosis111html.htm>. Accessed: September 2006.
4. Kanis JA, Melton LJ, III, Christiansen C, et al. The diagnosis of osteoporosis. *J Bone Miner Res* 1994;9:1137-41.
5. Van Daele PL, Burger H, Algra D, et al. Age-associated changes in ultrasound measurements of the calcaneus in men and women: the Rotterdam Study. *J Bone Miner Res* 1994;9:1751-7.
6. Massie A, Reid DM, Porter RW. Screening for osteoporosis: comparison between dual energy X-ray absorptiometry and broadband ultrasound attenuation in 1000 perimenopausal women. *Osteoporos Int* 1993;3:107-10.
7. Young H, Howey S, Purdie DW. Broadband ultrasound attenuation compared with dual-energy X-ray absorptiometry in screening for postmenopausal low bone density. *Osteoporos Int* 1993;3:160-4.
8. Langton CM, Ballard PA, Bennett DK, Purdie DW. A comparison of the sensitivity and specificity of calcaneal ultrasound measurements with clinical criteria for bone densitometry (DEXA) referral. *Clin Rheumatol* 1997;16:117-8.
9. Hans D, Dargent-Molina P, Schott AM, et al. Ultrasonographic heel measurements to predict hip fracture in elderly women: the EPIDOS prospective study. *Lancet* 1996;348:511-4.
10. Bauer DC, Gluer CC, Cauley JA, et al. Broadband ultrasound attenuation predicts fractures strongly and independently of densitometry in older women. A prospective study. Study of Osteoporotic Fractures Research Group. *Arch Intern Med* 1997;157:629-34.
11. Khaw, KT, Reeve, J, Luben, R, et al. Prediction of total and hip fracture risk in men and women by quantitative ultrasound of the calcaneus: EPIC-Norfolk prospective population study. *Lancet* 2004;363:197.
12. Gluer, CC, Eastell, R, Reid, DM, et al. Association of five quantitative ultrasound devices and bone densitometry with osteoporotic vertebral fractures in a population-based sample: the OPUS study. *J Bone Miner Res* 2004;19:782.
13. Hodson J, Marsh J. Quantitative ultrasound and risk factor enquiry as predictors of postmenopausal osteoporosis: comparative study in primary care. *BMJ* 2003;326:1250-1.
14. Stewart A, Reid DM. Quantitative ultrasound or clinical risk factors: which best identifies women at risk of osteoporosis? *Br J Radiol* 2000;73:165-71.
15. Diez-Perez, A, Marin, F, Vila, J, et al. Evaluation of calcaneal quantitative ultrasound in a primary care setting as a screening tool for osteoporosis in postmenopausal women. *J Clin Densitom* 2003;6:237.