

Phloroglucinol Attenuates Free Radical-induced Oxidative Stress

Mi Jung So and Eun Ju Cho

Department of Food Science and Nutrition, and Research Institute of Ecology for the Elderly, Pusan National University, Busan 609-735, Korea

ABSTRACT: The protective role of phloroglucinol against oxidative stress and stress-induced premature senescence (SIPS) was investigated *in vitro* and in cell culture. Phloroglucinol had strong and concentration-dependent radical scavenging effects against nitric oxide (NO), superoxide anions (O_2^-), and hydroxyl radicals. In this study, free radical generators were used to induce oxidative stress in LLC-PK1 renal epithelial cells. Treatment with phloroglucinol attenuated the oxidative stress induced by peroxy radicals, NO, O_2^- , and peroxyxynitrite. Phloroglucinol also increased cell viability and decreased lipid peroxidation in a concentration-dependent manner. WI-38 human diploid fibroblast cells were used to investigate the protective effect of phloroglucinol against hydrogen peroxide (H_2O_2)-induced SIPS. Phloroglucinol treatment attenuated H_2O_2 -induced SIPS by increasing cell viability and inhibited lipid peroxidation, suggesting that treatment with phloroglucinol should delay the aging process. The present study supports the promising role of phloroglucinol as an antioxidative agent against free radical-induced oxidative stress and SIPS.

Keywords: phloroglucinol, oxidative stress, nitric oxide, LLC-PK1 cells, WI-38 cells

INTRODUCTION

Oxidative stress and nitrosative stress are caused by the uncontrolled production of reactive oxygen species (ROS) and reactive nitrogen species (RNS) (1-3). Excessive production of ROS and RNS can damage cellular lipids, proteins, and DNA, thus inhibiting their normal function. Therefore, oxidative stress and nitrosative stress have been implicated in aging and in a number of human diseases, such as cardiovascular disease, cancer, neurological disorders, and diabetes (4-8). Antioxidants, which prevent free radical damage, have attracted much attention and there has been a marked effort to identify safe and effective therapeutic agents for the treatment of oxidative stress-related diseases. Compelling evidence indicates that increased consumption of dietary antioxidants may improve quality of life by delaying the onset and reducing the risk of degenerative diseases (9-11).

Marine brown algae accumulate a variety of phloroglucinol-based polyphenols in the form of phlorotannins, which consist of phloroglucinol units linked to each other in a variety of ways. Phlorotannins can be found in a variety of marine organisms, especially brown and red

algae (12). Several studies have demonstrated that phloroglucinol has broad therapeutic effects, including anti-diabetes (13), antioxidative (14), and anti-cancer (15) effects. Previous work indicates that phloroglucinol significantly decreases the level of radiation-induced intracellular ROS and damage to cellular components such as lipids, DNA, and proteins (16). In addition, phloroglucinol has been reported to reduce ROS levels and increase antioxidant defenses in human HepG2 cells (17). However, the protective effect of phloroglucinol against peroxy radical- and nitric oxide (NO)-induced oxidative stress in LLC-PK1 cells, which are susceptible to oxidative stress, has not been demonstrated.

Oxidative stress leads to premature senescence in cellular systems, but the effect of phloroglucinol on stress-induced premature senescence (SIPS) in human fibroblast WI-38 cells has not been reported. In the present study, we investigated the protective effect of phloroglucinol against oxidative stress induced by NO and peroxy radicals. Furthermore, SIPS was induced in human fibroblast WI-38 cells by treatment with hydrogen peroxide (H_2O_2), and then the protective effect of phloroglucinol against SIPS was investigated.

Received 14 May 2014; Accepted 22 July 2014; Published online 30 September 2014

Correspondence to Eun Ju Cho, Tel: +82-51-510-2837, E-mail: ejcho@pusan.ac.kr

Copyright © 2014 by The Korean Society of Food Science and Nutrition. All rights Reserved.

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

MATERIALS AND METHODS

Materials

Phloroglucinol, 3-morpholinosydnonimine (SIN-1), pyrogallol, and 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H tetrazolium bromide (MTT) were purchased from Sigma Chemical Co. (St. Louis, MO, USA). 2,2'-Azobis(2-amidinopropane) dihydrochloride (AAPH) and sodium nitroprusside (SNP) were obtained from Wako Pure Chemical Industries, Ltd. (Osaka, Japan). The LLC-PK1 porcine renal epithelial cells and WI-38 human embryonic lung-derived diploid fibroblasts were obtained from ATCC (Manassas, VA, USA). Dulbecco's modified eagle medium/nutrient mixture F-12 (DMEM/F-12), basal medium eagle (BME), and fetal bovine serum (FBS) were purchased from Invitrogen Co. (Grand Island, NY, USA).

NO scavenging activity

NO was generated from SNP and measured by the Griess reaction (18) according to the method of Sreejayan and Rao (19). SNP in phosphate buffered saline was mixed with different concentrations of samples and incubated at 25°C for 150 min. The amount of NO produced by SNP was determined by measuring nitrite accumulation with a microplate assay method based on the Griess reaction.

Superoxide anion (O_2^-) scavenging activity

O_2^- levels were measured following the method described by Ewing and Janero (20). Briefly, phloroglucinol was added to microplate wells containing 200 μ L of freshly prepared 0.125 mM EDTA, 62 μ M nitro blue tetrazolium (NBT), and 98 μ M NADH in 50 mM phosphate buffer (pH 7.4). The reaction was initiated with the addition of 25 μ L of freshly prepared 33 μ M 5-methylphenazinium methyl sulfate in 50 mM phosphate buffer (pH 7.4). A microplate reader (SpectraMax 340PC, Molecular Devices, Sunnyvale, CA, USA) was used to monitor NBT reduction by continuously measuring the absorbance (540 nm) of the solution over a 5 min period.

Hydroxyl radical (\cdot OH) scavenging activity

The reaction mixture contained 0.45 mL of 0.2 M sodium phosphate buffer (pH 7.0), 0.15 mL of 10 mM 2-deoxyribose, 0.15 mL of 10 mM $FeSO_4$ -EDTA, 0.15 mL of 10 mM H_2O_2 , 0.525 mL of H_2O , and 0.075 mL of sample solution. The reaction was started by the addition of H_2O_2 . After incubation at 37°C for 4 h, the reaction was stopped by adding 0.75 mL of 2.8% trichloroacetic acid and 0.75 mL of 1.0% 2-thiobarbituric acid in 50 mL of 0.05 N NaOH. The solution was boiled for 10 min and then cooled in water. The absorbance of the resulting solution was measured at 520 nm. The \cdot OH scavenging

activity of each sample was defined as the inhibition rate of 2-deoxyribose oxidation by \cdot OH (21).

Cell culture

Commercially available LLC-PK1 cells were maintained at 37°C in a humidified atmosphere of 5% CO_2 in culture plates containing 5% FBS-supplemented DMEM/F-12 medium. The cells were subcultured weekly with 0.05% trypsin-EDTA in calcium- and magnesium-free phosphate buffer. The WI-38 cells were cultured in BME supplemented with 10% FBS, 100 U/mL penicillin, and 100 μ g/mL streptomycin in a humidified incubator at 37°C and 5% CO_2 . The cells were subcultured with 0.05% trypsin-EDTA in PBS. Cells at early passages (i.e., between passage 26 and passage 33) were used for all experiments.

Treatment with radical generators

After reaching confluence, LLC-PK1 cells were seeded into 96-well culture plates at a cell density of 1×10^4 cells/mL. Two hours later, cells were treated with 1.0 mM AAPH, 1.2 mM pyrogallol, 1.2 mM SNP, or 1.0 mM SIN-1 for 24 h. After the free radical generator treatments, cells were treated with various concentrations of phloroglucinol for 24 h (22-23). The WI-38 cells were seeded into 96-well plates at a cell density of 1×10^4 cells/mL and then treated with 50 μ M of H_2O_2 for 60 min to induce SIPS. After SIPS induction, WI-38 cells were treated with various concentrations of phloroglucinol for 24 h.

MTT assay

Cell viability was determined with a MTT colorimetric assay (24). Fifty microliters of MTT solution (1 mg/mL) were added to the each well. After incubation for 4 h at 37°C, the MTT solution was removed from the wells. The formazan crystals present in the renal cells were solubilized with 100 μ L of dimethylsulfoxide. A microplate reader (SpectraMax 340PC) was used to read the absorbance of each well at 540 nm.

Thiobarbituric acid reactive substances (TBARS)

The level of lipid peroxidant released from cultured cells was estimated as TBARS according to the methods of Yagi (25) and Yokode et al. (26) with a slight modification. One aliquot of medium was mixed with 1.5 mL of 0.67% thiobarbituric acid aqueous solution and 1.5 mL of 20% trichloroacetic acid. The resulting mixture was boiled at 95~100°C for 45 min and then cooled and shaken vigorously with 3.0 mL of *n*-butanol. After centrifuging at 4,000 g for 10 min, a fluorescence spectrophotometer was used to measure the fluorescence of the *n*-butanol layer (excitation wavelength: 515 nm, emission wavelength: 553 nm).

Statistical analysis

The results for each group are expressed as mean±standard deviation. SAS software (SAS Institute Inc., Cary, NC, USA) was used to analyze the differences between the control and sample treated groups. Differences between groups were considered significant when the *P*-value was less than 0.05.

RESULTS

In vitro radical scavenging activity

Table 1 shows the *in vitro* radical scavenging activities of phloroglucinol. Phloroglucinol scavenged NO, O₂⁻ and ·OH radicals in a dose-dependent manner. The radical scavenging effect of phloroglucinol was strongest against NO. The 250 µg/mL, 500 µg/mL, and 1,000 µg/mL phloroglucinol concentrations were associated with scavenging effects of 59.3%, 64.4%, and 79.7%, respectively. In addition, the 1,000 µg/mL phloroglucinol concentration was associated with 55.5% and 59.1% scavenging effects against O₂⁻ and ·OH, respectively.

Protective activity against radical-induced oxidative stress

The effect of phloroglucinol on the viability and lipid peroxidation of LLC-PK1 renal epithelial cells treated with AAPH is shown in Table 2. Treatment with 1 mM

Table 1. *In vitro* radical scavenging effects of phloroglucinol

| Concentration (µg/mL) | Scavenging effect (%) | | |
|-----------------------|-----------------------|-----------------------------|-----------------------|
| | NO | O ₂ ⁻ | ·OH |
| 50 | 25.4±7.4 ^c | No effect | 13.7±0.4 ^e |
| 100 | 59.3±2.9 ^b | 1.5±0.3 ^d | 24.5±0.4 ^d |
| 250 | 59.3±2.9 ^b | 15.4±0.5 ^c | 38.5±0.4 ^c |
| 500 | 64.4±2.9 ^b | 36.1±0.6 ^b | 50.4±0.2 ^b |
| 1000 | 79.7±0.0 ^a | 55.5±0.3 ^a | 59.1±0.3 ^a |

Values are mean±SD.

^{a-e}Means with different superscripts within each column are significantly different (*P*<0.05) by Duncan's multiple range test.

Table 2. Protective effect of phloroglucinol against AAPH-induced oxidative stress in LLC-PK1 cells

| Concentration (µg/mL) | Cell viability (%) | MDA (nmol/mg protein) |
|--------------------------|--------------------------|------------------------|
| AAPH non-treated control | 100.00±0.66 ^a | 0.19±0.01 ^e |
| AAPH-treated control | 16.40±1.32 ^f | 0.80±0.01 ^a |
| 10 | 55.10±1.33 ^e | 0.55±0.03 ^b |
| 25 | 59.75±1.33 ^d | 0.43±0.01 ^c |
| 50 | 71.70±2.66 ^c | 0.30±0.04 ^d |
| 100 | 95.60±2.66 ^b | 0.21±0.01 ^e |

Values are mean±SD.

^{a-f}Means with different superscripts within each column are significantly different (*P*<0.05) by Duncan's multiple range test. Two hours after seeding, LLC-PK1 cells were treated with 1.0 mM AAPH for 24 h and then 10, 25, 50, or 100 µg/mL phloroglucinol for 24 h.

AAPH for 24 h reduced the viability of LLC-PK1 cells to 16.40%. However, phloroglucinol treatment exerted a protective effect against AAPH-induced cellular damage. Treatment with 25 µg/mL, 50 µg/mL, and 100 µg/mL phloroglucinol elevated cell viability from 16.40% to 59.75%, 71.70%, and 95.6%, respectively. Treatment with AAPH increased lipid peroxidation in LLC-PK1 renal tubular epithelial cells, resulting in a TBARS concentration of 0.80 nmol/mg protein. However, treatment with phloroglucinol significantly decreased the formation of TBARS in a concentration-dependent manner. The addition of 25 µg/mL and 100 µg/mL phloroglucinol to AAPH treated wells decreased lipid peroxidation, leading to TBARS concentrations of 0.43 nmol/mg protein and 0.21 nmol/mg protein, respectively.

Pyrogallol was used to evaluate the protective activity of phloroglucinol against O₂⁻. As shown in Table 3, O₂⁻ generated by pyrogallol decreased cell viability from 100% to 31.90%, and elevated malondialdehyde (MDA) formation from 0.21 nmol/mg protein to 0.76 nmol/mg protein. However, treatment with a 100 µg/mL phloroglucinol concentration increased cell viability to 79.1% and decreased MDA levels to 0.23 nmol/mg protein.

Table 4 shows the protective effect of phloroglucinol

Table 3. Protective effect of phloroglucinol against pyrogallol-induced oxidative stress in LLC-PK1 cells

| Concentration (µg/mL) | Cell viability (%) | MDA (nmol/mg protein) |
|--------------------------------|--------------------------|------------------------|
| Pyrogallol non-treated control | 100.00±0.65 ^a | 0.21±0.01 ^e |
| Pyrogallol-treated control | 31.90±0.66 ^f | 0.76±0.03 ^a |
| 10 | 58.66±2.63 ^e | 0.58±0.02 ^b |
| 25 | 67.23±2.64 ^d | 0.49±0.01 ^c |
| 50 | 72.50±3.29 ^c | 0.32±0.02 ^d |
| 100 | 79.10±1.32 ^b | 0.23±0.02 ^e |

Values are mean±SD.

^{a-f}Means with different superscripts within each column are significantly different (*P*<0.05) by Duncan's multiple range test. Two hours after seeding, LLC-PK1 cells were treated with 1.2 mM pyrogallol for 24 h and then 10, 25, 50, or 100 µg/mL phloroglucinol for 24 h.

Table 4. Protective effect of phloroglucinol against SNP-induced oxidative stress in LLC-PK1 cells

| Concentration (µg/mL) | Cell viability (%) | MDA (nmol/mg protein) |
|-------------------------|--------------------------|------------------------|
| SNP non-treated control | 100.00±0.67 ^a | 0.22±0.01 ^f |
| SNP-treated control | 26.30±0.67 ^f | 0.80±0.01 ^a |
| 10 | 52.06±1.34 ^e | 0.68±0.01 ^b |
| 25 | 61.41±2.67 ^d | 0.57±0.01 ^c |
| 50 | 69.42±1.33 ^c | 0.40±0.01 ^d |
| 100 | 80.10±2.67 ^b | 0.32±0.01 ^e |

Values are mean±SD.

^{a-f}Means with different superscripts within each column are significantly different (*P*<0.05) by Duncan's multiple range test. Two hours after seeding, LLC-PK1 cells were treated with 1.2 mM SNP for 24 h and then 10, 25, 50, or 100 µg/mL phloroglucinol for 24 h.

Table 5. Protective effect of phloroglucinol against SIN-1-induced oxidative stress in LLC-PK1 cells

| Concentration (µg/mL) | Cell viability (%) | MDA (nmol/mg protein) |
|---------------------------|--------------------------|------------------------|
| SIN-1 non-treated control | 100.00±0.66 ^a | 0.19±0.01 ^e |
| SIN-1-treated control | 21.40±1.32 ^f | 0.81±0.01 ^a |
| 10 | 52.80±1.32 ^e | 0.70±0.01 ^b |
| 25 | 66.00±1.32 ^d | 0.56±0.01 ^c |
| 50 | 71.28±1.98 ^c | 0.34±0.02 ^d |
| 100 | 85.80±1.98 ^b | 0.20±0.01 ^e |

Values are mean±SD.

^{a-f}Means with different superscripts within each column are significantly different ($P<0.05$) by Duncan's multiple range test. Two hours after seeding, LLC-PK1 cells were treated with 1.0 mM SIN-1 for 24 h and then 10, 25, 50, or 100 µg/mL phloroglucinol for 24 h.

against NO-induced oxidative stress. The viability of cells that had been treated with SNP was markedly lower than that of non-treated cells. However, phloroglucinol increased cell viability in a dose-dependent manner. At a 100 µg/mL phloroglucinol concentration, cell viability was elevated to 80.1%. In addition, SNP treatment increased lipid peroxidation from 0.22 nmol MDA/mg protein to 0.80 nmol MDA/mg protein, while treatment with 100 µg/mL phloroglucinol decreased lipid peroxidation to 0.32 nmol MDA/mg protein.

Table 5 shows the protective activity of phloroglucinol against peroxynitrite (ONOO⁻). Treatment with SIN-1 decreased cell viability to 21.4%, whereas treatment with 25 µg/mL and 100 µg/mL concentrations of phloroglucinol increased cell viability to 66.00% and 85.80%, respectively. Treatment with SIN-1 significantly increased MDA production by LLC-PK1 cells. However, treatment with 25 µg/mL and 100 µg/mL concentrations of phloroglucinol decreased MDA levels from 0.81 nmol/mg protein to 0.56 nmol/mg protein and 0.20 nmol/mg protein, respectively.

Protective effect of phloroglucinol against SIPS

Table 6 shows the effects of phloroglucinol on WI-38 cell viability and MDA production during H₂O₂-induced premature senescence. Treatment of WI-38 cells with 50 µM H₂O₂ for 60 min significantly decreased cell viability to 64.60%. However, treatment with phloroglucinol was associated with a significant dose-dependent increase in cell viability. Treatment with 25 µg/mL and 100 µg/mL phloroglucinol concentrations increased cell viability to 79.98% and 91.65%, respectively. In addition, MDA production by H₂O₂-treated, prematurely senescent cells was markedly greater than that by young, control WI-38 cells (0.45 nmol/mg protein vs. 1.02 nmol/mg protein). Treatment of WI-38 cells with phloroglucinol during SIPS is associated with a concentration-dependent decrease in MDA production; the MDA levels of prematurely-senescent WI-38 cells treated with 25 µg/mL

Table 6. Protective effect of phloroglucinol against H₂O₂-induced SIPS in WI-38 fibroblast cells

| Concentration (µg/mL) | Cell viability (%) | MDA (nmol/mg protein) |
|------------------------------|--------------------------|------------------------|
| Young control | 100.00±2.33 ^a | 0.45±0.02 ^e |
| Premature senescence control | 64.60±1.45 ^f | 1.02±0.01 ^a |
| 10 | 67.68±1.54 ^e | 0.65±0.02 ^b |
| 25 | 79.98±1.54 ^d | 0.57±0.01 ^c |
| 50 | 85.36±0.77 ^c | 0.48±0.01 ^d |
| 100 | 91.65±1.56 ^b | 0.42±0.02 ^f |

Values are mean±SD.

^{a-f}Means with different superscripts within each column are significantly different ($P<0.05$) by Duncan's multiple range test. Two hours after seeding, WI-38 cells were treated with 50 µM H₂O₂ for 60 min and then treated with 10, 25, 50, or 100 µg/mL phloroglucinol for 24 h.

and 100 µg/mL concentrations of phloroglucinol were 0.57 nmol/mg protein and 0.42 nmol/mg, respectively.

DISCUSSION

Phloroglucinol is the monomeric building unit of phlorotannins, which are phenolic compounds that can be found in brown algae. The free form of phloroglucinol can be found in some members of the order Fucales, including *Cystoseira discors* [$>0.5\%$ dry weight (DW)] (27), *Cystoseira tamariscifolia* ($>0.5\%$ DW) (27), and *Ecklonia cava* (*Laminariaceae*) (0.016% DW) (28).

Several studies have reported on the biological activities of phloroglucinol, which can be isolated from brown algae or purchased from Sigma Chemical Co. Kang et al. (28) demonstrated that phloroglucinol from *Ecklonia cava* has a ROS scavenging effect and protects Chinese hamster lung fibroblast (V79-4) cells against oxidative damage by enhancing cellular catalase activity and regulating extracellular signal-regulated kinase. In addition, Ishii et al. (29) demonstrated that phloroglucinol and phloroglucinol derivatives from *Mallotus japonicus* inhibit NO production by reducing inducible NO synthase protein induction and inhibiting enzyme activity in lipopolysaccharide- and interferon- γ -stimulated RAW 264.7 cells. Quéguineur et al. (17) reported that phloroglucinol from Sigma Chemical Co. has antioxidant activities in non-biological assays and alleviates oxidative stress in human HepG2 cells. Phloroglucinol has a direct scavenging effect against free radicals, increases antioxidant enzyme defenses, and protects against lipid peroxidation in cultivated liver cells. However, the protective role of phloroglucinol against free radical overproduction by LLC-PK1 renal epithelial cells, which are susceptible to oxidative stress, has not been studied. In addition, the effect of phloroglucinol on SIPS in human fibroblast WI-38 cells has not been reported. In the present study, we investigated the protective effects of phloroglucinol

against oxidative stress and SIPS *in vitro* and in cell culture.

Since the overproduction of free radicals results in a variety of pathological conditions, including cardiovascular disease, cancer, neurological disorders, diabetes, and aging. Antioxidants that prevent free radical damage have attracted much attention (30,31). Several studies have demonstrated that brown seaweeds play a protective role against pathological phenomena. Because phloroglucinol is one of the main active components of brown seaweeds (12-15), the present investigation was focused on the protective activity of phloroglucinol against free radical-induced oxidative stress *in vitro* and in cell culture.

NO is a free radical with a single unpaired electron. Overproduction of NO causes tissue damage and contributes to pathological conditions (30,31). In addition, reaction of O_2^- with NO leads to the formation of $ONOO^-$, which is far more reactive and toxic than its precursors (32,33). This reaction is extremely rapid, and can generate the most toxic radical, $\cdot OH$ (34). $\cdot OH$ is the most reactive radical, and it can react with the unsaturated fatty acids of membrane phospholipids to generate free radicals, which in turn react quickly with oxygen to form peroxides. These peroxides then act as free radicals, initiating an autocatalytic chain reaction that results in further loss of unsaturated fatty acids and extensive membrane damage. In this study, we found that phloroglucinol had strong and concentration-dependent radical scavenging effects against NO, O_2^- and $\cdot OH$. The present results provide evidence to indicate that phloroglucinol plays a protective role against oxidative stress by scavenging free radicals.

The use of LLC-PK1 cells, which are vulnerable to the oxidative stress induced by free radicals, as a cellular model of oxidative stress is well established and is useful when searching for agents that can provide effective protection from free radicals (22,23). Therefore, in the present study, cellular oxidative stress was induced in LLC-PK1 cells by free radical generators such as AAPH, SNP, pyrogallol, and SIN-1.

AAPH-intoxication experiments are useful for evaluating the biological activities of antioxidants. AAPH generates free radicals at a constant and well-defined rate by its thermal decomposition without biotransformation (35). The free radicals generated from AAPH rapidly react with oxygen molecules to yield peroxy radicals. The lipid peroxy radicals attack other lipid molecules to form lipid hydroperoxides and new lipid radicals. This reaction takes place repeatedly, with resultant attacks upon various biological molecules, and induces physicochemical alterations and cellular damage (36). In the present study, we used AAPH to produce peroxy radicals. The present study showed that AAPH leads to a decline

in cell viability and an elevation in lipid peroxidation in LLC-PK1 renal epithelial cells. However, treatment with phloroglucinol exerted a protective effect against AAPH-induced oxidative damage to LLC-PK1 cells, resulting in a dose-dependent increase in cell viability and decrease in lipid peroxidation.

The protective activity of phloroglucinol against NO-induced oxidative stress was also evaluated. NO and O_2^- generated by SNP and pyrogallol, respectively, significantly decreased cell viability and elevated lipid peroxidation compared to non-treated cells. However, phloroglucinol had significant protective effects against NO- and O_2^- -induced cell loss and lipid peroxidation. Peroxynitrite, which is derived from the reaction of NO with O_2^- , is a strong oxidant and nitrating agent that can lead to the production of the highly toxic $\cdot OH$ radical (32). SIN-1 simultaneously generates both NO and O_2^- , which then rapidly combine to form $ONOO^-$ (23). Compared with non-treated cells, treatment with SIN-1 led to a significant decrease in cell viability and increase in lipid peroxidation. However, phloroglucinol attenuated this oxidative stress by increasing cell viability and decreasing lipid peroxidation. These results indicate that phloroglucinol plays a protective role against NO-induced oxidative stress.

Oxidative reactions that occur in organisms generate free radicals, which cause multiple lesions in macromolecules and lead to aging (37,38). In addition, a decline in antioxidant defenses and progressive increase in oxidative stress is commonly associated with aging and age-related degenerative diseases. Human diploid fibroblasts (HDFs) are a typical model for studying the process of aging *in vitro*. Various oxidative stresses, such as treatment with H_2O_2 , have been used to study the onset of cellular senescence (39,40). Exposure of normal HDFs to H_2O_2 is widely used as a model to study cellular aging. The exposure of HDFs to a sublethal concentration of H_2O_2 induces permanent cell cycle arrest and phenotypic changes that mimic those seen in replicatively senescent cells (41). Therefore, an aging model of H_2O_2 -induced SIPS was used to investigate the protective effect of phloroglucinol against the oxidative stress induced by H_2O_2 . In the present study, H_2O_2 -induced SIPS led to a loss of cell viability, suggesting that H_2O_2 treatment led to premature replicative senescence (41,42). Treatment with phloroglucinol improved cell viability by protecting against H_2O_2 -induced oxidative damage. Intracellular oxidative stress has harmful effects on biomolecules such as lipids, proteins, and DNA and eventually leads to cellular senescence.

Previously, Morliere and Santus (43) reported that the exposure of cultured human skin fibroblasts to *tert*-butyl hydroperoxide, ultraviolet-A, and H_2O_2 results in the release of high levels of TBARS, an index of lipid pero-

oxidation. MDA is used as a biological marker of lipid peroxidation. In the present study, MDA levels were significantly increased by exposure to H₂O₂. In contrast, treatment with phloroglucinol inhibited lipid peroxidation in WI-38 cells under SIPS, suggesting that phloroglucinol may protect against oxidative stress. Although further study into the mechanism responsible for this protective effect has to be carried out, the present study supports the promising role of phloroglucinol as an antioxidative agent against free radical-induced oxidative stress and SIPS.

ACKNOWLEDGEMENTS

This work was supported by a 2-Year Research Grant of Pusan National University.

AUTHOR DISCLOSURE STATEMENT

The authors declare no conflict of interest.

REFERENCES

- Kovacic P, Jacintho JD. 2001. Mechanisms of carcinogenesis: focus on oxidative stress and electron transfer. *Curr Med Chem* 8: 773-796.
- Valko M, Morris H, Mazúr M, Raptá P, Bilton RF. 2001. Oxygen free radical generating mechanisms in the colon: do the semiquinones of vitamin K play a role in the aetiology of colon cancer? *Biochim Biophys Acta* 1527: 161-166.
- Ridnour LA, Isenberg JS, Espey MG, Thomas DD, Roberts DD, Wink DA. 2005. Nitric oxide regulates angiogenesis through a functional switch involving thrombospondin-1. *Proc Natl Acad Sci USA* 102: 13147-13152.
- Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. 2006. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact* 160: 1-40.
- Dhalla NS, Temsah RM, Netticadan T. 2000. Role of oxidative stress in cardiovascular diseases. *J Hypertens* 18: 655-673.
- Sayre LM, Smith MA, Perry G. 2001. Chemistry and biochemistry of oxidative stress in neurodegenerative disease. *Curr Med Chem* 8: 721-738.
- Jenner P. 2003. Oxidative stress in Parkinson's disease. *Ann Neurol* 53: S26-S36.
- Dalle-Donne I, Rossi R, Colombo R, Giustarini D, Milzani A. 2006. Biomarkers of oxidative damage in human disease. *Clin Chem* 52: 601-623.
- Ames BN, Shigenaga MK, Hagen TM. 1993. Oxidants, antioxidants, and the degenerative diseases of aging. *Proc Natl Acad Sci USA* 90: 7915-7922.
- Steinmetz KA, Potter JD. 1996. Vegetables, fruit, and cancer prevention: a review. *J Am Diet Assoc* 96: 1027-1039.
- Meydani M, Lipman RD, Han SN, Wu D, Beharka A, Martin KR, Bronson R, Cao G, Smith D, Meydani SN. 1998. The effect of long-term dietary supplementation with antioxidants. *Ann NY Acad Sci* 854: 352-360.
- Balboa EM, Conde E, Moure A, Falqué E, Domínguez H. 2013. *In vitro* antioxidant properties of crude extracts and compounds from brown algae. *Food Chem* 138: 1764-1785.
- Dellabella M, Milanese G, Muzzonigro G. 2005. Randomized trial of the efficacy of tamsulosin, nifedipine and phloroglucinol in medical expulsive therapy for distal ureteral calculi. *J Urol* 174: 167-172.
- Rosa A, Deiana M, Atzeri A, Corona G, Incani A, Melis MP, Appendino G, Dessì MA. 2007. Evaluation of the antioxidant and cytotoxic activity of arzanol, a prenylated α -pyrone-phloroglucinol etherodimer from *Helichrysum italicum* subsp. *microphyllum*. *Chem Biol Interact* 165: 117-126.
- Hashida W, Tanaka N, Kashiwada Y, Sekiya M, Ikeshiro Y, Takaishi Y. 2008. Tomoeones A-H, cytotoxic phloroglucinol derivatives from *Hypericum ascyron*. *Phytochemistry* 69: 2225-2230.
- Kang KA, Zhang R, Chae S, Lee SJ, Kim J, Kim J, Jeong J, Lee J, Shin T, Lee NH, Hyun JW. 2010. Phloroglucinol (1,3,5-trihydroxybenzene) protects against ionizing radiation-induced cell damage through inhibition of oxidative stress *in vitro* and *in vivo*. *Chem Biol Interact* 185: 215-226.
- Quéguineur B, Goya L, Ramos S, Martín MA, Mateos R, Bravo L. 2012. Phloroglucinol: antioxidant properties and effects on cellular oxidative markers in human HepG2 cell line. *Food Chem Toxicol* 50: 2886-2893.
- Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. 1982. Analysis of nitrate, nitrite, and [¹⁵N]nitrate in biological fluids. *Anal Biochem* 126: 131-138.
- Sreejayan, Rao MN. 1997. Nitric oxide scavenging by curcuminoids. *J Pharm Pharmacol* 49: 105-107.
- Ewing JF, Janero DR. 1995. Microplate superoxide dismutase assay employing a nonenzymatic superoxide generator. *Anal Biochem* 232: 243-248.
- Gutteridge JM. 1987. Ferrous-salt-promoted damage to deoxyribose and benzoate. The increased effectiveness of hydroxyl-radical scavengers in the presence of EDTA. *Biochem J* 243: 709-714.
- Yokozawa T, Cho EJ, Hara Y, Kitani K. 2000. Antioxidative activity of green tea treated with radical initiator 2,2'-azobis (2-amidinopropane) dihydrochloride. *J Agric Food Chem* 48: 5068-5073.
- Yokozawa T, Satoh A, Cho EJ, Kashiwada Y, Ikeshiro Y. 2005. Protective role of Coptidis Rhizoma alkaloids against peroxynitrite-induced damage to renal tubular epithelial cells. *J Pharm Pharmacol* 57: 367-374.
- Mosmann T. 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65: 55-63.
- Yagi K. 1976. A simple fluorometric assay for lipoperoxide in blood plasma. *Biochem Med* 15: 212-216.
- Yokode M, Kita T, Kikawa Y, Ogorochi T, Narumiya S, Kawai C. 1988. Stimulated arachidonate metabolism during foam cell transformation of mouse peritoneal macrophages with oxidized low density lipoprotein. *J Clin Invest* 81: 720-729.
- Ragan MA, Glombitza KW. 1986. Phlorotannins, brown algal polyphenols. *Prog Phycol Res* 4: 129-141.
- Kang KA, Lee KH, Chae S, Zhang R, Jung MS, Ham YM, Baik JS, Lee NH, Hyun JW. 2006. Cytoprotective effect of phloroglucinol on oxidative stress induced cell damage via catalase activation. *J Cell Biochem* 97: 609-620.
- Ishii R, Horie M, Saito K, Arisawa M, Kitanaka S. 2001. Inhibitory effects of phloroglucinol derivatives from *Mallotus japonicus* on nitric oxide production by a murine macrophage-like cell line, RAW 264.7, activated by lipopolysaccharide and interferon- γ . *Biochim Biophys Acta* 1568: 74-82.
- Halliwell B. 1987. Oxidants and human disease: some new concepts. *FASEB J* 1: 358-364.
- Halliwell B, Zhao K, Whiteman M. 1999. Nitric oxide and

- peroxynitrite. The ugly, the uglier and the not so good: a personal view of recent controversies. *Free Radic Res* 31: 651-669.
32. Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA. 1990. Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci USA* 87: 1620-1624.
 33. Patel RP, McAndrew J, Sellak H, White CR, Jo H, Freeman BA, Darley-Usmar VM. 1999. Biological aspects of reactive nitrogen species. *Biochim Biophys Acta* 1411: 385-400.
 34. Blough NV, Zafriou OC. 1985. Reaction of superoxide with nitric oxide to form peroxynitrite in alkaline aqueous solution. *Inorg Chem* 24: 3502-3504.
 35. Terao K, Niki E. 1986. Damage to biological tissues induced by radical initiator 2,2'-azobis(2-amidinopropane) dihydrochloride and its inhibition by chain-breaking antioxidants. *Free Radic Biol Med* 2: 193-201.
 36. Miki M, Tamai H, Mino M, Yamamoto Y, Niki E. 1987. Free-radical chain oxidation of rat red blood cells by molecular oxygen and its inhibition by α -tocopherol. *Arch Biochem Biophys* 258: 373-380.
 37. Knight JA. 2000. The biochemistry of aging. *Adv Clin Chem* 35: 1-62.
 38. Oliveira BF, Nogueira-Machado JA, Chaves MM. 2010. The role of oxidative stress in the aging process. *Sci World J* 10: 1121-1128.
 39. Fripiat C, Chen QM, Zdanov S, Magalhaes JP, Remacle J, Toussaint O. 2001. Subcytotoxic H₂O₂ stress triggers a release of transforming growth factor- β 1, which induces biomarkers of cellular senescence of human diploid fibroblasts. *J Biol Chem* 276: 2531-2537.
 40. Fripiat C, Dewelle J, Remacle J, Toussaint O. 2002. Signal transduction in H₂O₂-induced senescence-like phenotype in human diploid fibroblasts. *Free Radic Biol Med* 33: 1334-1346.
 41. Chen QM. 2000. Replicative senescence and oxidant-induced premature senescence: beyond the control of cell cycle checkpoints. *Ann NY Acad Sci* 908: 111-125.
 42. Toussaint O, Medrano EE, von Zglinicki T. 2000. Cellular and molecular mechanisms of stress-induced premature senescence (SIPS) of human diploid fibroblasts and melanocytes. *Exp Gerontol* 35: 927-945.
 43. Morlière P, Santus R. 1998. Pro-oxidant role of superoxide dismutase in ultraviolet-A-induced lipid peroxidation in cultured normal human skin fibroblasts. *Eur J Biochem* 256: 184-189.